

GOAL[®] 2E

HERBICIDE

WARNING

KEEP OUT OF REACH OF CHILDREN

Read and follow all directions, precautions, and restrictions on the label. Do not use if directions, precautions, and restrictions are not followed. Do not use if directions, precautions, and restrictions are not followed.

ACTIVE INGREDIENT


glyphosate (isopropylamine salt) 41.0%
polyvinylpyrrolidone 0.5%
inert ingredients 58.5%

INERT INGREDIENTS

polyvinylpyrrolidone 0.5%
inert ingredients 58.5%

NET CONTENTS

5 GAL
(18.9 L)



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Caution the user and children to not get on or near the product. If contact occurs, wash immediately with soap and water. If contact occurs with eyes, wash with water for 15 minutes. Call a physician in case of eye contact. Remove contaminated clothing and wash skin immediately with soap and water.

This material may be harmful if swallowed. Do not induce vomiting. Call a physician. Do not induce vomiting or give anything by mouth to an unconscious person.

ENVIRONMENTAL HAZARDS

Keep out of streams, ponds, or streams. Do not contaminate water by cleaning of equipment or disposal of waste.

The user should not use this product in areas where it is prohibited by law. Do not apply where prohibited by law. Do not apply where prohibited by law. Do not apply where prohibited by law. Do not apply where prohibited by law.

PHYSICAL OR CHEMICAL HAZARDS

FLAMMABLE.

Keep away from heat and open flame.

DIRECTIONS FOR USE

General classification

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION

Preemergence and Postemergence Weed Control in Conifer Seedbeds

DOAL 15 is a selective herbicide containing 2,4-D and picloram. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds.

Preemergence and Postemergence Weed Control in Conifer Seedbeds

DOAL 15 can be used as a selective herbicide for the preemergence and postemergence control of certain annual grass and broadleaf weeds in conifer seedbeds, nurseries and outplantings.

SEEDBEDS

CONIFER SPECIES

DOAL 15 may be applied to conifer seedbeds of numerous species including the following:

Colorado Blue	Pinus baltica
Jack pine	Pinus edulis
Loblolly pine	Pinus juliflora
Shortleaf pine	Pinus echinata
Eastern white pine	Pinus strobus
Virginia pine	Pinus virginiana
Ponderosa pine	Pinus ponderosa
Lodgepole pine	Pinus contorta
Douglas fir	Pseudotsuga menziesii
Turbo blue spruce	Abies balsamea

Conifers are tolerant of preemergence and postemergence applications of DOAL 15. DOAL 15 applied postemergence will provide both postemergence and residual preemergence control of many broadleaf weeds and grasses.

PREEMERGENCE USE

WEEDS CONTROLLED

When DOAL 15 is applied as a preemergence herbicide, it will control the following weeds and grasses:

Barnyardgrass	Echinochloa crusgalli
Common foxtail	Setaria viridis
Goosegrass	Eleusine indica
Large crabgrass	Digitaria sanguinalis
Proso millet	Panicum capillare

Emergent Weeds Controlled by DOAL 15

Large crabgrass	Digitaria sanguinalis
Common foxtail	Setaria viridis
Common lambsquarters	Chenopodium album
Common Purslane	Portulaca oleracea
Common hemp dogwood	Tripsacum daniellii
Pennsylvania smartweed	Polypogon monspeliensis
Proso millet	Digitaria sanguinalis
Proso millet	Digitaria sanguinalis

DOSAGE

DOAL 15 is a selective herbicide. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds.

TIMING AND METHOD OF APPLICATION

DOAL 15 is a selective herbicide. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds. It is used for the control of weeds in conifer seedbeds.

7. *How do you feel about the way the company is doing?*

DOSEAGE

100% of birds 105 to 150 pounds in weight in 1964. The average total weight of the application was 1.33 pounds per bird. The average weight of the birds was 125 pounds.

[illegible]

NOT FOR CONIFER RELEASE IN FOREST MANAGEMENT PROGRAMS OR FOR FOREST
REGENERATION APPLICATIONS.

Goal 15 may be applied to many conifer species in both interiors and transplants. The following species have been shown to be resistant to Goal 1. It is suggested that other species be tested in infested areas prior to using Goal for pre and post-transference seed control.

Demand interviewee, to see officer involved in incident, long time,
New Orleans service, longest service, was a 1st time, longer than
Table "N, Norway Service and Chicago Service

deeds company, ac-

The following needs are considered when JAL is used as a reference in consequence of the hypotheses 12-18:

[illegible]

THESE ARE THE FIRST EFFECTIVE MEASURES TO BE TAKEN IN ORDER TO
IMPROVE THE SITUATION OF THE PEOPLE WHO ARE BEING TREATED AS
AND WITH NO OTHER REASON FOR THEIR PRESENCE.

includes both the ... and ...

Annual Normality
Redspot Pigweed
Scarlet Pigweed
Wayweed Dog Fen
el. Mustard
Shepherd's Purse
Lemon Thistle
Black Chickweed
Common Amaranth
Buttercup
Climbing
Wild Buckwheat
Blackberry
Prostrate Knotweed
Laysanweed
Red Sorrel
Common Immortelle
Marsh Marigold
Annual
Common
Red Clover
Blackberry
Common
Red Clover
Blackberry
Common

1994-1995 1995-1996 1996-1997 1997-1998 1998-1999 1999-2000 2000-2001 2001-2002 2002-2003 2003-2004 2004-2005 2005-2006 2006-2007 2007-2008 2008-2009 2009-2010 2010-2011 2011-2012 2012-2013 2013-2014 2014-2015 2015-2016 2016-2017 2017-2018 2018-2019 2019-2020 2020-2021 2021-2022 2022-2023 2023-2024 2024-2025 2025-2026 2026-2027 2027-2028 2028-2029 2029-2030 2030-2031 2031-2032 2032-2033 2033-2034 2034-2035 2035-2036 2036-2037 2037-2038 2038-2039 2039-2040 2040-2041 2041-2042 2042-2043 2043-2044 2044-2045 2045-2046 2046-2047 2047-2048 2048-2049 2049-2050 2050-2051 2051-2052 2052-2053 2053-2054 2054-2055 2055-2056 2056-2057 2057-2058 2058-2059 2059-2060 2060-2061 2061-2062 2062-2063 2063-2064 2064-2065 2065-2066 2066-2067 2067-2068 2068-2069 2069-2070 2070-2071 2071-2072 2072-2073 2073-2074 2074-2075 2075-2076 2076-2077 2077-2078 2078-2079 2079-2080 2080-2081 2081-2082 2082-2083 2083-2084 2084-2085 2085-2086 2086-2087 2087-2088 2088-2089 2089-2090 2090-2091 2091-2092 2092-2093 2093-2094 2094-2095 2095-2096 2096-2097 2097-2098 2098-2099 2099-2100 2100-2101 2101-2102 2102-2103 2103-2104 2104-2105 2105-2106 2106-2107 2107-2108 2108-2109 2109-2110 2110-2111 2111-2112 2112-2113 2113-2114 2114-2115 2115-2116 2116-2117 2117-2118 2118-2119 2119-2120 2120-2121 2121-2122 2122-2123 2123-2124 2124-2125 2125-2126 2126-2127 2127-2128 2128-2129 2129-2130 2130-2131 2131-2132 2132-2133 2133-2134 2134-2135 2135-2136 2136-2137 2137-2138 2138-2139 2139-2140 2140-2141 2141-2142 2142-2143 2143-2144 2144-2145 2145-2146 2146-2147 2147-2148 2148-2149 2149-2150 2150-2151 2151-2152 2152-2153 2153-2154 2154-2155 2155-2156 2156-2157 2157-2158 2158-2159 2159-2160 2160-2161 2161-2162 2162-2163 2163-2164 2164-2165 2165-2166 2166-2167 2167-2168 2168-2169 2169-2170 2170-2171 2171-2172 2172-2173 2173-2174 2174-2175 2175-2176 2176-2177 2177-2178 2178-2179 2179-2180 2180-2181 2181-2182 2182-2183 2183-2184 2184-2185 2185-2186 2186-2187 2187-2188 2188-2189 2189-2190 2190-2191 2191-2192 2192-2193 2193-2194 2194-2195 2195-2196 2196-2197 2197-2198 2198-2199 2199-2200 2200-2201 2201-2202 2202-2203 2203-2204 2204-2205 2205-2206 2206-2207 2207-2208 2208-2209 2209-2210 2210-2211 2211-2212 2212-2213 2213-2214 2214-2215 2215-2216 2216-2217 2217-2218 2218-2219 2219-2220 2220-2221 2221-2222 2222-2223 2223-2224 2224-2225 2225-2226 2226-2227 2227-2228 2228-2229 2229-2230 2230-2231 2231-2232 2232-2233 2233-2234 2234-2235 2235-2236 2236-2237 2237-2238 2238-2239 2239-2240 2240-2241 2241-2242 2242-2243 2243-2244 2244-2245 2245-2246 2246-2247 2247-2248 2248-2249 2249-2250 2250-2251 2251-2252 2252-2253 2253-2254 2254-2255 2255-2256 2256-2257 2257-2258 2258-2259 2259-2260 2260-2261 2261-2262 2262-2263 2263-2264 2264-2265 2265-2266 2266-2267 2267-2268 2268-2269 2269-2270 2270-2271 2271-2272 2272-2273 2273-2274 2274-2275 2275-2276 2276-2277 2277-2278 2278-2279 2279-2280 2280-2281 2281-2282 2282-2283 2283-2284 2284-2285 2285-2286 2286-2287 2287-2288 2288-2289 2289-2290 2290-2291 2291-2292 2292-2293 2293-2294 2294-2295 2295-2296 2296-2297 2297-2298 2298-2299 2299-2300 2300-2301 2301-2302 2302-2303 2303-2304 2304-2305 2305-2306 2306-2307 2307-2308 2308-2309 2309-2310 2310-2311 2311-2312 2312-2313 2313-2314 2314-2315 2315-2316 2316-2317 2317-2318 2318-2319 2319-2320 2320-2321 2321-2322 2322-2323 2323-2324 2324-2325 2325-2326 2326-2327 2327-2328 2328-2329 2329-2330 2330-2331 2331-2332 2332-2333 2333-2334 2334-2335 2335-2336 2336-2337 2337-2338 2338-2339 2339-2340 2340-2341 2341-2342 2342-2343 2343-2344 2344-2345 2345-2346 2346-2347 2347-2348 2348-2349 2349-2350 2350-2351 2351-2352 2352-2353 2353-2354 2354-2355 2355-2356 2356-2357 2357-2358 2358-2359 2359-2360 2360-2361 2361-2362 2362-2363 2363-2364 2364-2365 2365-2366 2366-2367 2367-2368 2368-2369 2369-2370 2370-2371 2371-2372 2372-2373 2373-2374 2374-2375 2375-2376 2376-2377 2377-2378 2378-2379 2379-2380 2380-2381 2381-2382 2382-2383 2383-2384 2384-2385 2385-2386 2386-2387 2387-2388 2388-2389 2389-2390 2390-2391 2391-2392 2392-2393 2393-2394 2394-2395 2395-2396 2396-2397 2397-2398 2398-2399 2399-2400 2400-2401 2401-2402 2402-2403 2403

1

For persistence or post-persistence seed control apply 2 to 4 quarts of 1000 pounds active ingredient of 20AL per acre.

For optimum weed control, preemergence applications should be made immediately after transplanting and postemergence applications should be made to weeds less than 4 inches in height. Two applications may be necessary, in 3" transplanted corn fields, for season long weed control. The addition of 0.25% (1 qt./100 gal.) of sorby solution of Tordon AG-38 enhances GOAL 15 activity on emerged weeds. GOAL 15 must be applied only to confirm transplants which are dormant. GOAL 15 should be thoroughly mixed with clear water at recommended concentrations and applied at a minimum of 20 gallons of water per treated acre. Apply at 20 to 40 psi of pressure. Sorby GOAL 15 over the top of transplants and allow 24 hours before initiating heavy rainfall immediately following application to emerged weeds may reduce effectiveness.

PROHIBITIONS Do not contaminate water; do not dispose of chemicals in drains or on land; do not mix contents. Do not reuse empty containers.

PESTICIDE DISPOSAL: Pesticide spray materials should not be poured down drains, into storm sewers, or into water bodies. Pesticide containers should be triple rinsed and then disposed of in a secure container. Do not burn or incinerate pesticides. Do not place away from water supplies.

CONTAINER DISPOSAL. Do not use or dispose of any container that has not been approved for pesticide containers at your local waste place.

GENERAL. Consult Federal State or Local Social Security Administration approved state health care
 bureaus such as United Open Burning

[illegible]

222