

ROHM AND HAAS COMPANYINDEPENDENCE MALL WEST
PHILADELPHIA, PENNSYLVANIA 19105EPA REG. NO. 707-145-AA
EPA EST. NO. 707-PA-1SUPPLEMENTAL LABELING FOR GOAL[®] 2E HERBICIDE USE ON FALLOW LAND
(FOR USE ONLY IN IDAHO, OREGON AND WASHINGTON)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Precautions and Storage and Disposal Instructions on back panel and attached use directions. If the conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

GENERAL INFORMATION

GOAL 2E is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with Roundup for the control of certain annual broadleaf weeds in a fallow land system. GOAL 2E herbicide can be used as an effective tool to reduce weed growth prior to the establishment of a dry soil mulch. Use of this product is restricted to summer fallow land which will be planted back the following year to winter wheat, barley or oats.

GOAL 2E HERBICIDE USED ALONE

DOSAGE

GOAL 2E herbicide should be used at 0.5 to 2.0 pints (0.13 to 0.5 lb. active) per broadcast acre.

WEEDS CONTROLLED

GOAL 2E herbicide will provide postemergence control and preemergence activity of the following broadleaf weeds when used at recommended dosages.

FIDDLENECK, COAST	Amsinckia intermedia
HENBIT	Lamium amplexicaule
LETTUCE, PRICKLY (China lettuce)	Lactuca serriola
MUSTARD, BLUE	Chorispora tenella
MUSTARD, TUMBLE (Jim Hill mustard)	Sisymbrium altissimum
PIGWEEED, REDROOT	Amaranthus retroflexus
PURSLANE, COMMON	Portulaca oleraceae
SHEPHERDSPURSE	Capsella bursa-pastoris
SOWTHISTLE, ANNUAL	Sonchus oleraceus

TIMING AND METHOD OF APPLICATION

The most effective postemergence weed control is achieved when GOAL 2E herbicide is applied to seedling weeds (less than 4 inches in height). Seedling weeds are controlled as they come in contact with the soil applied herbicide during emergence.

GOAL 2E herbicide should be applied in 20 to 40 gallons of water per acre depending upon density of emerged weeds. Use a low pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

MIXING DIRECTIONS

Fill the spray tank at least one-third full of clean water and add the recommended amount of GOAL 2E herbicide while the pump and agitator are running. Complete filling of the spray tank with water and then add 1 quart of TRITON AG-98 or comparable 80% active nonionic surfactant cleared for use on growing crops, per 100 gallons of spray. Maintain agitation until spraying is complete.

GOAL 2E HERBICIDE - TANK MIXES

DOSAGE

For postemergence control of annual grassy weeds, GOAL 2E herbicide can be tank mixed with Roundup^R. Tank mix 0.5 to 2.0 pints (0.13 to 0.5 lb. active) of GOAL 2E herbicide with 0.75 to 1 pint (0.38 to 0.5 lb. active) of Roundup for each acre treated. Refer to the "Fallow and Reduced Tillage System" Section on the Roundup label for specific use directions and restrictions.

Fill the spray tank at least one-third full of clean water and add the recommended amounts of GOAL 2E herbicide and Roundup while the pump and agitator are running. Complete filling of the spray tank with water. Add 1 quart of TRITON AG-98 or comparable 80% active nonionic surfactant cleared for use on growing crops, per 100 gallons of spray. Maintain agitation until spraying is complete.

FALLOW LAND - GENERAL USE RESTRICTIONS

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not contaminate irrigation water or water used for domestic purposes.
- Do not use any plants treated with GOAL 2E herbicide for feed or forage.
- Do not feed or graze animals on any areas treated with GOAL 2E herbicide.
- GOAL 2E herbicide should be applied only by ground application equipment.
- Do not apply when weather conditions favor drift. Avoid drift to all non-target areas. GOAL 2E herbicide is phytotoxic to plant foliage.
- Thoroughly flush spray equipment (tank, pump, hoses and boom) with clean water before and after each use. Residual GOAL 2E herbicide remaining in spray equipment may damage other crops. To assist removal of GOAL herbicide residues in spray equipment, TRITON AG-98 or TRITON CS-7 may be added at the rate of 1 quart per 100 gallons of water during flushing.

- Do not rotate any crops other than cotton, soybeans, or spearmint/peppermint within a 10-month period after treatment.
- Use GOAL 2E herbicide only for recommended purposes and at recommended rates.
- Do not treat ditch banks or waterways with GOAL 2E herbicide.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

JAH:flp
(8095E/49Z)
12/16/83

ROHM AND HAAS COMPANYINDEPENDENCE MALL WEST
PHILADELPHIA PENNSYLVANIA 19103EPA REG. NO. 707-145-AA
EPA EST. NO. 707-PA-1SUPPLEMENTAL LABELING FOR GOAL^R 2E HERBICIDE USE ON FALLOW LAND
(FOR USE ONLY IN IDAHO, OREGON AND WASHINGTON)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Precautions and Storage and Disposal Instructions on back panel and attached use directions. If the conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

GENERAL INFORMATION

GOAL 2E is effective as a preemergence and/or postemergence herbicide when used alone or in a tank mix combination with Roundup for the control of certain annual broadleaf weeds in a fallow land system. GOAL 2E herbicide can be used as an effective tool to reduce weed growth prior to the establishment of a dry soil mulch. Use of this product is restricted to summer fallow land which will be planted back the following year to winter wheat, barley or oats.

GOAL 2E HERBICIDE USED ALONE

DOSAGE

GOAL 2E herbicide should be used at 0.13 to 0.5 lb. active per broadcast acre.


WEEDS CONTROLLED

GOAL 2E herbicide will provide postemergence control and preemergence activity of the following broadleaf weeds when used at recommended dosages.

FIDDLENECK, COAST	Amsinckia intermedia
HENBIT	Lamium amplexicaule
LETTUCE, PRICKLY (China lettuce)	Lactuca serriola
MUSTARD, BLUE	Chorispora tenella
MUSTARD, TUMBLE (Jim Hill mustard)	Sisymbrium altissimum
PIGWEEED, REDROOT	Amaranthus retroflexus
PURSLANE, COMMON	Portulaca oleraceae
SHEPHERDSPURSE	Capsella bursa-pastoris
SOWTHISTLE, ANNUAL	Sonchus oleraceus

TIMING AND METHOD OF APPLICATION

The most effective postemergence weed control is achieved when GOAL 2E herbicide is applied to seedling weeds (less than 4 inches in height). Seedling weeds are controlled as they come in contact with the soil applied herbicide during emergence.



757-145

GOAL 2E herbicide should be applied in 20 to 40 gallons of water per acre depending upon density of emerged weeds. Use a low pressure sprayer equipped with flat fan nozzles. Spray equipment should be calibrated carefully before each use.

MIXING DIRECTIONS

Fill the spray tank at least one-third full of clean water and add the recommended amount of GOAL 2E herbicide while the pump and agitator are running. Complete filling of the spray tank with water and then add 1 quart of TRITON AG-98 or comparable 80% active nonionic surfactant cleared for use on growing crops, per 100 gallons of spray. Maintain agitation until spraying is complete.

GOAL 2E HERBICIDE - TANK MIXES

1,2,4 3.6

DOSAGE

For postemergence control of annual grassy weeds, GOAL 2E herbicide can be tank mixed with Roundup^R. Tank mix 0.5 to 2.0 pints (0.13 to 0.5 lb. active) of GOAL 2E herbicide with 0.75 to 1 pint (0.38 to 0.5 lb. active) of Roundup for each acre treated. Refer to the "Fallow and Reduced Tillage System" Section on the Roundup label for specific use directions and restrictions.

Fill the spray tank at least one-third full of clean water and add the recommended amounts of GOAL 2E herbicide and Roundup while the pump and agitator are running. Complete filling of the spray tank with water. Add 1 quart of TRITON AG-98 or comparable 80% active nonionic surfactant cleared for use on growing crops, per 100 gallons of spray. Maintain agitation until spraying is complete.

FALLOW LAND - GENERAL USE RESTRICTIONS

- Read and observe all label directions before using. When tank mixing, always read all individual manufacturers' labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.
- Do not contaminate irrigation water or water used for domestic purposes.
- Do not use any plants treated with GOAL 2E herbicide for feed or forage.
- Do not feed or graze animals on any areas treated with GOAL 2E herbicide.
- GOAL 2E herbicide should be applied only by ground application equipment.
- Do not apply when weather conditions favor drift. Avoid drift to all non-target areas. GOAL 2E herbicide is phytotoxic to plant foliage.
- Thoroughly flush spray equipment (tank, pump, hoses and boom) with clean water before and after each use. Residual GOAL 2E herbicide remaining in spray equipment may damage other crops. To assist removal of GOAL herbicide residues in spray equipment, TRITON AG-98 or TRITON CS-7 may be added at the rate of 1 quart per 100 gallons of water during flushing.

- Do not rotate any crops other than cotton, soybeans, or spearmint/peppermint within a 10-month period after treatment.
- Use GOAL 2E herbicide only for recommended purposes and at recommended rates.
- Do not treat ditch banks or waterways with GOAL 2E herbicide.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrant that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

JAH:flp
(8095E/49Z)
12/16/83