

## ROHM AND HAAS COMPANY

INDEPENDENCE MALL WEST  
PHILADELPHIA, PENNSYLVANIA 19105



### SUPPLEMENTAL LABELING

BLAZER<sup>®</sup> 2L herbicide and Fusilade<sup>®</sup> herbicide for postemergence control of broadleaf and grass weeds

BLAZER 2L (EPA Reg. No. 707-149)  
Fusilade (EPA Reg. No. 10182-67)

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Storage and Disposal Instruction on container label. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within 30 days of purchase to the place of purchase.

### POSTEMERGENCE CONTROL OF GRASS AND BROADLEAF WEEDS

BLAZER 2L herbicide and Fusilade 4E may be used together in a postemergence program for broad spectrum weed control in soybeans.

BLAZER 2L is a selective postemergence herbicide for the control of certain broadleaf weeds. Fusilade 4E is a selective postemergence herbicide for control of annual and perennial grass weeds in soybeans. BLAZER 2L herbicide and Fusilade 4E may be applied sequentially or in a tank mix. The growth stage of weeds at the time of application will determine which method of application will result in the most satisfactory results. Both BLAZER 2L herbicide and Fusilade 4E should be applied to actively growing weeds. See BLAZER 2L label for defined environmental conditions that promote active growth. Do not apply either herbicide if weeds appear stressed due to unfavorable temperatures, drought and/or low soil fertility. Do not apply BLAZER 2L or BLAZER 2L/Fusilade 4E tank mix if soybeans show injury from prior herbicide applications. Read and observe all applicable label directions and limitations for both BLAZER 2L herbicide and Fusilade 4E before using.

### METHOD 1: Sequential Application Fusilade 4E Followed by BLAZER 2L Herbicide

(Annual and/or perennial grass weeds at proper growth stage for treatment, prior to broadleaf weed treatment)

Apply Fusilade 4E and an approved adjuvant at the recommended rate and growth stage for the grass weeds being treated, as per the Fusilade 4E label directions. Fusilade 4E is a systemic herbicide, therefore, when treating annual grasses, allow at least 3 days, and when treating perennial grasses allow at least 5 days to transpire prior to a BLAZER 2L herbicide application. This will enable Fusilade 4E to be adequately translocated to the meristematic regions of the plant. After the appropriate time interval has elapsed, apply BLAZER 2L herbicide and TRITON AG-9S surfactant for susceptible broadleaf weeds as outlined on the BLAZER 2L herbicide label.

METHOD 2: Sequential Application - BLAZER 2L Herbicide Followed by Fusilade 4E

(Broadleaf weed, at proper growth stage for treatment, prior to annual and/or perennial grass weed treatment)

Apply BLAZER 2L herbicide with TRITON AG-90 surfactant at the recommended rate and growth stage for susceptible broadleaf weeds, as per the BLAZER 2L herbicide label directions. Broadleaf herbicides can stress and affect the growth of grass species. Fusilade 4E must be applied to actively growing grasses for best results. A sequential application of Fusilade 4E may be applied following a BLAZER 2L herbicide application when grasses resume active growth with the development of a new leaf. Follow the recommended rates and growth stage for the grass weeds being treated, as per the Fusilade 4E label.

METHOD 3: Tank Mix Applications of BLAZER 2L Herbicide and Fusilade 4E

(Only annual grasses and broadleaf weeds are present and at the proper growth stage for treatment, as per the respective labels)

A tank mix of BLAZER 2L herbicide and Fusilade 4E may be applied at the recommended rates and growth stages to susceptible broadleaf weeds and annual grass species in a manner as described on the respective labels. Use an approved nonionic surfactant, as per the Fusilade 4E label recommendations. Follow water volume and spray pressure recommendations, as per the BLAZER 2L herbicide label. Load the spray tank with half the amount of required water and add the recommended amounts of Fusilade 4E, BLAZER 2L, and nonionic surfactant while the agitator is running. Add the remaining quantity of water. Do not make more than one application of this tank mix per season.

NOTE: Tank mix applications sometimes have resulted in reduced grass weed control and possible increases in crop injury as compared to either product used alone. If grass regrowth occurs following an application of the tank mix or an additional flush of grasses emerge, make a second application of Fusilade 4E to actively growing annual grass weeds, as per the label recommendations. A tank mix application is not recommended if perennial grass weeds are the predominant grasses to be controlled.

POSTEMERGENCE USE RESTRICTIONS (PARTIAL LIST)

- . Always read and follow the restrictions and limitations for all products whether used alone or in a tank mix. The most restrictive labeling applies in tank mixtures.
- . Do not apply a total of more than 1 pint of Fusilade 4E per acre per season to soybeans.
- . Do not apply more than 4 pints (1.0 lb. active) per acre of BLAZER 2L herbicide per growing season for soybeans.
- . Make the last Fusilade 4E application before soybean bloom. Do not apply within 100 days of harvest.

- . Do not apply BLAZER 2L herbicide within 50 days of harvest of soybeans.
- . Do not use treated plants for feed or forage.
- . Do not plant rotational crops other than cotton and soybeans within 60 days after last Fusilade 4E application.
- . Crop Rotation Restriction: Root Crops (such as carrots, turnips, sweet potatoes, etc.) must not be planted in fields treated with BLAZER 2L herbicide for a period of 18 months following treatment.
- . In the case of crop failure, only soybeans may be immediately replanted.
- . Avoid drift to all other crops and non-target areas.
- . Thoroughly flush spray tanks with water before and after each use.
- . BLAZER 2L herbicide requires a six hour rain free period for best results. Do not apply BLAZER 2L herbicide or BLAZER 2L/Fusilade tank mix if rain is threatening.
- . Do not make more than one application of BLAZER 2L/Fusilade 4E herbicide tank mix in a single season.
- . Do not use during dry weather conditions when crop and weeds are under stress or wilt and not actively growing, as weed control may be reduced.
- . Do not apply BLAZER 2L herbicide or BLAZER 2L/Fusilade 4E herbicide tank mix when crop and weeds are under stress from cold temperature, when maximum day temperature is below 70°F, or when soil temperature is below 60°F, as weeds will not be actively growing and control may be reduced.
- . Do not apply BLAZER 2L or BLAZER 2L/Fusilade 4E tank mix when crops are exhibiting injury from previously applied pesticides or are exhibiting stress symptoms from disease, nematodes or insects, as increased crop response may result.
- . Do not apply BLAZER 2L or BLAZER 2L/Fusilade 4E tank mix when crop and weeds are under stress from excessive fertilizer or soil salt, wind injury, frost damage, or high temperature stress or wilt, as increased crop response may result.
- . Do not mix BLAZER 2L, or BLAZER 2L/Fusilade 4E tank mix with other pesticides or liquid fertilizers except as specified on this label or on other approved Rohm and Haas Company supplemental labeling.

WRU:fs  
(Doc. #8683E/217  
2/20/84