

**ROHM AND HAAS COMPANY**

INDEPENDENCE MALL WEST  
PHILADELPHIA, PENNSYLVANIA 19105



SUPPLEMENTAL LABELING

BLAZER<sup>R</sup> 2S herbicide with Poast<sup>(TM)</sup> herbicide

FOR POSTEMERGENCE BROADLEAF AND GRASS WEED CONTROL IN SOYBEANS

BLAZER 2S Herbicide (EPA Reg. No. 707-150)

Poast (EPA Reg. No. 7969-53)

DIRECTIONS FOR USE

General Classification.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, and Storage and Disposal Instructions on container label. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

TANK MIXTURE OF BLAZER 2S HERBICIDE  
WITH POAST HERBICIDE FOR CONTROL OF  
ANNUAL BROADLEAF/GRASS WEEDS IN SOYBEANS

BLAZER 2S herbicide at 1.5 to 2.0 pints (0.38 to 0.5 lb. active) per acre may be tank mixed with Poast herbicide at 1.5 pints for postemergence control of selected annual broadleaf/grass weeds in soybeans. In order to determine the correct application rate of BLAZER 2S herbicide to use in the tank mixture, see the BLAZER 2S herbicide use rate table.

For grass species and timing refer to Poast label.

Always add Poast herbicide in the tank mixture at 1.5 pints per acre with 2 pints per acre of a recommended nonphytotoxic oil concentrate. Use 20 gallons of water per acre with a minimum of 40 psi pressure at the nozzle. Air applications should be made in 10 gallons of water per acre. See your Poast supplier for a list of oil concentrates recommended by BASF Wyandotte Corporation. Oil concentrate must be used with the tank mixture in place of TRITON AG-98 surfactant or any other surfactant.

#### POAST HERBICIDE IN SEQUENTIAL APPLICATIONS WITH BLAZER 2S HERBICIDE

When using BLAZER 2S herbicide as a separate application be sure and allow a minimum of 7 days between the BLAZER 2S herbicide and the Poast grass herbicide application. This applies when BLAZER 2S herbicide is applied first. When Poast herbicide is applied first, wait 24 hours before applying BLAZER 2S herbicide. It is important that treated grass has resumed active growth before spraying with Poast grass herbicide. This waiting period is important in achieving maximum activity with the Poast grass herbicide. For additional information on rate, timing and use restrictions for Poast herbicide refer to the Poast label.

#### POSTEMERGENCE USE RESTRICTIONS (PARTIAL LIST)

- Always read and follow the restrictions and limitations for all products when used alone or in a tank mix. The most restrictive labeling must apply in tank mixtures.
- Do not apply BLAZER 2S herbicide or Poast herbicide when soybeans are exhibiting injury from previously applied pesticides or are exhibiting stress symptoms from disease, nematodes or insects, as increased crop response may result.
- Do not apply BLAZER 2S within 50 days of harvest and do not apply Poast within 90 days of harvest.
- Do not graze treated soybean fields and do not feed treated soybean forage, ensilage or hay to livestock (see BLAZER 2S herbicide and Poast herbicide labels).
- Do not apply BLAZER 2S herbicide or Poast herbicide when soybeans and weeds are under stress from excessive fertilizer or soil salts, wind injury, frost damage or high temperature stress or wilt, as increased crop response and/or decreased weed control may result.

BLAZER<sup>R</sup> is a Registered Trademark of Rohm and Haas Co.

Poast<sup>TM</sup> is a Trademark of BASF Wyandotte Corporation

#### CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use.

ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

(8209E/85Z)  
1/3/84

**ROHM AND HAAS COMPANY**

INDEPENDENCE MALL WEST  
PHILADELPHIA, PENNSYLVANIA 19106



11/17/84  
707-150-1  
SUPPLEMENTAL LABELING

BLAZER<sup>R</sup> 2S herbicide and Basagran<sup>R</sup> herbicide with Poast<sup>(TM)</sup> herbicide.

FOR POSTEMERGENCE BROADLEAF AND GRASS WEED CONTROL IN SOYBEANS

BLAZER 2S Herbicide (EPA Reg. No. 707-150)

Basagran (EPA Reg. No. 7969-45) Poast (EPA Reg. No. 7969-53) 54

DIRECTIONS FOR USE

General Classification.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, and Storage and Disposal Instructions on container label. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

TANK MIXTURE OF BLAZER 2S HERBICIDE  
WITH BASAGRAN AND POAST HERBICIDES FOR  
CONTROL OF ANNUAL BROADLEAF/GRASS WEEDS IN SOYBEANS

BLAZER 2S herbicide at 1.5 to 2.0 pints (0.38 to 0.5 lb. active) per acre may be tank mixed with Basagran herbicide at 1.0 to 1.5 pints (0.5 to 0.75 lb. active) per acre and Poast herbicide at 1.5 pints for postemergence control of selected annual broadleaf/grass weeds in soybeans. In order to determine the correct application rate of BLAZER 2S herbicide to use in the tank mixture, see the BLAZER 2S herbicide use rate table.

For control of cocklebur up to the 6 leaf stage and velvetleaf up to the 4 leaf stage add 1 pint (0.5 lb. active) per acre of Basagran to BLAZER 2S herbicide and Poast herbicide. For prickly sida (teaweed) up to the 6 leaf stage, add 1.5 pints (0.75 lb. active) per acre of Basagran to BLAZER 2S herbicide and Poast herbicide. Basagran may also be used in the tank mixture at rates up to 2 pints (1.0 lb. active) per acre.

For grass species and timing refer to Basagran/Poast label.

Always add Poast herbicide in the tank mixture at 1.5 pints per acre with 2 pints per acre of a recommended nonphytotoxic oil concentrate. Use 20 gallons of water per acre with a minimum of 40 psi pressure at the nozzle. Air applications should be made in 10 gallons of water per acre. See your Poast/Basagran supplier for a list of oil concentrates recommended by BASF Wyandotte Corporation. Oil concentrate must be used with the tank mixture in place of TRITON AG-98 surfactant or any other surfactant.

#### POAST HERBICIDE IN SEQUENTIAL APPLICATIONS

BLAZER ALONE OR AS A TANK MIX OF BLAZER/BASAGRAN HERBICIDE.

When using BLAZER 2S herbicide or BLAZER/Basagran herbicides as a separate application be sure and allow a minimum of 7 days between the BLAZER 2S or BLAZER/Basagran herbicides in the tank mix and the Poast grass herbicide application. This applies when BLAZER 2S herbicide or the tank mix of BLAZER/Basagran herbicides are applied first. When Poast herbicide is applied first, wait 24 hours before applying BLAZER 2S herbicide or a tank mixture of BLAZER/Basagran herbicides. It is important that treated grass has resumed active growth before spraying with Poast grass herbicide. This waiting period is important in achieving maximum activity with the Poast grass herbicide. For additional information on rate, timing and use restrictions for Poast herbicide refer to the Poast label.

#### POSTEMERGENCE USE RESTRICTIONS (PARTIAL LIST)

- Always read and follow the restrictions and limitations for all products when used alone or in a tank mix. The most restrictive labeling must apply in tank mixtures.
- Do not apply BLAZER 2S herbicide or Poast herbicide when soybeans are exhibiting injury from previously applied pesticides or are exhibiting stress symptoms from disease, nematodes or insects, as increased crop response may result.
- Do not apply BLAZER 2S within 50 days of harvest and do not apply Poast within 90 days of harvest.
- Do not graze treated soybean fields and do not feed treated soybean forage, ensilage or hay to livestock (see BLAZER 2S herbicide and Poast herbicide labels).
- Do not apply BLAZER 2S herbicide or Poast herbicide when soybeans and weeds are under stress from excessive fertilizer or soil salts, wind injury, frost damage or high temperature stress or wilt, as increased crop response and/or decreased weed control may result.

BLAZER<sup>R</sup> is a Registered Trademark of Rohm and Haas Co.

Basagran<sup>R</sup> is a Registered Trademark of BASF Wyandotte Corporation

Poast<sup>TM</sup> is a Trademark of BASF Wyandotte Corporation

**CONDITIONS OF SALE AND WARRANTY**

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use.

ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Crop injury, ineffectiveness or other unintended consequences resulting from such factors as failure to follow label directions, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property, handling, storage and manner of use are beyond the control of Rohm and Haas and Seller. All such risks will be assumed by the Buyer or User. In no case will Rohm and Haas or Seller be held liable for consequential, special or indirect damages resulting from the use or handling of this product.

(8210E/85Z)  
1/3/84