

8842-R2a

DITHANE M-45

FLOWABLE

AGRICULTURAL FUNGICIDE

Protective clothing consisting of long pants, long sleeve shirt, gloves, hat and boots must be worn during mixing and loading.

ACTIVE INGREDIENTS

A coordination product of zinc ion and manganese ethylene bisdithiocarbamate 35.0%

In which the ingredients are:

Manganese	7.0%
Zinc	0.9%
Ethylene bisdithiocarbamate ion $(C_2H_4N_2S_4)^{2-}$	27.1%

INERT INGREDIENTS

65.0%
100%

KEEP OUT OF REACH OF CHILDREN


CAUTION

Before using this product, read all additional precautionary statements, storage and disposal instructions, use restrictions and limit of warranty and liability statements on side panel and attached use directions.

EPA REG. NO. 100-100
EPA REG. NO. 100-100
EPA REG. NO. 100-100
EPA REG. NO. 100-100

NET CONTENTS
2.5 GALS.
(9.46 L)

MADE IN U.S.A.



**ROHM
HAAS**

PHILADELPHIA, PA. 19106

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

1. Causes irritation of nose, throat, eyes and skin. Avoid breathing dust. If irritation occurs, stop spraying. If contact with skin or eyes, wash with plenty of water. Get medical attention if irritation persists. Wash thoroughly after using and before eating or drinking. Contaminated clothing before reuse.

ENVIRONMENTAL HAZARD

2. This pesticide is toxic to fish. Keep out of lakes, ponds or streams. When weather conditions favor drift from areas treated, wash by cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL

PROHIBITIONS AND RECOMMENDATIONS: Do not store in areas accessible to children or livestock. Do not dump in waterways.

3. Do not use in areas where it may be harmful to plants, animals or humans. Do not use in areas where it may be harmful to the environment. Do not use in areas where it may be harmful to the environment.

PESTICIDE DISPOSAL: Pesticide spray mixture or residue should be disposed of in accordance with local, state or federal regulations.

CONTAINER DISPOSAL: Empty container should be disposed of in accordance with local, state or federal regulations.

GENERAL: Do not use in areas where it may be harmful to plants, animals or humans. Do not use in areas where it may be harmful to the environment. Do not use in areas where it may be harmful to the environment.

Protective clothing consisting of long pants, long sleeve shirt, gloves, hat and boots must be worn during mixing and loading.

INERT INGREDIENTS	65.0-
	100

EPA Ref. No. 707-156
EPA Est. No. 707-FA-2
U.S. Export Administration
Regulations

MADE IN U S A

2 This pesticide is toxic to fish. ~~Keep out of~~ ^{Do not apply to water} ponds or streams. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

GENERAL COMMENTS: The above information was obtained from a review of the file of the above named individual. The information was obtained from a review of the file of the above named individual. The information was obtained from a review of the file of the above named individual.



ROHM AND
PHILADELPHIA

EPA REG NO
EPA EST NO
Copyright 19

DITHANE

DIRECTIONS

It is a violation of law
to use this product in a
manner inconsistent with
these directions.

GENERAL INFORMATION

Flow DITHANE M 45 FLOWABLE is a fungicide for use on a wide range of crops. It is applied as a spray or dust. For detailed information on use, see the label for the specific product.

VEGETABLES

ASPARAGUS—Rust. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

CARROTS—Alternaria. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

CELERY, FENNEL—Eschschoria. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

CORN (sweet, popcorn)—Blight. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

CUCUMBERS, CANTALOUPE, WATERMELON—Downy mildew. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

ONIONS (dry bulb)—Eschschoria. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

OKRA (New York)—Blight. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

POTATOES—Late blight. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

POTATO SEED TREATMENT—Late blight. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

TOMATOES—Late blight. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

TOMATOES—Bacterial wilt. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

TOMATOES—Bacterial wilt. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

FRUIT—Apples. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

APPLES—Crabapple. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

COTTON—Fungal diseases. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

PEANUTS—Fungal diseases. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

BERMUDAGRASS—Fungal diseases. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

Begin application—10 to 14 days after planting or when disease symptoms first appear. Repeat at 10 to 14 day intervals.

IMPORTANT—Do not use DITHANE M 45 FLOWABLE on crops for which it is not registered. Do not use for seed treatment.

SUGAR BEETS—Fungal diseases. Apply 1.3 to 1.7 qts. per acre at 10 day intervals until plants are in full leaf. Do not use for seed treatment.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

SEED TREATMENT—Do not use DITHANE M 45 FLOWABLE for seed treatment. Use a separate seed treatment product.

STORAGE AND DISPOSAL

PROHIBITIONS AND RECOMMENDATIONS—Do not contaminate water, food, or feed by storage or disposal. Do not use in a way that will cause injury to humans or animals.

CONTAINER DISPOSAL—Triple rinse immediately and thoroughly. Do not reuse for any purpose.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

PESTICIDE DISPOSAL—Pesticide spray mixture or residue that cannot be used for its intended purpose should be disposed of in a safe manner.

CONTAINER DISPOSAL—Triple rinse immediately and thoroughly. Do not reuse for any purpose.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

GENERAL—Consult Federal, State, and local authorities for disposal instructions.

DITHANE is a trademark of Rohm and Haas Company or of its subsidiaries or affiliates.

8842-80a

ROHM AND HAAS COMPANY
PHILADELPHIA PA 19105

EPA REG NO 707 156
EPA EST NO 707 PA 2
Copyright 1979 Rohm and Haas Company

DITHANE® M-45 FLOWABLE FUNGICIDE

ROHM AND HAAS COMPANY
PHILADELPHIA PA 19105

EPA REG NO 707 156
EPA EST NO 707 PA 2
Copyright 1979 Rohm and Haas Company

DITHANE® M-45 FLOWABLE FUNGICIDE

DIRECTIONS FOR USE

Read and follow the label directions for use of this product. Do not use in a manner inconsistent with the label directions.

GENERAL INSTRUCTIONS

Use DITHANE M-45 FLOWABLE fungicide to control a wide range of plant diseases caused by fungi. It is a systemic fungicide which is absorbed by the plant and moves to the site of infection. It is effective against a wide range of fungi including Ascomycetes, Basidiomycetes, and Zygomycetes. For complete list of diseases controlled, see the label directions. For more information on the use of DITHANE M-45 FLOWABLE, see the label directions. For more information on the use of DITHANE M-45 FLOWABLE, see the label directions. For more information on the use of DITHANE M-45 FLOWABLE, see the label directions.

VEGETABLES

ASPARAGUS—Rust (Uromyces asparagi). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after spears have been harvested.

CARROTS—Alternaria leaf spot and Cercospora. Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in 75 to 125 gallons of water. Start applications when disease first appears and repeat every 7 to 10 days as needed. Do not apply within 7 days of harvest. Do not use treated tops for food or feed.

CELERY FENNEL—Early blight and late blight. Use DITHANE M-45 FLOWABLE at 1.7 qts. per acre in sufficient water for thorough coverage. Start applications in the plant bed when plants first come through the ground. Make further applications at 3 to 5 day intervals until plants are set in the field. Continue field applications on a 7 day schedule. Do not apply within 7 days of harvest or fennel or 14 days of harvest for celery. Remove excess residue remaining at harvest by chopping, trimming, and washing.

CORN (sweet popcorn)—Helminthosporium leaf blight. Use DITHANE M-45 FLOWABLE at 1.3 qts. per acre in sufficient water for thorough coverage. Start applications when disease first appears and repeat at 4 to 7 day intervals. Do not apply within 7 days of harvest. Do not use treated tops for food or feed.

CUCUMBERS CANTALOUPES HONEYDEW MELONS MUSKMELONS SUMMER SQUASH WATERMELONS—Alternaria and Cercospora leaf spots, anthracnose, downy mildew, gummy stem, light and dark. Use DITHANE M-45 FLOWABLE at 1.7 to 2.1 qts. per acre in sufficient water for thorough coverage. Start applications when plants are in the field. Repeat at 5 to 7 day intervals. Direct spray for thorough coverage of both upper and lower leaf surfaces. Do not apply within 5 days of harvest.

ONIONS (dry bulb)—Botrytis leaf blight, downy mildew, purple blotch, and neck rot. Use DITHANE M-45 FLOWABLE at 1.7 to 2.5 qts. per acre in 100 gallons of water. For a protective spray schedule start when diseases are first reported in the area and repeat at 7 day intervals throughout the season. Do not apply within 7 days of harvest and do not allow spray or drift to contact bulbs after coming from soil.

ONIONS (New York Only)—Apply 2.5 qts. of DITHANE M-45 FLOWABLE per acre as a fungicide for the control of Botrytis leaf blight, downy mildew, purple blotch, and neck rot. Use DITHANE M-45 FLOWABLE at 1.7 to 2.5 qts. per acre in 100 gallons of water. For a protective spray schedule start when diseases are first reported in the area and repeat at 7 day intervals throughout the season. Do not apply within 7 days of harvest and do not allow spray or drift to contact bulbs after coming from soil.

POTATOES—Early blight and late blight. Apply every 7 to 10 days throughout the season. For a protective spray schedule start when diseases are first reported in the area and repeat at 7 day intervals throughout the season. Do not apply within 7 days of harvest and do not allow spray or drift to contact tubers after coming from soil.

POTATO SEED TREATMENT—For control of Fusarium seed rot and seed blight. Use DITHANE M-45 FLOWABLE at 1.7 to 2.5 qts. per acre in 100 gallons of water. For a protective spray schedule start when diseases are first reported in the area and repeat at 7 day intervals throughout the season. Do not apply within 7 days of harvest and do not allow spray or drift to contact tubers after coming from soil.

TOMATOES—Early blight, late blight, gray leaf spot, and leaf mold. Use DITHANE M-45 FLOWABLE at 1.7 to 2.5 qts. per acre in 100 gallons of water. For a protective spray schedule start when diseases are first reported in the area and repeat at 7 day intervals throughout the season. Do not apply within 7 days of harvest and do not allow spray or drift to contact fruit after coming from soil.

TOMATOES—Early blight, late blight, gray leaf spot, and leaf mold. Use DITHANE M-45 FLOWABLE at 1.7 to 2.5 qts. per acre in 100 gallons of water. For a protective spray schedule start when diseases are first reported in the area and repeat at 7 day intervals throughout the season. Do not apply within 7 days of harvest and do not allow spray or drift to contact fruit after coming from soil.

FRUIT

APPLES CRABAPPLES PEARS QUINCE—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

Use DITHANE M-45 FLOWABLE fungicide to control a wide range of plant diseases caused by fungi. It is a systemic fungicide which is absorbed by the plant and moves to the site of infection. It is effective against a wide range of fungi including Ascomycetes, Basidiomycetes, and Zygomycetes. For complete list of diseases controlled, see the label directions. For more information on the use of DITHANE M-45 FLOWABLE, see the label directions. For more information on the use of DITHANE M-45 FLOWABLE, see the label directions.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.

APPLES—Rust (Uromyces malvacearum). Use DITHANE M-45 FLOWABLE at 1.3 to 1.7 qts. per acre in sufficient water for thorough coverage. Start applications when rust first appears and repeat at 10 day intervals as long as rust persists. Four applications will usually be sufficient. Apply only in a parasitic form after fruit has been harvested.