

P1116

11/1/89

11/1/89

RESTRICTED USE PESTICIDE

Because pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.


ACCEPTED

11/1/89
77-159

SEE PAMPHLET IN BOTTOM OF BAG FOR COMPLETE USE DIRECTIONS

KERB®

WSP



TURF & ORNAMENTAL HERBICIDE

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Take special care to avoid contact with eyes, skin or clothing. Do not breathe the dust or spray mist. Wash clothing and gloves after use.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person. **Note to Physician:** Emesis is recommended. **If Inhaled:** Move subject to fresh air. **If on Skin:** Wash affected skin areas with soap and water. Consult a physician if irritation persists. **If in Eyes:** Flush eyes with large amounts of water for at least 15 minutes. Consult a physician if irritation persists.

PROTECTIVE CLOTHING: When applying this product, wear mid-forearm waterproof gloves, long-sleeved shirt and long pants, preferably one piece (coveralls). Hand-spray or hand-spreaders also require the use of waterproof boots or shoe coverings. Wash nondisposable gloves, boots and shoe coverings thoroughly with soap and water before removing. Protective clothing/equipment is not needed during application if an enclosed tractor cab with filtered air supply or enclosed cockpit is used.

Any article of clothing worn while handling product must be cleaned before reusing. Clothing should be laundered separately from household articles. Clothing that has been drenched or heavily contaminated should be disposed of in accordance with state or local regulations.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on front and back panel and attached use directions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.



ROHM AND HAAS

PHILADELPHIA PA 19105
TEL 2151 592-3000

ACTIVE INGREDIENT

Pronamide
3,5-dichloro-N-(1,1-dimethyl-2-propynyl)-benzamide 50.0%

INERT INGREDIENTS 50.0%

TOTAL 100.0%

EPA Reg. No 707-159 EPA Est. No. 33967-NJ-1 (B) 65387-AR-001 (C)
Superscript used is first letter of Lot Number

NET CONTENTS: 3 LBS (1.36 KG)

3 x 1 LB WATER-SOLUBLE POUCHES

BEST AVAILABLE COPY

(1 inch)

317 250 Hwy - Rohm and Haas - DC - 11/1/91

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SEE PAMPHLET IN BOTTOM OF BAG FOR COMPLETE USE DIRECTIONS.

USE RESTRICTIONS: Hand-spray applications of pronamide may be made only to ornamentals and nursery stock.

RE-ENTRY AND WORKER PROTECTION STATEMENTS

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. SEE USE DIRECTIONS FOR ADDITIONAL RE-ENTRY STATEMENTS.

GENERAL INFORMATION

MIXING: The enclosed pouches of KERB WSP herbicide are water-soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately five minutes from the time they are added to the water.

DOSAGE: Refer to the specific crop for dosage recommendations. Determine the total amount of KERB WSP to be added to the spray tank. For each pound of KERB WSP, use one one-pound pouch.

Example: If two pounds of KERB WSP should be added to the spray tank, add two one-pound pouches. The type of equipment used will determine the concentration required; however, the use of these water-soluble pouches is not recommended for making highly concentrated mixtures such as those used in aircraft spraying unless pouches are premixed in nurse tanks with adequate agitation.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in a cool, dry place but not below 32°F (0°C). Do not remove package from container except for immediate use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.


CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Sweep up and place in containers for recovery or disposal. Keep dusting to a minimum. Wear MSHA/NIOSH respirator for dust encountered. Flush area with water. Keep spill out of all sewers and open bodies of water. Protective Clothing — REFER TO PRECAUTIONARY STATEMENTS.

RESTRICTED USE PESTICIDE

Because pronamide has produced tumors in laboratory animals, this product is for retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

KERB®



TURF & ORNAMENTAL HERBICIDE

USE DIRECTIONS FOR:

- Bermudagrass (Ornamental)
- Woody Ornamentals/Nursery Stock and Christmas Trees

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed. Take special care to avoid contact with eyes, skin or clothing. Do not breathe the dust or spray mist. Wash clothing and gloves after use.

PROTECTIVE CLOTHING: When applying this product, wear midforearm waterproof gloves, long-sleeved shirts and long pants, preferably one-piece coveralls. Hand-spray or hand-spreaders also require the use of waterproof boots or shoe coverings. Wash nondisposable gloves, boots and shoe coverings thoroughly with soap and water before removing.

Protective clothing/equipment is not needed during application if an enclosed tractor cab with filtered air supply or enclosed cockpit is used.

Any article of clothing worn while handling product must be cleaned before reusing. Clothing should be laundered separately from household articles. Clothing which has been drenched or heavily contaminated should be disposed of in accordance with state or local regulations.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Emesis is recommended.

IF INHALED: Move subject to fresh air.

IF ON SKIN: Wash affected skin areas with soap and water. Consult a physician if irritation persists.

IF IN EYES: Flush eyes with large amounts of water for at least 15 minutes. Consult a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

ACTIVE INGREDIENT

Pronamide
3,5-dichloro-N (1,1-dimethyl-2-propynyl)-
benzamide 50.0%

INERT INGREDIENTS 50.0%

Total 100.0%

EPA Reg. No. 707-159

©Copyright 1991 Rohm and Haas Company



CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seiler. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

Hand-spray applications of pronamide may be made only to ornamentals and nursery stock.

CHEMIGATION

Do not apply this product through any type of irrigation system.

RE-ENTRY AND WORKER PROTECTION STATEMENTS

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

CAUTION

(Insert area or field description) treated with KERB WSP herbicide on (insert date of application). Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to KERB WSP herbicide and describe his condition. For further information see the "Statement of Practical Treatment."

Oral warnings must include the same information that is in the written warnings.

BEST AVAILABLE COPY

1 of 6

GENERAL INFORMATION

KERB wsp selective herbicide is formulated as a wettable powder containing 50% active ingredient. It is effective for the control of a wide range of grasses and certain broadleaf weeds. The product is a soil active herbicide with uptake by sensitive weeds occurring through the roots. Before using this herbicide for a specific crop use, study the following general use information which provides important instructions for the safe and effective application of the product.

MIXING

The enclosed pouches of KERB wsp herbicide are water-soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately five minutes from the time they are added to the water.

DOSAGE

Refer to the specific crop for dosage recommendations. Determine the total amount of KERB wsp to be added to the spray tank. For each pound of KERB wsp, use one-pound pouch.

Example: If two pounds of KERB wsp should be added to the spray tank, add two one-pound pouches.

The type of equipment used will determine the concentration required; however, the use of these water-soluble pouches is not recommended for making highly concentrated mixtures such as those used in aircraft spraying unless pouches are premixed in nurse tanks with adequate agitation.

WEED SPECTRUM

KERB wsp herbicide is recommended for both preemergence and early postemergence control of winter annual and perennial grasses and chickweed and for preemergence control only of certain other broadleaf weeds and certain other grasses listed.

WEEDS CONTROLLED BOTH PREEMERGENCE AND EARLY POSTEMERGENCE

BARLEY, FOXTAIL	<i>Horaeum jubatum</i>
BARLEY, VOLUNTEER	<i>Horaeum vulgare</i>
BLUEGRASS, ANNUAL	<i>Poa annua</i>
BLUEGRASS, KENTUCKY	<i>Poa pratensis</i>
BROME, DOWNY (CHEATGRASS)	<i>Bromus tectorum</i>
CHICKWEED, COMMON	<i>Stellaria media</i>
CHICKWEED, MOUSE-EAR	<i>Cerastium vulgatum</i>
OAT, VOLUNTEER	<i>Avena sativa</i>
ORCHARDGRASS	<i>Dactylis glomerata</i>
QUACKGRASS	<i>Agropyron repens</i>
RYE, VOLUNTEER	<i>Secale cereale</i>
RYEGRASS, ITALIAN	<i>Lolium multiflorum</i>
RYEGRASS, PERENNIAL	<i>Lolium perenne</i>
WHEAT, VOLUNTEER	<i>Triticum aestivum</i>

WEEDS CONTROLLED ONLY PREEMERGENCE

MUSTARD, WILD	<i>Brassica kaber</i>
ROCKET, LONDON	<i>Sisymbrium irio</i>
SHEPHERDSPURSE	<i>Capsella bursa-pastoris</i>
SORREL, RED (from seed)	<i>Rumex acetosella</i>

NOTE: The weed species controlled by KERB wsp herbicide are dependent on the rate used, specific crop culture involved, and the associated conditions of temperature, soil type and moisture availability. Refer to specific crop use directions for weed species controlled.

DOSAGE

The rate of KERB wsp herbicide required will vary depending on the crop culture involved and weed species to be controlled. See specific crop use directions for all dosage recommendations. All dosage recommendations listed in this label are in terms of pounds KERB wsp herbicide per broadcast acre. For banded application, the amount of KERB wsp herbicide used per acre should be reduced according to the following formula:

$$\frac{\text{Band width (in inches)}}{\text{Row width (in inches)}} \times \text{Rate per Acre Broadcast} = \text{Amount Needed per Acre for Band Application}$$

TIMING AND APPLICATION

Unless specific directions are given under the crop to be treated, KERB wsp herbicide should be applied in the fall or early winter, when temperatures do not exceed 55°F, BUT PRIOR TO SOIL FREEZE-UP. Best weed control results when KERB wsp herbicide is applied preemergence to the weeds and when application is followed by rainfall or irrigation to move the product into the root zone of the germinating weeds.

KERB wsp herbicide should be thoroughly mixed in clean water at the recommended concentration and applied uniformly as a spray. For ground application, use a conventional low pressure herbicide sprayer equipped with flat fan nozzles spaced and calibrated to uniformly deliver 20 to 50 gallons of spray per acre. For aerial applications apply in a coarse droplet spray at 5 to 10 gallons per acre. Accurately calibrate spray equipment prior to each use.

EFFECT OF SOIL TYPE, MOISTURE AND TEMPERATURE

KERB wsp herbicide is most active in coarse to medium-textured soils of low organic matter and is relatively inactive in peat or muck soils or mineral soils high in organic matter content at rates recommended in this label. Herbicidal activity is best in soils containing less than 4 percent organic matter. Use in soils with higher organic matter may result in inconsistent or incomplete weed control.

The herbicidal activity of KERB wsp is mainly through root absorption in sensitive weed species. Rain, melting snow or irrigation is ESSENTIAL following treatment to move KERB wsp herbicide into the root zone of germinating weeds.

Under field conditions, KERB wsp herbicide will remain relatively stable with little loss of herbicidal activity when soil temperatures are less than 55°F. As soil temperatures increase, degradation of the active ingredient takes place. KERB wsp herbicide may degrade rather quickly if left exposed on the soil surface in warm weather. If KERB wsp herbicide is applied when air temperatures exceed 85°F, it should be soil incorporated to a shallow depth (top two to three inches) or watered into the soil as soon as possible.

CULTURAL CONSIDERATIONS

For best results apply KERB wsp herbicide to a trash free soil surface. Clean cultivation before application is preferable, but not necessary. To obtain optimum weed control in areas not clean cultivated, the area to be treated should be free of surface litter (dead or decaying crop and weed debris, mowing clippings, etc.) Trash free areas create ideal conditions for rapid movement of KERB wsp into the weed root zone following rain or irrigation.

ROTATION CROP PLANTING INFORMATION

Follow the directions given below when rotation crops will be planted to areas previously treated with KERB wsp herbicide:

1. Whether KERB wsp herbicide is bed-topped, banded or broadcast, the beds should be knocked down and the field cross-disked before rotation crops other than artichokes, lettuce, endive or escarole are planted.
2. Where the KERB wsp herbicide treatment is to be followed by a rotation crop within six months of application, bed-topped or banded applications are suggested.
3. Where small grains or grasses will be planted within one year to areas where KERB wsp herbicide is to be applied at a dosage rate of 1.0 pound of product per acre or higher, KERB wsp herbicide should not be broadcast. Treatments should be limited to band or bed-top applications only.
4. Where rotation crops are to follow within one year of the KERB wsp herbicide treatment, the following rotational plan applies:

BEST AVAILABLE COPY

Amount of KERB wsp Applied/Planted Acre ²	Waiting Period in Months Before Planting the Crops Indicated ¹			
	Group A	Group B	Group C	Group D
1.0 lb.	0	3	3	6
2.0 lb.	0	3	4	9
3.0 lb.	0	4	6	9
4.0 lb.	0	5	7	12

- ¹Crop Group A — Artichokes, lettuce, endive, escarole
- Crop Group B — Beans, corn, cotton, sorghum, carrots, celery
- Crop Group C — Broccoli, cabbage, cauliflower, cucurbits, spinach, sugar-beets, onion, tomatoes
- Crop Group D — Wheat, barley, oats, grasses

NOTE: "Crops not listed in Groups A, B, C or D (above) should not be planted back sooner than 12 months after the last KERB wsp herbicide application regardless of the rate of KERB wsp used."

²Planted Acre refers to Acreage planted. KERB wsp at 4.0 pounds of product per treated acre on two 5 inch treated bands (40-inch centers) is the equivalent of one pound KERB wsp per planted acre. KERB wsp at 4.0 pounds of product as a broadcast treatment (total acreage sprayed) is the equivalent of 4 pounds of KERB wsp per planted acre.

BERMUDAGRASS (ORNAMENTAL)

GROWN FOR TURF OR SEED

GENERAL INFORMATION

KERB wsp is a selective herbicide recommended for the preemergence and postemergence control of annual bluegrass (*Poa annua*) in ornamental bermudagrass (*Cynodon dactylon*) turf or bermudagrass grown for seed.

WEED CONTROL

KERB wsp herbicide will control annual bluegrass from pregermination and seedling stages through tillering, heading and seed formation. KERB wsp herbicide acts slowly on seedling to mature annual bluegrass. Following application of KERB wsp annual bluegrass may first become dark green and then gradually turn yellow and die over a 3 to 5 week period.

For effective control of annual bluegrass in bermudagrass turf or seed crops, moisture is necessary to move KERB wsp herbicide in the weed root zone. Refer to the Moisture Requirement section of this label for details.

DOSAGE AND TIMING

KERB wsp herbicide is recommended at the rate of 1.0 to 3.0 pounds of product (0.5 to 1.5 pounds active ingredient) per acre broadcast application. The dosage rate required is dependent on the growth stage of annual bluegrass at time of application. Follow the dosage rate and timing recommendations given below:

Annual Bluegrass Growth Stage	KERB wsp	
	Oz. per 1000 Sq. Feet	Lbs. per Acre Broadcast Application
Preemergence or early postemergence	0.4 - 0.75	1.0 - 2.0*
Postemergence — early tillering to heading	0.6 - 0.75	1.5 - 2.0
Postemergence-seed forming stages	0.75 - 1.1	2.0 - 3.0

*Use the higher rate when longer preemergence residual control is desired.

APPLICATION

Mix the recommended amount of KERB wsp herbicide in clean water and apply uniformly with a low pressure ground sprayer in 20 to 50 gallons of water per acre or 0.5 to 1 gallon of water per 1000 sq. feet. The sprayer should be equipped with flat fan nozzles, spaced to provide uniform distribution without skips or excessive overlapping of spray patterns.

NOTE: Avoid spraying on fairways or approaches that may drain onto bentgrass greens or to areas overseeded with sensitive cool season grasses.

MOISTURE REQUIREMENTS

KERB wsp herbicide acts mainly through root absorption in sensitive weed species. If no rainfall occurs within a day or so of the application, a light overhead irrigation should be made to move the chemical into the weed root zone. Avoid heavy irrigations of more than 1 inch to reduce the possibility of excess washing or leaching of the chemical from the area of application.

BERMUDAGRASS (ORNAMENTAL)

TURF OR SEED CROP

— SPECIFIC USE RESTRICTIONS

- Do not apply KERB wsp herbicide to areas that are to be overseeded with susceptible cool season grasses within 90 days of treatment.
- Do not apply KERB wsp to dicentra, perennial bluegrass, annual and perennial ryegrasses, fescues and bentgrasses.
- KERB wsp herbicide is not recommended for use on golf greens.
- Do not graze treated areas and do not feed clippings to livestock.

WOODY ORNAMENTALS/NURSERY STOCK AND CHRISTMAS TREES

GENERAL INFORMATION

KERB wsp is a selective herbicide recommended for fall applications to established woody ornamentals, nursery stock and Christmas trees for the control of winter annual and perennial grasses and certain broadleaf weeds.

CROP TOLERANCE

At recommended rates of KERB wsp herbicide the following trees and shrubs are tolerant to topical applications made in the fall:

ASH	FLORIBUNDA	MOUNTAIN ASH
ARBORVITAE	FLOWERING CHERRY	MOUNTAIN LAUREL
AZALEA	FLOWERING CRABAPPLE	OHIO BUCKEYE
BARBERRY	FLOWERING QUINCE	OAK
BASSWOOD	FOXSYPHIA	PINE
BEECH	GINKGO	POPLAR
BIRCH	HAWTHORN	PRIVET
BOXWOOD	HEMLOCK	RHODODENDRON
BRADFORD PEAR	HOLLY	SPIREA
CEDAR	HONEY LOCUST	SPRUCE
COTONEASTER	JUNIPER	SWEETGUM
DOGWOOD	LILAC	SYCAMORE
DOUGLAS FIR	LINDEN	TUL-FR-FR
EASTERN REDBUD	LONDON PLANE	VIBURNUM
ELM	MAGNOLIA	WALNUT
EUONYMUS	MAPLE	WILLOW
FIR	MOCK ORANGE	YEW

KERB wsp herbicide should only be used on established trees and woody ornamentals. KERB wsp herbicide is not recommended for use on seedling trees or shrubs less than one year old or to fall transplanted stock transplanted less than one year or to spring transplanted stock transplanted less than six months.

WEED CONTROL

KERB wsp herbicide is effective in fall applications at the rate of 2.0 to 4.0 pounds of product (1.0 to 2.0 pounds active ingredient) per broadcast acre for the preemergence and postemergence control of susceptible winter annual and perennial grasses and chickweed and for preemergence control only of other broadleaf weeds listed on this label. Refer to chart in dosage rate section below for specific weeds controlled.

DOSAGE AND TIMING

KERB wsp herbicide is recommended in a single, fall application, either directed or topically applied, to woody ornamentals, nursery stock or Christmas trees at the rate of 2.0 to 4.0 pounds of product (1.0 to 2.0 pounds active ingredient) per broadcast acre. Application of KERB wsp herbicide should be made in the fall prior to leaf drop and soil freeze-up for control of winter annual or perennial grasses or chickweed. Applications can be made either preemergence or postemergence to the weeds. For control of other labeled broadleaf weeds, preemergence applications must be used to achieve control.

The dosage rate required will depend on the weed species present in the area to be treated. Follow the weed control recommendations given in the chart below:

<u>Weeds Controlled</u>	<u>Pounds KERB wsp Per Acre Broadcast Application</u>
BARLEY, FOXTAIL BLUEGRASS, ANNUAL BROME, DOWNY (CHEATGRASS) CHICKWEED GRAIN VOLUNTEER RYEGRASS, ITALIAN SORREL RED (from seed)	2.0
MUSTARD, WILD ROCKET, LONDON SHEPHERDSPURSE	3.0
BLUEGRASS, KENTUCKY ORCHARDGRASS QUACKGRASS RYEGRASS, PERENNIAL	4.0

APPLICATION

Mix the recommended amount of KERB wsp herbicide in clean water and apply uniformly in 20 to 50 gallons per acre. Use a low pressure ground sprayer equipped with flat fan nozzles spaced to provide uniform distribution.

Dosage recommendations listed on this label are for surface broadcast application. For banded treatments down the row, the amount of KERB wsp used per acre should be reduced according to the following formula:

$$\frac{\text{Band width (in inches)}}{\text{Row width (in inches)}} \times \text{Rate per Acre Broadcast} = \text{Amount Needed per Acre for Band Application}$$

KERB wsp herbicide should not be soil incorporated.

NOTE: Most ornamental turf grass species and ground covers are sensitive to KERB wsp. Care should be exercised to avoid contact of KERB wsp herbicide with these plants from either direct application, spray drift or from applications to areas that may drain onto established ornamental turf and ground cover.

SOIL AND MOISTURE REQUIREMENTS

KERB wsp herbicide is most active in coarse to medium-textured soils of low organic matter and is relatively inactive in peat or muck soils or mineral soils high in organic matter content at rates recommended in this label. Herbicidal activity is best in soils containing less than 4 percent organic matter. Use in soils of higher organic matter content may result in inconsistent or incomplete weed control.

KERB wsp herbicide acts mainly through root absorption in sensitive weed species. Dependable rainfall or overhead irrigation is essential following application for effective weed control.

WOODY ORNAMENTALS/NURSERY STOCK AND CHRISTMAS TREES

SPECIFIC USE RESTRICTIONS

- Apply KERB wsp herbicide in the fall prior to soil freeze-up.
- Do not soil incorporate KERB wsp herbicide.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

STORAGE: Store in a cool, dry place but not below 32°F (0°C). Do not remove package from container except for immediate use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Sweep up and place in containers for recovery or disposal. Keep dusting to a minimum. Wear MSHA/NIOSH respirator for dusts encountered. Flush area with water. Keep spill out of all sewers and open bodies of water. Protective Clothing — See Precautionary Statements.

8384-80

MADE IN U.S.A.

KERB is a registered trademark of the Rohm and Haas Company.

BEST AVAILABLE COPY