

# ROHM AND HAAS COMPANY

INDEPENDENCE MALL WEST  
PHILADELPHIA, PENNSYLVANIA 19105



## CAPTAN 50W Agricultural Fungicide

EPA Reg. No. 707-160  
EPA Est. No. 279-FL-1

**ACCEPTED**  
*Per self-cert. RFR*  
**OCT 19 1987**  
Use: the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the pesticide registered under EPA Reg. No. 707-160

### Active Ingredient

- \*Captan ..... 48.9%
- Related Derivatives ..... 1.1%
- Inert Ingredients ..... 50.0%
- \*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**

**PELIGRO**

Precaucion Al Usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

### Precautionary Statements

#### Hazards to Humans and Domestic Animals

Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin or clothing. Remove and separately launder clothing before reuse.

#### Statement of Practical Treatment

**If in Eyes:** Flush eyes with a large amount of water for at least 15 minutes. Get medical attention.

**If on Skin:** Wash affected skin areas with plenty of soap and water.

**If Swallowed:** Dilute by giving 2 glasses of water to drink. Contact physician, poison control center, or emergency clinic before inducing vomiting.

**If Inhaled:** Move victim to fresh air.

#### Environmental Hazards

This product is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

In event that treated seeds are spilled and cannot be salvaged, completely cover them with soil.

Storage and Disposal

Storage Keep pesticide in original container. Do not put concentration or dilute into food or drink containers. Do not store or transport near feed or food. Store in a cool, dry place.

Pesticide Disposal Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Step to be taken if Material is Released or Spilled: Scoop or shovel solid material into suitable container for recovery or disposal. Keep dust to a minimum. Refer to all Precautionary Statements. Keep spills and cleaning runoffs out of municipal sewers and open bodies of water.

Directions for Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

Endangered Species Restriction

The following restrictions apply to use of this product after February 1, 1988.

Before using this pesticide on corn, cotton, soybeans, sorghum, oats, barley and rye in the counties listed below you must obtain the Pesticide Use Bulletin for Protection of Endangered Species for the county in which the product is to be used. The bulletin is available from your County Extension Agent, State Fish and Game Office, or your pesticide dealer. Use of this product in a manner inconsistent with the Pesticide Use Bulletin for Protection of Endangered Species is a violation of Federal law.

ALABAMA - Lauderdale, Limestone and Madison

ARIZONA - Graham, Maricopa, Mohave, Pima, Pinal and Santa Cruz


ARKANSAS - Benton and Polk

CALIFORNIA - Imperial, Modoc and Riverside

MISSISSIPPI - Claiborne, Copiah and Hinds

MISSOURI - Barry, Benton, Camden, Christian, Dallas, Greene, Hickory, Jasper, Lawrence, Miller, Newton, Osage, Polk, St. Clair, Stone and Webster

NEVADA - Clark



8/20/82

NEW MEXICO - Chaves, DeBaca and Eddy

OHIO - Pickaway

OKLAHOMA - Delaware, McCurtain and Pushmataha

OREGON - Lake

TENNESSEE - Lawrence and Wayne

TEXAS - Bastrop, Burleson, Correll, Harris, Hays, Jeff Davis, Pecos and Reeves

UTAH - Utah and Washington

VIRGINIA - Grayson, Lee, Scott, Smyth, Washington

5 2  
3 3

RW/dda  
(0222g/11g)

**REENTRY STATEMENT**

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift.

Do not allow persons to enter treated areas within 4 days following application unless a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Conspicuously post reentry information at site of application.

Certain states may require more restrictive reentry intervals; consult your State Department of Agriculture for further information.

Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

**DIRECTIONS**

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves if entering treated areas within 4 days following application. Leather or fabric gloves are not acceptable.

A dust mask must be worn when mixing and loading this product.

During application only, if from a tractor with a completely enclosed cab a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab and worn while exiting.

**AFTER WORK:** Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

**NOTE:** For concentrate sprayer applications, the same quantity (per acre) of CAPTAN 50 Wettable as would be normally applied when spraying dilute mixture, should be applied to each acre in sufficient water for coverage.


**CHEMIGATION**

Do not apply this product through any type of irrigation system.

**ALLIUMS:** Brown Rot, Twig and Blossom Blight, Shot Hole, Scab, Leaf Blight—Apply 9 lbs. CAPTAN 50 Wettable per acre in 20 to 300 gals. of water using ground equipment. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. Do not apply within 30 days of harvest. Hulls may be fed to livestock. Subsequent applications may be made up to 12 days of harvest in accordance with pest infestations and State Agricultural Extension Service recommendations, provided the hulls are not fed to dairy animals or animals being finished for slaughter.

**APPLES (Eastern U.S.):** Primary Scab, Black Rot, Botrytis Blossom End Rot—Apply 8 lbs. CAPTAN 50 Wettable per acre in 20 to 400 gals. of water using ground equipment. Apply at 5 to 7 day intervals as needed to maintain control in prebloom, bloom, petal fall and first cover sprays.

Secondary Scab, Brooks Fruit Spot, Sooty Blotch, Fly Speck, Black Rot, Black Pox, Botryospheria Rot, Bitter Rot—Apply 4 to 8 lbs. CAPTAN 50 Wettable per acre in 20 to 400 gals. of water using ground equipment. Apply at 10 to 14 day intervals in second and latter cover sprays up to start of harvest. Powdery Mildew—If powdery mildew is a problem add 6 to 12 lbs. sulfur or 1 to 2 lbs. of Karathane per acre to all postbloom sorays until the foliage matures. Use 2 lbs. of Karathane if interval between sprays is longer than 7 days. **CAUTION—Do**



**BEST AVAILABLE COPY**

not use CAPTAN 50 Wettable in combination with or closely following or in alternation with wettable sulfur products or sulfur sensitive varieties of apples such as Red Delicious, Staymen, Baldwin King, etc. as severe injury and defoliation may occur.

(Western U.S.): Apple Scab—Apply 8 lbs. CAPTAN 50 Wettable per acre in a high volume application for preblossom sprays. For preblossom sprays using low volume application, apply 16 lbs. per acre. In mid-summer, the dosage may be reduced to 4 lbs. per acre in high volume applications or 10 lbs. per acre in low volume applications.

(Pacific Northwest): Bull's Eye Rot, Botrytis Rot—Apply 8 lbs. CAPTAN 50 Wettable per acre in 20 to 400 gals. of water using ground equipment. Make 1 or 2 applications with late cover sprays and 1 to 2 preharvest applications.

APPLES—POST HARVEST DIP: For control of Various Molds and Storage Rots (Botrytis, Gloeosporium, Rhizopus)—Use as a post harvest dip or wash. Mix 2½ lbs. CAPTAN 50 Wettable per 100 gals. water. Apply as a spray or in a dip tank. When used as a dip, recharge solution when tank volume is reduced by 25%. Bring water back to volume and add 1 lb. for each 25 gals. added. At the end of every 8 hour shift empty tank, flush and mix with a fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

APRICOTS: Brown Rot (Twig Blight) Jacket Rot—Apply 5 lbs. CAPTAN 50 Wettable per acre in 20 to 250 gals. of water using ground equipment. Apply at red bud, bloom, 75% petal fall stages and in cover sprays.

CANTALOUPE: Angular Leaf Spot, Anthracnose—Apply 4 lbs. CAPTAN 50 Wettable per acre in sufficient water for thorough coverage. Begin applications at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control.

CHERRIES (Eastern U.S.): Brown Rot, Leaf Spot, Botrytis Rot—Apply 4 lbs. CAPTAN 50 Wettable per acre in 20 to 200 gals. of water using ground equipment. Apply in pre-bloom, bloom, petal fall, shuck, cover and preharvest sprays. Application at 3 to 4 day intervals may be necessary during bloom to control Blossom Blight. Repeat applications at 7 to 10 day intervals as needed to maintain control up to start of harvest. If powdery mildew is a problem add 6 lbs. sulfur per acre to the petal fall, shuck and early cover sprays. If sulfur is added, CAPTAN 50 Wettable may be reduced to 2 lbs. per acre in these sprays.

POST HARVEST SPRAYS—Leaf Spot: Apply 4 lbs. CAPTAN 50 Wettable per acre in 20 to 200 gals. of water using ground equipment. Apply immediately after harvest and repeat application in 10 to 14 days. (Western U.S.): Brown Rot, Blossom Blight, Brown Rot (fruit), Leaf Spot: Apply 4 lbs. CAPTAN 50 Wettable per acre in 20 to 200 gals. of water using ground equipment. Apply in prebloom, bloom, petal fall, shuck, cover and preharvest sprays.

CHERRIES—POST HARVEST DIP: For control of Various Molds and Storage Rots (Botrytis, Gloeosporium, Rhizopus)—Use as a post harvest dip or wash. Mix 2½ lbs. CAPTAN 50 Wettable per 100 gals. water. Apply as a spray or in a dip tank. When used as a dip, recharge solution when tank volume is reduced by 25%. Bring water back to volume and add 1 lb. for each 25 gals. added. At the end of every 8 hour shift empty tank, flush and mix with a fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

CUCUMBER: Angular Leaf Spot, Anthracnose—Apply 4 lbs. CAPTAN 50 Wettable per acre in sufficient water for thorough coverage. Begin applications at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control.

DAMP OFF on Cullings of Azaleas, Carnations and Chrysanthemums—Dip cuttings in a suspension of 4 lbs. per 100 gals. spray before bedding.

GRAPES (Eastern U.S.): Black Rot, Downy Mildew—Apply 4 lbs. CAPTAN 50 Wettable per acre in 10 to 200 gals. of water using ground equipment. Apply first application just before bloom, repeat application immediately after bloom and at 10 to 14 day intervals for 1 to 3 additional applications depending on disease pressure and susceptibility of the grape variety.

Dead Arm (current season infection)—Apply 2 to 4 lbs. CAPTAN 50 Wettable per acre in 10 to 200 gals. of water using ground equipment. Apply first application when shoots are ½ to 1 inch long and repeat application when shoots are 4 to 6 inches long.

GRAPES (California): Bunch Rot (Botrytis)—Apply 4 lbs. CAPTAN 50 Wettable per acre in 10 to 200 gals. of water using ground equipment. Apply 2 applications before bloom and 1 immediately after bloom. Repeat periodically making 3 cover sprays before bunches close.

Dead Arm (current season infection)—Apply 3 to 6 lbs. CAPTAN 50 Wettable per acre in 10 to 200 gals. of water using ground equipment. Apply first spray when green tissue begins to show but before shoots are 1 inch long and repeat application when shoots are 6 to 8 inches long.

NECTARINES: Brown Rot, Scab—Apply 5 lbs. CAPTAN 50 Wettable per acre in 20 to 250 gals. of water using ground equipment. Apply in full pink, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control Blossom Blight. Repeat applications at 7 to 14 day intervals as necessary to maintain control. Continue applications through harvest if conditions favor Brown Rot. If powdery mildew is a problem, add 7.5 lbs. sulfur per acre to the petal fall and shuck sprays and early cover spray. If sulfur is added, CAPTAN 50 Wettable may be reduced to 2.5 lbs. per acre in these sprays.

Coryneum Blight (Peach Blight, Shot Hole)—Apply 5 lbs. CAPTAN 50 Wettable per acre in 20 to 250 gals. of water using ground equipment. Apply in pink bud, full bloom, petal fall and cover sprays as necessary and as a post harvest spray (but before leaves drop).

ORNAMENTALS: ROSES—Black Spot, CARNATIONS—Rust and Leaf Spot, CHRYSANTHEMUMS—Flower Blight—2 lbs. per 100 gals. spray. Apply at first signs of diseases, cover upper and lower leaf surfaces thoroughly. Repeat every 7 to 10 days.

PEACHES: Brown Rot, Scab—Apply 8 lbs. CAPTAN 50 Wettable per acre in 20 to 400 gals. of water using ground equipment. Apply at full pink, bloom, petal fall, shuck stages and in cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control Blossom Blight. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown Rot. If powdery mildew is a problem, add 12 lbs. of sulfur per acre to the petal fall and shuck sprays and early cover spray. If sulfur is added, CAPTAN 50 Wettable may be reduced to 4 lbs. per acre in these sprays.

Coryneum Blight (Peach Blight, Shot Hole)—Apply 8 lbs. CAPTAN 50 Wettable per acre in 20 to 400 gals. of water using ground equipment. Apply at pink bud, full bloom and petal fall stages and in cover sprays as necessary and as a post harvest spray (but before leaf drop).

PEARS (Eastern U.S.): Scab—Apply 8 lbs. CAPTAN 50 Wettable per acre in 20 to 200 gals. of water using ground equipment. Apply in preblossom, petal fall and cover sprays. Repeat cover spray applications at 7 to 14 day intervals under severe scab infection periods. Russeting may be reduced on Bosc pears. Do not use on D'Anjou pears.

... Use 2 to 3 lbs. per acre in 20 to 300 gals. of water using ground equipment. Apply during early finger stage, petal fall and first cover sprays. Under severe scab infection periods additional treatments may be necessary. Russeting may be reduced on Bosc pears. Do not use on D'Anjou pears.

**PEARS—POST HARVEST DIP:** For control of Various Molds and Storage Rots (Botrytis, Gloeosporium, Rhizopus)—Use as a post harvest dip or wash. Mix 2½ lbs. CAPTAN 50 Wettable per 100 gals. of water. Apply as a spray or in a dip tank. When used as a dip, recharge solution when tank volume is reduced by 25%. Bring water back to volume and add 1 lbs. for each 25 gals. added. At the end of every 8 hour shift empty tank, flush and mix a fresh dilution. Do not allow dip tank solution to stand overnight. Maintain continuous agitation during dipping operation.

**PLUMS, PRUNES (Western U.S.): Brown Rot**—Apply 6 lbs. CAPTAN 50 Wettable per acre in 20 to 300 gals. of water using ground equipment. Apply at green bud, popcorn, bloom and petal fall stages. Repeat in cover sprays as conditions warrant.

**California: Prune Russet Scab (lacy scab)**—Apply 6 lbs. CAPTAN 50 Wettable per acre in 20 to 300 gals. of water using ground equipment. Apply at full bloom.

**PLUMS, PRUNES (Eastern U.S.): Brown Rot**—Apply 6 lbs. CAPTAN 50 Wettable per acre in 20 to 300 gals. of water using ground equipment. Apply in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown Rot. The addition of a neutral spreader has improved coverage.

**SOIL AND GREENHOUSE BENCH TREATMENT:** Preplanting for Damp Off and Root Rot Diseases on Seedlings or Transplants of Roses (and other shrubs and trees). Many Flowers and Lawn Seedbeds—Apply as a dust at 4 to 5 oz. per 1000 sq. ft. or spray at 2 lbs. to 100 gals. spray, using 15 gals. to each 1000 sq. ft. Cultivate into upper 3 to 4 inches of soil before planting. Only the applicator is permitted to be in the greenhouse during application of captan. Open vents to greenhouse during application and for at least 1 hour after application. Workers planting in captan treated soil in greenhouses must wear chemical resistant gloves.


**STRAWBERRIES: Botrytis (Gray Mold, Leaf Spot)**—Apply 6 lbs. CAPTAN 50 Wettable per acre in sufficient water for thorough coverage. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at weekly intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking.

**TOMATOES: Anthracnose, Gray Leaf Spot, Septoria Leaf Spot**—Apply 6 lbs. CAPTAN 50 Wettable per acre in sufficient water for thorough coverage. Under severe infection conditions use 6 to 8 lbs. per acre in sufficient water for thorough coverage. Apply at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control.

**TURF, LAWN: Brown Patch, Leaf Spot, Damp Off, Root Rot**—2 lbs. per 100 gals. spray. Apply 10 gals. to each 1000 sq. ft. of turf or apply 8 lbs. per acre in water for coverage. Apply prior to infection. Repeat at 7 to 10 day intervals as necessary. Do not graze treated areas or feed clippings to livestock.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

**COMPATIBILITY:** CAPTAN 50 Wettable may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline materials such as hydrated lime, which reduces the fungicidal activity. Do not use this product in combination with oil sprays or with solvent formulations of Parathion products. Do not use this product on foliage closely following oil sprays.



**BEST AVAILABLE COPY**

Smut, Seed Rot, Molds, Storage Rot, Seed and Soil Borne Fungi, Not for Smuts of Barley, Oats, Rye or Wheat.

7008

CROP	METHOD OF APPLICATION			Ounces (actual)	SLURRY	DRY	PLANTER BOX	Ounces (actual)
	OZ. CAPTAN 50-W/ 100 lbs.	OZ. CAPTAN 50-W/ 100 lbs.	OZ. CAPTAN 50-W/ 100 lbs.		OZ. CAPTAN 50-W/ 100 lbs.	OZ. CAPTAN 50-W/ 100 lbs.	OZ. CAPTAN 50-W/ 100 lbs.	
Alfalfa	4.4-8.26	12.0	0.8	—	—	—	—	1.5-2.0/lbs. seed
Barley	1.2-3.8	1.2-4.0	1.6-4.0	—	—	—	—	8.0 oz./lbs.
Beans	1.6-3.0	3.0-3.8	0.4-2.0	—	—	—	—	—
Beans (limb)	1.6-3.0	3.0-3.8	0.4-2.0	—	—	—	—	—
Beets, sugar	1.2-19.2	1.2-18.0	—	—	—	—	—	—
Beets, table	5.8-12.0	18.0	—	—	—	—	—	—
Bluegrass	4.4-8.26	12.0	—	—	—	—	—	—
Broccoli	0.8-4.5	2.25-3.6	—	—	—	—	—	—
Brussels sprouts	0.8-4.5	2.25-3.6	—	—	—	—	—	—
Cantaloupe	2.3-3.2	3.0-4.5	1.5	—	—	—	—	—
Carrot	9.0	13.5	—	—	—	—	—	—
Cauliflower	0.8-4.5	2.5-3.6	0.8	—	—	—	—	—
Clover	4.4-8.26	12.0	0.8	—	—	—	—	—
Collards	0.8	—	—	—	—	—	—	—
Conifers (Douglas fir, Red pine, Scotch pine, Norway spruce)	—	3.0	—	—	—	—	—	—
Corn (field)	1.0-2.4	1.5-1.7	3.2-3.6	—	—	—	—	—
Corn (sweet)	1.6-4.0	3.0-3.8	—	—	—	—	—	—
Cucumbers	1.4-3.2	3.0-4.5	1.6	—	—	—	—	—
Eggplant	6.0	9.0	—	—	—	—	—	—
Flax	2.0-3.8	3.0-5.4	—	—	—	—	—	—
Grasses	4.4-8.26	12.0	—	—	—	—	—	—
Kate	0.8	—	—	—	—	—	—	—
Legumes (small-seeded)	—	12.0	—	—	—	—	—	—
Lentils	2.54	—	—	—	—	—	—	—
Lescudera	4.4-8.26	12.0	—	—	—	—	—	—
Millet	4.31	—	—	—	—	—	—	—
Milo	4.0-6.0	6.0	1.6	—	—	—	—	—
Muskmelon	1.0-2.0	3.0	—	—	—	—	—	—
Mustard	0.8-4.5	0.6-20.0	—	—	—	—	—	—
Oats	1.2-5.6	1.2-5.0	0.8-4.0	—	—	—	—	—
Peas	1.6-3.6	1.6-3.8	—	—	—	—	—	—
Peppers	1.6-3.0	4.5	—	—	—	—	—	—
Radish	1.5	2.25	—	—	—	—	—	—
Rice	1.8-7.5	1.5-7.5	—	—	—	—	—	—
Rutabaga	4.5	6.75	—	—	—	—	—	—
Rye	1.2-3.2	1.2-3.0	—	—	—	—	—	—
Safflower	3.8811	0.6	—	—	—	—	—	—
Sesame	1.5-2.0	—	—	—	—	—	—	—
Sorghum	1.6-4.6	2.0-6.0	—	—	—	—	—	—
Soybeans	1.4-2.5	1.6-4.0	—	—	—	—	—	—
Spruce	3.6-6.0	9.0	—	—	—	—	—	—
Sugar beets	—	—	—	—	—	—	—	(see under Beets, sugar)
Sunflower	—	1.0	—	—	—	—	—	—
Swiss chard	6.8-12.0	18.0	—	—	—	—	—	—
Tomato	1.6	—	—	—	—	—	—	—
Trefoil	4.4-8.26	12.0	—	—	—	—	—	—
Turkey	0.8-3.0	4.5	—	—	—	—	—	—
Wheat	1.2-4.0	1.2-3.0	0.8-4.0	—	—	—	—	—

BEST AVAILABLE COPY

1 In combination with equal amounts of Botran.  
 2 In combination with equal amounts of Forder.  
 3 In combination with equal amounts of Maneo.  
 4 In combination with Maneo (Maneo to CAPTAN 50-W ratio 1.5:1).  
 5 In combination with equal amounts of PCNB.  
 6 In combination with 2.1 oz. actual Thiram/100 lbs.  
 7 In combination with Zineo (CAPTAN 50-W to Zineo ratio 22:21).  
 8 In combination with equal amounts of Thiram.  
 9 In combination with 0.6 oz. actual of Botran.  
 10 In combination with equal amounts of Maneo plus 0.4 to 0.6 oz. actual PCNB and 0.1 to 0.15 oz. actual 5-ethoxy-3-trifluoromethyl-1,2,4-triazole.  
 11 In combination with equal amounts of Carbazin.  
 12 In combination with WCR and Maneo (CAPTAN 50-W to WCR to Maneo ratio of 1:4:1).

**CONDITIONS OF SALE AND WARRANTY**

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

RQ Hazardous substance, solid,  
N.O.S., NA 9188, (Captan)

ORM-E

