

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive. Causes irreversible eye damage. Causes skin irritation. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Do not breathe spray mist. Wash thoroughly with copious water after handling. Remove contaminated clothing and wash before reuse. Wear goggles or face shield when applying. Wear natural rubber gloves when mixing or handling.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink promptly a large quantity of milk or egg whites, or at least a quart, if these are not available, drink large quantities of water. Get medical attention.

If In Eyes: Flush with plenty of water. Get immediate medical attention.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation continues.

NOTE TO PHYSICIAN: Product may cause damage to eyes. Irritation to the eyes if water is washed.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift. Do not apply to water. Do not apply to water by any means. Do not apply to water by any means.

STORAGE AND DISPOSAL

PROHIBITIONS AND RECOMMENDATIONS: Do not contaminate water, food, or feed by storage or disposal. Open burning is prohibited. Do not reuse empty container. Keep container closed when not in use.

STORAGE: Keep from freezing. Store above 32°F.

PESTICIDE DISPOSAL: Empty container, spray mixture or residue that cannot be used as directed by label instructions must be disposed of according to Federal, State, or local regulations under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and offer for recycling or reuse, if allowed, or dispose of in a sanitary landfill, or by other means if permitted by State and local authorities.

NOTICE: This warranty is given by the manufacturer of this product. It is given on the condition that the product is used in accordance with the directions on the label. The manufacturer does not warrant the product for use in any other manner. The manufacturer does not warrant the product for use in any other manner. The manufacturer does not warrant the product for use in any other manner. The manufacturer does not warrant the product for use in any other manner.



KELTHANE 4F FLOWABLE AGRICULTURAL MITICIDE

DIRECTIONS FOR USE

FOR AGRICULTURAL USE ONLY OR
ONLY FOR SALE TO, USE, AND
STORAGE BY SERVICEPERSONS

Before using this product read all precautionary statements including use precautions, storage and disposal instructions, use directions, limit of warranty and liability statements in these use directions and on the container label.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive. Causes irreversible eye damage. Causes skin irritation. Harmful if swallowed. Do not get in eyes. Do not wear contact lenses. Do not breathe spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Wear goggles or face shield when applying. Wear natural rubber gloves when mixing or handling.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Drink promptly a large quantity of milk, egg white, gelatin solution or if these are not available, drink large quantities of water. Get immediate medical attention.

If In Eyes: Flush with plenty of water. Get immediate medical attention.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation continues.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

GENERAL INFORMATION

KELTHANE 4F is a flowable formulation that gives high miticidal activity against most species of agricultural mites, including the following:

apple rust	peach silver
Bermuda	plum nursery
Brown	privet
Carmin	Schnee
citrus flat	two spotted
citrus red (purple)	strawberry
citrus rust	strawberry (Atlantic)
clover	tea and fruit
cycamen	tropical russet
Desert	tropical
dryberry	two spotted (red) (Desert)
European red	Willamette
McDaniel	yellow (orange)
Pacific	Yuma

KELTHANE is a specific miticide and will not kill bees and other beneficial insect predators when used as recommended.

KELTHANE 4F is believed to be compatible with most other commonly used insecticides and fungicides. It may be combined with some fungicides. KELTHANE may be applied in dust, granulate, ground sprays. For aerial sprays, use 3 to 10 gallons of water per acre and provide thorough coverage of all plant parts. Coverage of the underside of leaves is especially important. For further details, see specific recommendations in these use directions. For current State Agricultural Experiment Station spray schedules, contact State Extension Service Specialists for a list of contacts and publications dealing with mites.

USE PRECAUTIONS

- Read and observe all directions and precautions on labels of all pesticides used.
- Do not use KELTHANE 4F in combination with any other insecticide or fungicide unless specifically recommended.
- Do not use KELTHANE 4F in combination with any other insecticide or fungicide unless specifically recommended.
- Do not use KELTHANE 4F in combination with any other insecticide or fungicide unless specifically recommended.
- Do not use KELTHANE 4F in combination with any other insecticide or fungicide unless specifically recommended.

KELTHANE 4F FLOWABLE AGRICULTURAL MITICIDE

DIRECTIONS FOR USE

FOR AGRICULTURAL USE ONLY OR
ONLY FOR SALE TO, USE, AND
STORAGE BY SERVICEPERSONS

CROP RECOMMENDATIONS

ALFALFA (GROWN FOR SEED ONLY — USE RESTRICTED TO CALIFORNIA, IDAHO, NEVADA, OREGON AND WASHINGTON) — Carmin, Pacific, strawberry (Atlantic) and two spotted mites. Use 1 to 1 1/2 quarts of KELTHANE 4F per acre in sufficient water for thorough coverage. Apply when mites become active and repeat if necessary. Treated fields are not to be grazed, and crop residues and cuttings are not to be fed to livestock or baled for mulch use. The fields must be cleared by burning crop residues or by discing into the soil.

APPLES — To clean up brown and European red mite and to control apple rust. McDaniel, Pacific, Schnee, two spotted, Willamette and yellow (carpin) mites. Use 1/2 to 1 pint KELTHANE 4F per 100 gallons dilute spray (1 gallon maximum per acre). Make applications in the second or later cover sprays or whenever mite population appears. In areas where the preventive schedule is accepted practice, use 2 successive applications of 0.4 pint per 100 gallons dilute spray applied at 10 to 14 day intervals. Make thorough full coverage application. Do not apply within 7 days of harvest. To avoid possible injury, do not use glyodin fungicide with KELTHANE in late cover sprays. Before using on this crop, read USE PRECAUTIONS section in these use directions.

APRICOTS — Brown, European red, McDaniel, Pacific, peach silver, plum nursery, Schnee, two spotted, Willamette and yellow (carpin) mites. Use 1/2 to 1 pint KELTHANE 4F per 100 gallons dilute spray (6.4 pints maximum per acre). Apply in full coverage at petal fall to first cover or whenever mite population appears. Do not apply within 14 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

BEANS (dry, green, lima) — CALIFORNIA ONLY — Pacific, tropical, russet, tropical and two spotted mites. Use 3 pints KELTHANE 4F per acre. Apply at first signs of mite build up and repeat as necessary. Do not apply within 21 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

BEANS (dry, green, lima) — Pacific, tropical and two spotted mites. Use 1/2 to 1 pint KELTHANE 4F per acre. Apply at first signs of mite build up and repeat if necessary. Do not apply within 7 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

BLACKBERRIES — Brown, European red, McDaniel, Pacific, two spotted, Willamette and yellow (carpin) mites. Apply 6 to 12 quarts per acre just after bloom or whenever mites appear, and repeat as necessary. Do not apply within 2 days of harvest.

BOYSENBERRIES — Refer to recommendations for blackberries.

CHERRIES — Brown, European red, McDaniel, Pacific, peach silver, plum nursery, Schnee, two spotted, Willamette and yellow (carpin) mites. Use 1/2 to 1 pint KELTHANE 4F per 100 gallons dilute spray (6.4 pints maximum per acre). Apply in full coverage spray at petal fall, first cover or whenever mite population appears. For repeat applications, wait 30 days to avoid mite resistance. Do not apply within 14 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

CHESTNUTS — Brown, European red, McDaniel, Pacific, two spotted, Willamette and yellow (carpin) mites. Use 1 to 4 quarts per acre just after bloom or whenever mites appear, and repeat as necessary. Do not apply within 24 days of harvest.

CITRUS (grapefruit, kumquats, lemons, limes, oranges, tangelos, tangerines) — Citrus flat, citrus red (purple), citrus rust, citrus yellow, Yuma mite. Apply KELTHANE 4F in full coverage spray at petal fall, first cover or whenever mite population appears. For repeat applications, wait 30 days to avoid mite resistance. Do not apply within 14 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

CLOVER (GROWN FOR SEED ONLY — USE RESTRICTED TO CALIFORNIA, IDAHO, NEVADA, OREGON AND WASHINGTON) — Carmin, Pacific, strawberry (Atlantic) and two spotted mites.

