



751-164

FOR AGRICULTURAL USE ONLY OR
ONLY FOR SALE TO, USE, AND STORAGE BY SERVICEPERSONS

KELTHANE AF



FLOWABLE AGRICULTURAL MITICIDE



KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

PRECAUCIONAL USUARIO: Sustancia tóxica. Evite el contacto con la piel y los ojos. Evite la inhalación de la nebulización.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Precautions and Storage and Disposal Instructions on side panels and attached use directions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ACTIVE INGREDIENT

2,2-Dimethyl-4-chlorophenol 4.0%

INERT INGREDIENTS

96.0%

EPA Reg. No. 707-264

EPA Est. No. 707-PA-1

Manufactured by Rohm and Haas Company

NET CONTENTS
2.5 GALS
(9.46 L)

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury and skin irritation. Harmful if swallowed or inhaled. Do not get in eyes or on skin or clothing. Avoid breathing spray mist. Wear goggles or face shield when applying. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Flush with plenty of water. Get immediate medical attention.

If Inhaled: Move subject to fresh air.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation continues.

If Swallowed: Drink 2-3 glasses of water and induce vomiting by touching back of throat with finger. Avoid alcohol and fat bearing liquids like milk. Get medical attention. Do not induce vomiting if you are having difficulty swallowing or are unconscious.

NOTE TO PHYSICIAN: Adrenergic amines may cause myocardial instability in the presence of chlorinated hydrocarbons and blurred vision may be a prominent symptom of poisoning.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to the description and directions on the label. It does not warrant that the product will be effective when used in accordance with the label directions. ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE, handling, storage and disposal. Even if users are beyond the control of Rohm and Haas, it does not warrant that the product will be effective against the quantities resulting from but not limited to, weather conditions, presence of other materials, disease, pest resistance, or other products or failure to use label directions. In no case will Rohm and Haas or seller be held liable for consequential, special or indirect damages resulting from the handling, storage or use of this product.

549 1471 Rev. - Rohm and Haas - (C) - E. S. H. 4
Disc. 1 - File E

BEST DOCUMENT AVAILABLE

BEST DOCUMENT AVAILABLE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SEE ATTACHED PAMPHLET ON BACK PANEL FOR COMPLETE USE DIRECTIONS

RE-ENTRY AND WORKER PROTECTION STATEMENTS

Do not enter treated areas without protective clothing and sprays have dried. Because certain states may require more restrictive re-entry intervals than those specified on this product, check with your State Department of Agriculture for further information. Do not apply this product to water bodies as to direct contact with fish, shellfish, or other aquatic life. The area being treated must be completely unoccupied per se. SEE USE DIRECTIONS FOR ADDITIONAL RE-ENTRY STATEMENTS.

STORAGE AND DISPOSAL

PROHIBITIONS AND RECOMMENDATIONS Do not store in water, food, or feed. Do not store in areas where children, pets, or livestock are present. Do not use empty containers. See label for other restrictions.

STORAGE Keep from freezing. Store at 50° to 90°F.

PESTICIDE DISPOSAL Pesticide wastes are acutely toxic. Immediately clean spills with water. Do not get on skin or clothing. Do not breathe vapors. Do not eat, drink, or smoke while handling. Do not use household vacuum to clean up spills. Do not use household vacuum to clean up spills. Do not use household vacuum to clean up spills. Do not use household vacuum to clean up spills.

CONTAINER DISPOSAL Triple rinse, if necessary, with water. Empty containers must be disposed of in a safe manner. Do not reuse or refill containers.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED Avoid breathing vapors. Avoid contact with skin and clothing. If on skin, wash with plenty of water. If on clothing, remove clothing and wash thoroughly. If in eyes, flush with plenty of water for at least 15 minutes. If swallowed, do not induce vomiting. Seek medical attention immediately.

BEST DOCUMENT AVAILABLE

ROHM AND HAAS COMPANY
 PHILADELPHIA, PA 19106
 EPA REG. NO. 707-164
 © Copyright 1984 Rohm and Haas Company

ROHM AND HAAS COMPANY
 PHILADELPHIA, PA 19106
 EPA REG. NO. 707-164
 © Copyright 1984 Rohm and Haas Company

KELTHANE® 4F
FLOWABLE AGRICULTURAL MITICIDE
DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FOR AGRICULTURAL USE ONLY OR
 ONLY FOR SALE TO, USE, AND
 STORAGE BY SERVICE PERSONS**

Notice: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Precautions and Storage, and Disposal Instructions on this label. The Conditions of Sale and Warranty are not acceptable if you return the product unopened within thirty days of purchase to the place of purchase.

KELTHANE® 4F
FLOWABLE AGRICULTURAL MITICIDE
DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**FOR AGRICULTURAL USE ONLY OR
 ONLY FOR SALE TO, USE, AND
 STORAGE BY SERVICE PERSONS**

KEEP OUT OF REACH OF CHILDREN
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS
WARNING AVISO

PRECAUCION AL USUARIO: Sustitua no leeringes no use este producto hasta que le etiqueta haya sido expandida completamente. Causes substantial but temporary eye injury and skin irritation. Harmful if swallowed or inhaled. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist. Wear goggles or face shield when applying. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Flush with plenty of water. Get immediate medical attention.

If Inhaled: Move subject to fresh air.

If On Skin: Wash with plenty of soap and water. Get medical attention if irritation continues.

If Swallowed: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Avoid alcohol and fat based liquids like milk. Get medical attention. Do not induce vomit if you give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Adrenergic amines may cause myocardial arrhythmia in the presence of a preexisting cardiovascular disease. Sinus tachycardia may be a prominent symptom of poisoning.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label, only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or user are beyond the control of Rohm and Haas and any loss, such as crop injury, insect, vermin, or other unintended consequences, resulting from, but not limited to, weather or other conditions, or presence of other materials, diseases, pests, drift, etc., is the responsibility of the user. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

RE-ENTRY AND WORKER PROTECTION STATEMENTS

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

WARNING

(Insert area or field description) treated with Kelthane 4F on (insert date of application). Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure, Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Kelthane 4F and describe his condition. For further information, see the "Statement of Practical Treatment."

Oral warnings must include the same information that is in the written warnings.

GENERAL INFORMATION

KELTHANE 4F Miticide is a water formulation that gives the miticide a liquid spray application of agricultural mites, including the following:

- | | |
|-------------------|--------------------------|
| apple rust | peach silver |
| Bermuda | plum nursery |
| brus | prick |
| Carmine | Schoene |
| citrus flat | six spotted |
| citrus red spider | spruce |
| citrus rust | strawberry (Atlantic) |
| clover | Texas citrus |
| cyclopent | tomato russet |
| desert | Tropica |
| dratens | two spotted (red spider) |
| European red | Williams |
| Miticide | yellow scarab |
| Parasite | Yuma |

KELTHANE 4F Miticide is a specific miticide and will not kill beneficial insect predators when used as recommended.

KELTHANE 4F Miticide may be applied in dilute or concentrated ground sprays. For area sprays use 3 to 10 gallons of water per acre and provide thorough coverage of all plant parts. Coverage on the underside of leaves is especially important. For further details see spray recommendations in the use directions. For complete current State Agricultural Experiment Station spray schedules, State Extension Service, State University, address details on crop parts to be sprayed and use.

BEST DOCUMENT AVAILABLE

PESTICIDE COMPATIBILITY

KELTHANE 4F miticide is believed to be compatible with most commonly used insecticides and fungicides with the exception of the following lime and other highly alkaline materials: Cypres fungicide and AarEE growth regulator. In addition, KELTHANE 4F should not be used with Glycor fungicide in late cover sprays in apples.

USE PRECAUTIONS

- Read and observe all label directions including use precautions before using.
- Use only for recommended purposes and at recommended rates.
- Do not apply to any crop within two weeks following a fall or dust or spray.
- Do not apply to apples, apricots, cherries, crab apples, figs, grapes, nectarines, peaches, plums or quince after fruit formation in California, Oregon or Washington. Use KELTHANE 35 Miticide instead.
- Do not apply to prunes or strawberries after fruit formation in California. Use KELTHANE 35 Miticide instead.

CROP RECOMMENDATIONS

ALFALFA (GROWN FOR SEED ONLY — USE RESTRICTED TO CALIFORNIA, IDAHO, NEVADA, OREGON AND WASHINGTON) — For Carmine, Pacific strawberry (Atlantic) and two spotted mites. Use 2 to 3 pints KELTHANE 4F miticide per acre in sufficient water for thorough coverage. Apply when mites appear and repeat if necessary. Treated fields are not to be grazed and crop residues and cuttings are not to be fed to livestock or baled for sale. The fields must be cleared by burning crop residues or by disking into the soil.

APPLES — Apple rust, clover, European red, McDanie, Pacific, Schoene, two spotted, Willamette and yellow (carpini) mites.

Dosage — For dilute sprays use 1/2 to 1 pint KELTHANE 4F miticide per 100 gallons. Use the 1 pint rate per 100 gallons spray for large trees or heavy mite populations. KELTHANE 4F miticide is recommended at a minimum rate of 3 pints per acre in the dilute or concentrate sprays. The maximum rate for KELTHANE 4F miticide is 8 pints per acre. For concentrate ground applications use 3 to 5 pints of KELTHANE 4F miticide per acre depending on tree size and number of trees per acre, foliage density and mite pressure. When heavy mite populations are encountered, two successive sprays applied 7 to 10 days apart are recommended.

Application — Dilute applications should be made in sufficient volume to provide thorough coverage of all parts of the tree. Concentrate applications are recommended at 1/2 gallon of water per acre or greater with sufficient volume to provide thorough coverage of all tree parts. The amount of KELTHANE 4F miticide applied per acre should be the same whether dilute or concentrate sprays are used.

Timing of Application — Applications are recommended in the spring or later cover sprays at the first signs of mite buildup. Repeat applications should be made as necessary for adequate seasonal control.

USE RESTRICTIONS

- Do not apply KELTHANE 4F miticide to bare varieties before the second cover spray.
- Applications of KELTHANE 4F miticide on deer and wild turkey should not exceed 4 pints per acre per spray.
- Do not apply KELTHANE 4F miticide within 7 days of harvest.
- Refer to label section on pesticide compatibility.

APRICOTS — Brown, European red, McDanie, Pacific peach silver plum nursery, Schoene, two spotted, Willamette and yellow (carpini) mites. Use 1/2 to 1 pint per 100 gallons dilute spray. 4 pints maximum per acre. Apply in full coverage spray at petal fall, first cover or whenever mite population appears. Do not apply within 14 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

BEANS (dry green lima) — CALIFORNIA ONLY — Pacific tomato russet, tropical and two spotted mites. Use 3 pints per acre. Apply at first signs of mite buildup and repeat as necessary. Do not apply within 21 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

BEANS (dry green lima) — Pacific, tropical and two spotted mites. Use 1/2 to 1 pint per acre. Apply at first signs of mite buildup and repeat if necessary. Do not apply within 7 days of harvest. Do not feed treated vines (or crop residues) to meat or dairy animals.

BLACKBERRIES — Brown, European red, McDanie, Pacific two spotted, Willamette and yellow (carpini) mites. Apply 0.6 to 1.2 quarts per acre just after bloom or whenever mites appear and repeat as necessary. Do not apply within 2 days of harvest.

BOYSENBERRIES — Refer to recommendations for blackberries.

CHERRIES — Brown, European red, McDanie, Pacific peach silver plum nursery, Schoene, two spotted, Willamette and yellow (carpini) mites. Use 1/2 to 1 pint per 100 gallons dilute spray (4 pints maximum per acre). Apply in full coverage spray at petal fall, first cover or whenever mite population appears. For repeat application, wait 30 days to avoid excessive residues, but do not apply within 7 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

CHESTNUTS — Brown, European red, McDanie, Pacific two spotted, Willamette and yellow (carpini) mites. Use 1.6 to 4 quarts per acre just after bloom or whenever mites appear and repeat as necessary. Do not apply within 14 days of harvest.

CITRUS (grapefruit, kumquats, lemons, limes, oranges, tangelos, tangerines) — Citrus flat, citrus red, citrus rust, six spotted, Texas citrus and Yuma mites. Apply in full coverage sprays using 1 pint per 100 gallons of dilute spray (2 gallons maximum per acre). If concentrate sprays are used, adjust dosage accordingly. Apply when mites appear in threatening numbers and repeat as required for satisfactory control. Do not apply within 7 days of harvest.

CLOVER (GROWN FOR SEED ONLY — USE RESTRICTED TO CALIFORNIA, IDAHO, NEVADA, OREGON AND WASHINGTON) — Refer to recommendations for alfalfa.

COTTON — Desert, Pacific and two spotted mites. Use 1 to 1.5 quarts per acre depending on size of plant and density of foliage. Apply in sufficient water to provide thorough coverage. Begin application when mites are first noted and repeat if necessary, and do not apply within 14 days of harvest. Do not feed cotton stalks or trash to meat or dairy animals.

CRABAPPLES — Refer to recommendations for apples. Before using on this crop, read USE PRECAUTIONS section in these use directions.

CUCUMBERS — European and two spotted mites. Use 1/2 to 1 pint KELTHANE 4F miticide per acre in sufficient water to provide thorough coverage of all plant parts. Adjust spray nozzle to give adequate coverage to underside of leaves. Use the higher maximum rate of 1 1/2 pints when vines begin to overgrow. Make applications at 7 to 10 day intervals and repeat if necessary. Do not feed treated vines or crop residues to meat or dairy animals. Do not apply KELTHANE 4F miticide within 2 days of harvest.

DEWBERRIES — Refer to recommendations for blackberries.

FIGS — Brown, European red, McDanie, Pacific peach silver plum nursery, Schoene, two spotted, Willamette and yellow (carpini) mites. Use 1/2 to 1 pint per 100 gallons dilute spray. 4 pints maximum per acre. Apply in full coverage spray at petal fall, first cover or whenever mite population appears. KELTHANE 4F Miticide may cause fruit splitting of figs and should be used with caution. For repeat application, wait 30 days to avoid excessive residues but do not apply within 7 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

FILBERTS — Refer to recommendations for chestnuts.

GRAPES — European red, McDanie, Pacific two spotted, McDanie and Willamette mites. Use 1/2 to 1 pint KELTHANE 4F miticide per acre in sufficient water to provide thorough coverage. Adjust spray nozzle to give adequate coverage to underside of leaves. Make applications in full coverage spray with 10 to 15 gallons per acre and repeat as necessary. Do not apply within 7 days of harvest.

BEST DOCUMENT AVAILABLE

HICKORY NUTS — Refer to recommendations for chestnuts.

HOPS — For Pacific and two spotted mites. Use 2 to 2 1/2 pints KETHANE 4F miticide per acre in sufficient water to provide full coverage to all plant parts. Use lower recommended dosage rate when plants are small and increase dosage rate with development of vines. Apply at first sign of mite build up and repeat when necessary. Do not apply KETHANE 4F miticide within 7 days of harvest. Do not feed treated vines or crop residues to meat or dairy animals.

LOGANBERRIES — Refer to recommendations for blackberries.

MELONS — Refer to recommendations for cucumbers.

MINT (peppermint and spearmint for oil production) — For two spotted mites. Use 1 1/2 to 2 pints KETHANE 4F miticide per acre in sufficient water for thorough coverage. Make application when mites become active. Do not apply within 30 days of harvest and do not make more than one application per crop per season. Do not feed treated fresh or spent hay to livestock.

NECTARINES — Brown, European red, McDaniel, Pacific peach silver plum nursery, Schone, two spotted, Willamette and yellow (carpenter) mites. Use 1/2 to 1 pint per 100 gallons dilute spray (6 1/4 pints maximum per acre). Apply in full coverage spray at petal fall to first cover or whenever mite population appears. Do not apply within 14 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

OLALLIEBERRIES — Refer to recommendations for blackberries.

PEACHES — Brown, European red, McDaniel, Pacific peach silver plum nursery, Schone, two spotted, Willamette and yellow (carpenter) mites. Use 1/2 to 1 pint per 100 gallons dilute spray (6 1/4 pints maximum per acre). Apply in full coverage spray at petal fall to first cover or whenever mite population appears. Do not apply within 14 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

PEARS — Refer to recommendations for apples. KETHANE 4F Miticide may cause fruit splits on pears under certain conditions and should be used at lower dosages and with caution on this crop.

PECANS — Refer to recommendations for chestnuts.

PEPPERS — Pacific tomato, russet, tropical and two spotted mites. Apply 0.6 to 1.6 pints per acre at first signs of mite build up and repeat as necessary. Do not feed crop residues to dairy or meat animals. Do not apply within 7 days of harvest.

PLUMS — Brown, European red, McDaniel, Pacific peach silver plum nursery, Schone, two spotted, Willamette and yellow (carpenter) mites. Use 1/2 to 1 pint per 100 gallons dilute spray (6 1/4 pints maximum per acre). Apply in full coverage spray at petal fall to first cover or whenever mite population appears. For repeat application wait 30 days to avoid excessive residue build up. Do not apply within 7 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

PRUNES — Brown, European red, McDaniel, Pacific peach silver plum nursery, Schone, two spotted, Willamette and yellow (carpenter) mites. Use 1/2 to 1 pint per 100 gallons dilute spray (6 1/4 pints maximum per acre). Apply in full coverage spray at petal fall to first cover or whenever mite population appears. For repeat application wait 30 days to avoid excessive residue build up. Do not apply within 7 days of harvest. Before using on this crop, read USE PRECAUTIONS section in these use directions.

PUMPKINS — Refer to recommendations for cucumbers.

QUINCE — Refer to recommendations for apples. Before using on this crop, read USE PRECAUTIONS section in these use directions.

RASPBERRIES — Refer to recommendations for blackberries.

SQUASH (winter and summer) — Refer to recommendations for cucumbers.

STRAWBERRIES — Six spotted, straw, yellow, Atlantic and two spotted mites. Use 1 quart per acre. Apply when mites appear and repeat as necessary. Do not apply within 7 days of harvest. Do not feed treated fruit to dairy animals. Before using on this crop, read USE PRECAUTIONS section in these use directions.

TOMATOES — Pacific tomato, russet, tropical and two spotted mites. Apply 0.6 to 1.6 pints per acre at first signs of mite build up and repeat as necessary. Do not feed crop residues to dairy or meat animals. Do not apply within 7 days of harvest.

WALNUTS — Brown, European red, McDaniel, Pacific two spotted, Willamette and yellow (carpenter) mites. Use a full coverage spray of 1/2 to 1 pint per 100 gallons dilute spray (1 gallon maximum per acre). Apply just after bloom or whenever mites appear and repeat if necessary. Do not apply within 14 days of harvest.

OTHER USES

LAWN AND TURF GRASSES — Bermuda, brown and clover mites. Use 0.6 to 0.8 pint per 100 gallons (1 teaspoon per gallon). Apply 2 1/2 gallons dilute spray per 1000 square feet. Make application when mites appear, covering the entire lawn area with a thorough spray, and repeat in 10 to 14 days if necessary.

FLOWERING PLANTS (including African violets, chrysanthemums, sweet peas) — European red and two spotted mites. Use 0.6 to 0.8 pint per 100 gallons of water. Apply full coverage spray at first sign of mites and repeat as necessary.

ORCHIDS — European red and two spotted mites. Use 0.6 to 0.8 pint per 100 gallons of water. Apply full coverage spray at first sign of mites and repeat as necessary.

SHRUBS, TREES, VINES (ornamental) — DO NOT APPLY TO CANAETH, JUNIPER OR CHINESE HOLLY. — Clover, European red, spruce and two spotted mites. Use 0.6 to 0.8 pint per 100 gallons of water. Apply full coverage spray at first sign of mites and repeat as necessary.

DOMESTIC DWELLINGS, MEDICAL FACILITIES, SCHOOLS — Clover mites. Use 0.6 to 0.8 pint per 100 gallons of water. Thoroughly spray only the outside walls, foundations and windowsills, as well as plants and lawns at base of infested buildings.

GREENHOUSE PLANTS (including carnations, chrysanthemums, roses) — Spider mites. Use 4 fluid ounces per 50,000 cubic feet. Apply by fog generator. Close all doors, windows and ventilators. Turn off all misting systems, fans and sulfur pots. Be sure all blossoms and foliage are dry. Adjust inside temperature to 70 to 80°F. Keep greenhouse tightly closed for 1 1/2 hours following treatment. Ventilate thoroughly for at least 1 hour before entering.

NOTE — KETHANE 4F Miticide may be used without any spreader sticker. When necessary to improve the spray coverage on foliage, dilute with water and to smooth out spray deposits on maturing fruit, TRITON® B 1956® is recommended. Other suitable pesticide spreader stickers may also be used except in situations where TRITON® B 1956 is recommended as the only acceptable spreader sticker.

STORAGE AND DISPOSAL

PROHIBITIONS AND RECOMMENDATIONS — Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not re-use empty container. Keep container closed when not in use.

STORAGE — Keep from freezing. Store above 32°F (10°C).

Pesticide Disposal — Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agent, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Disposal — Triple rinse (or equivalent). Puncture and dispose of in a sanitary landfill, or incinerate, or if allowed by state and local authority, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED — ventilate area. Avoid breathing vapors. Use MSHA NIOSH self contained breathing apparatus or air respirator for large spills in confined areas. Dike the spill with inert material (sand, earth, etc.) and transfer the liquid or solid diking material to separate container for recovery or disposal. Remove contaminated clothing promptly and wash exposed skin areas with soap and water. Wash clothing before re-use. Keep spill out of all sewers and open bodies of water.

ALABAMA: Alabama Department of Conservation, 1000 North University Blvd., Montgomery, AL 36102. ALASKA: Alaska Department of Fish and Game, 1225 West 7th Ave., Anchorage, AK 99515. ARIZONA: Arizona Department of Agriculture, 1500 North Central Ave., Phoenix, AZ 85004. ARKANSAS: Arkansas Department of Agriculture, 1000 North Main St., Little Rock, AR 72201. CALIFORNIA: California Department of Pesticide Regulation, 2800 California Street, Sacramento, CA 95833. COLORADO: Colorado Department of Agriculture, 1700 North Lincoln Ave., Denver, CO 80202. CONNECTICUT: Connecticut Department of Agriculture, 1000 Main St., Hartford, CT 06103. DELAWARE: Delaware Department of Agriculture, 1000 North Main St., Dover, DE 19901. DISTRICT OF COLUMBIA: District of Columbia Department of the Environment, 1400 Constitution Ave., NW, Washington, DC 20004. FLORIDA: Florida Department of Agriculture and Consumer Services, 1900 North 17th Ave., Tallahassee, FL 32304. GEORGIA: Georgia Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. HAWAII: Hawaii Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. ILLINOIS: Illinois Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. INDIANA: Indiana Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. IOWA: Iowa Department of Agriculture and Land Stewardship, 1000 North 17th Ave., Tallahassee, FL 32304. KANSAS: Kansas Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. KENTUCKY: Kentucky Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. LOUISIANA: Louisiana Department of Agriculture and Forestry, 1000 North 17th Ave., Tallahassee, FL 32304. MAINE: Maine Department of Agriculture, Conservation and Forestry, 1000 North 17th Ave., Tallahassee, FL 32304. MARYLAND: Maryland Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. MASSACHUSETTS: Massachusetts Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. MICHIGAN: Michigan Department of Agriculture and Rural Affairs, 1000 North 17th Ave., Tallahassee, FL 32304. MINNESOTA: Minnesota Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. MISSISSIPPI: Mississippi Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. MISSOURI: Missouri Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. MONTANA: Montana Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. NEBRASKA: Nebraska Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. NEVADA: Nevada Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. NEW HAMPSHIRE: New Hampshire Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. NEW JERSEY: New Jersey Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. NEW MEXICO: New Mexico Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. NEW YORK: New York Department of Agriculture and Markets, 1000 North 17th Ave., Tallahassee, FL 32304. NORTH CAROLINA: North Carolina Department of Agriculture and Consumer Services, 1000 North 17th Ave., Tallahassee, FL 32304. NORTH DAKOTA: North Dakota Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. OHIO: Ohio Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. OKLAHOMA: Oklahoma Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. OREGON: Oregon Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. PENNSYLVANIA: Pennsylvania Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. RHODE ISLAND: Rhode Island Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. SOUTH CAROLINA: South Carolina Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. SOUTH DAKOTA: South Dakota Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. TENNESSEE: Tennessee Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. TEXAS: Texas Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. UTAH: Utah Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. VERMONT: Vermont Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. VIRGINIA: Virginia Department of Agriculture and Forestry, 1000 North 17th Ave., Tallahassee, FL 32304. WASHINGTON: Washington Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. WEST VIRGINIA: West Virginia Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. WISCONSIN: Wisconsin Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304. WYOMING: Wyoming Department of Agriculture, 1000 North 17th Ave., Tallahassee, FL 32304.