

Container Label

**ACCEPTED**

NOV - 5 1985

U.S. Environmental Protection Agency  
 Office of Pesticide Registration  
 400 ...  
 EPA Reg. No. ...


# STAMPEDE CM

HERBICIDE

KEEP OUT OF REACH OF CHILDREN

## CAUTION

BEFORE USING THIS PRODUCT, READ THE ENTIRE PRECAUTIONARY STATEMENTS, CONDITIONS OF SALE AND WARRANTY, DIRECTIONS FOR USE, USE RESTRICTIONS, AND STORAGE AND DISPOSAL INSTRUCTIONS ON SIDE PANELS AND ATTACHED USE DIRECTIONS. IF THE CONDITIONS OF SALE AND WARRANTY ARE NOT ACCEPTABLE RETURN THE PRODUCT UNOPENED WITHIN THIRTY DAYS OF PURCHASE TO THE PLACE OF PURCHASE.



**ROHM  
AND  
HAAS**  
PHILADELPHIA, PA. 19105

ACTIVE INGREDIENTS.....48%  
 propyl 1,3,4-Dichloropropionamide.....35%\*  
 2-Methyl-4-chlorophenoxyacetic acid,  
 isobutyl ester.....15%\*\*

INERT INGREDIENTS.....52%  
 100%

\*Equivalent to 5lbs active ingredient per gallon

\*\*Equivalent to 1.4lbs per gallon of

2-Methyl-4-chlorophenoxyacetic acid, isobutyl ester

(0.85 lbs 2-Methyl-4-chlorophenoxyacetic acid equivalent)

EPA REG NO. 707-182


EPA EST NO. 4990-DV-2

Copyright 1985 Rohm and Haas Company

STAMPEDE is a trademark of  
 Rohm and Haas Company

NET CONTENTS:

2.5 GALS  
 (9.46 L)



**BEST AVAILABLE COPY**

### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Remove and wash clothing before re-use.

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES** — Flush eyes with plenty of water for at least 15 minutes. Call a physician if irritation persists.

**IF INHALED** — Move subject to fresh air.

**IF ON SKIN** — Wash affected area with soap and water.

**IF SWALLOWED** — Dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN** — Emesis is recommended.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Keep out of lakes, ponds or streams. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

### PHYSICAL AND CHEMICAL HAZARDS

#### COMBUSTIBLE

Do not use, pour, or store near heat or open flame.

### CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather, soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

### RE-ENTRY AND WORKER PROTECTION STATEMENTS

Do not enter treated areas without protective clothing until sprays have dried. Because certain states have restrictive re-entry intervals for various crops and products, consult your State Department of Agriculture for further information. Do not apply this product directly or through drift, expose workers to the area being treated must be vacated by persons.

### STORAGE AND DISPOSAL

**STORAGE** — Do not store at temperature below 15°F and crystals form, warm to room temperature, periodically shaking container to reconstitute.

**PESTICIDE DISPOSAL** — Do not contaminate water by storage or disposal. Pesticide wastes are a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or Waste representatives at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL** — Triple rinse (or equivalent) and dispose of in a sanitary landfill, or if allowed by state and local authorities, by other means that will prevent escape and stay out of smoke.

**STEPS TO BE TAKEN IN CASE MATERIAL IS SPILLED** — Eliminate ignition sources. Ventilate area to avoid breathing vapors. Use MSHA/NIOSH self-contained breathing apparatus or air mask for large spills in confined spaces. For small spills, absorb spill with inert material (sand, earth, fuller's earth, etc.) and, if appropriate, transfer the liquid and solid to separate containers for recovery or disposal. Do not discharge into sewers and open bodies of water. Refer to the Statements.

MADE IN U.S.A.

**NOTICE:** Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

## KEEP OUT OF REACH OF CHILDREN CAUTION

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes or clothing. Remove and wash clothing before re-use.

### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES** — Flush eyes with plenty of water for at least 15 minutes. Call a physician if irritation persists.

**IF INHALED** — Move subject to fresh air.

**IF ON SKIN** — Wash affected area with soap and water.

**IF SWALLOWED** — Dilute by giving 2 glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

**NOTE TO PHYSICIAN** — Emesis is recommended.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Keep out of lakes, ponds or streams. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

### PHYSICAL AND CHEMICAL HAZARDS

#### COMBUSTIBLE

Do not use, pour, or store near heat or open flame.

### CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### RE-ENTRY AND WORKER PROTECTION STATEMENTS

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift, expose workers or other persons. The area being treated must be vacated by unprotected persons.

### GENERAL INFORMATION

STAMPEDE CM is a selective postemergence herbicide for the control of green and yellow foxtail grass, wild millet, pigeongrass and specific broadleaf weeds in spring barley, durum and hard red spring wheat in North Dakota, South Dakota, Minnesota and Montana. When used according to label directions the product will provide effective control of the following weeds:

#### PIGEONGRASS

GREEN FOXTAIL

YELLOW FOXTAIL

WILD BUCKWHEAT

PEDROOT PIGWEED

PROSTRATE PIGWEED

LAMBSQUARTERS

WILD MUSTARD

KOCHIA

*Setaria viridis*

*Setaria lutescens*

*Polygonum convolvulus*

*Amaranthus retroflexus*

*Amaranthus bi-lobes*

*Chenopodium album*

*Brassica kaber*

*Kochia scoparia*

STAMPEDE CM herbicide is an emulsifiable concentrate containing 4.4 pounds active ingredient per U.S. gallon. The product kills susceptible weeds mainly by direct contact action. For this reason, thorough coverage of emerged weeds is essential for optimum results. Only weeds that have emerged and are exposed at time of application will be controlled. The product has no preemergence or residual herbicidal activity. STAMPEDE CM is most effective if applied when pigeongrass and susceptible broadleaf weeds are small and growing actively under favorable soil moisture and weather conditions. Early weed control removes competition, saves moisture and generally contributes to increased yields.

### DOSAGE RATE RECOMMENDATIONS — SPRING BARLEY, DURUM AND HARD RED SPRING WHEAT

For control of pigeongrass, wild buckwheat and other labeled broadleaf weeds growing under favorable soil moisture conditions, apply STAMPEDE CM herbicide in a single broadcast application at the rate of 2 to 3 pints per acre. Do not apply STAMPEDE CM at rates higher than 3 pints per acre.

**APPLICATION TIMING**

**WEED CONTROL** — For optimum weed control and crop tolerance, the importance of early spray timing cannot be overemphasized. STAMPEDE CM herbicide is most effective in controlling pigeongrass if spray applications are made when the majority of the emerged pigeongrass are in the 2 to 3 LEAF STAGE. Effectiveness of STAMPEDE CM in controlling pigeongrass declines rapidly as the 4th leaf begins to emerge. Since pigeongrass at these early stages is often less than one inch tall, close inspection of the field is essential to determine correct application timing. Fields should be checked frequently for emerging pigeongrass starting 2 to 2½ weeks after planting. The best time to apply STAMPEDE CM often will be 10 to 17 days after crop emergence when the grain is in the 3 to 4 LEAF STAGE.

For effective control of labeled weeds, applications of STAMPEDE CM herbicide must be made to actively growing weeds in the seedling stages listed below.

Weeds Controlled	Seedling Stage*
Pigeongrass	1 to 3 leaf
Buckwheat (wild)	1 to 4 leaf
Kochia	1 to 4 leaf
Mustard (wild)	1 to 4 leaf
Lambsquarters	1 to 4 leaf
Pigweed (prostrate)	1 to 4 leaf
Pigweed (redroot)	1 to 4 leaf

\*Refers to weed seedling stages for control. Pigeongrass greater than the 3 leaf stage or listed broadleaf weeds beyond the 4 leaf stage will not be controlled.

**CROP TOLERANCE**

STAMPEDE CM herbicide causes temporary yellowing or tip browning in the grain crop 2 to 5 days after application. The effects are temporary and usually will disappear 10 to 14 days after application. New leaves will have normal green color. Application of STAMPEDE CM under adverse growing conditions may result in greater crop injury and slower recovery. Care should be taken not to spray STAMPEDE CM on grain crops that are under growth stress caused by drought, flooding, excess soil salts, hail damage, or extreme temperatures. Do not apply STAMPEDE CM if frost is expected within 24 hours or when temperatures are above 85°F, especially with drying winds.

Applications of STAMPEDE CM herbicide should be made only to actively growing (healthy) grain crops in the seedling stages recommended below:

Crop	Seedling Stage*
Hard Red Spring Wheat	2 to 5 leaf
Durum Wheat	2 to 4 leaf
Barley	2 to 4 leaf

\*Refers to crop seedling stages for best tolerance. To avoid crop injury, do not apply STAMPEDE CM herbicide to hard red spring wheat beyond the 5 leaf stage or to spring barley or durum wheat beyond the 4 leaf stage.

**MIXING AND EQUIPMENT**

STAMPEDE CM herbicide is an emulsifiable concentrate that mixes readily with water. With agitator running, pour the measured quantity of STAMPEDE CM into partially filled spray tank and complete filling. Spray water should be clean and 50°F or warmer when mixed. STAMPEDE CM may be applied with conventional low pressure herbicide sprayers.

**Field Sprayers:** Spra-Coupe, tractor drawn or truck-mounted ground sprayers.

Use flat fan nozzles of sufficient size to provide a minimum of 10 to 12 gallons volume per acre at a pressure of not less than 40 psi. Recommended ground speeds are 5 to 8 mph (1st or 2nd gears only) for Spra-Coupe, 4 to 5 mph for tractor drawn sprayers and 8 to 10 mph for truck-mounted sprayers. Hollow cone or floodjet nozzles are not recommended on these sprayers.

Floaters: Big A, Terragator, etc.

Floaters equipped with floodjet nozzles are satisfactory for STAMPEDE CM herbicide application. A minimum of 15 to 20 gallons spray per acre is recommended because of the coarser droplet size produced by the floodjet nozzles. The nozzles should be no more than 60" apart and of such a size to provide the recommended volume at normal operating speeds of 12 to 15 mph. Pressure should be sufficient to produce proper overlap of spray patterns between nozzles to avoid skips.

If floaters are equipped with other nozzles use a minimum pressure of 40 psi to insure proper overlap of spray patterns between nozzles.

**AERIAL APPLICATION**

STAMPEDE CM herbicide can be applied either by fixed wing aircraft or helicopter. Make applications at the recommended dosage and timing in a minimum of 5 to 10 gallons of water per acre. Aircraft should be equipped with a well designed spray distribution system that is adjusted and operated to provide a uniform pattern of medium sized droplets (400 micron size) distributed over an optimum width swath. Only hollow cone nozzles with cores are recommended for STAMPEDE CM aerial application. Nozzles should be angled 0° to 20° downward and to the rear, depending on air speed. Nozzles should not be placed within 2 to 3 feet of the wing tips in fixed wing aircraft and extras should be placed on the right side of the fuselage. Use a spray pump of at least 1½ to 2 inches in diameter and operate at 20 to 50 psi. Application should be made with boom height 6 to 10 feet above crop. The optimum or effective spray swath width depends on operating conditions and type of aircraft being used. For uniform spray coverage with fixed wing aircraft do not exceed a spray swath width 10% greater than the wingspan or the length of the boom in helicopters. Measure the swaths accurately for flagging.

**CAUTION: DO NOT APPLY STAMPEDE CM HERBICIDE IN AERIAL APPLICATIONS WITHIN 600 FEET OF ANY PERMANENT BODY OF WATER OR WHEN WIND SPEED EXCEEDS 10 MPH.**

**CALIBRATION**

Sprayers should be checked carefully for pressure, ground speed, and uniformity of spray pattern, and calibrated accurately for spray volume. Replace any worn nozzles.

**CLEANING SPRAYERS**

Drain and flush spray tank and lines thoroughly before mixing STAMPEDE CM herbicide. Repeat this procedure at the end of each day's spraying and before changing to or from another pesticide. This cleaning operation of tank and lines is especially important if insecticides such as SEVIN® or any of the organophosphorus insecticides are used in the sprayer prior to or after STAMPEDE CM. Interaction of these insecticides and STAMPEDE CM can injure the crop severely. See COMPATIBILITY WITH OTHER CHEMICALS section.

**EFFECT OF CLIMATIC CONDITIONS AND CULTURAL PRACTICES ON WEED CONTROL**

**Field and Seedbed Preparation**

Fields under conventional tillage should have well prepared seedbeds, free of large clods, to encourage more uniform and rapid emergence of the crop, pigeongrass, and broadleaf weeds. This permits better timing of sprays. In no-till grain fields, pigeongrass and other weeds may emerge rapidly and very thickly because of the firm seedbed. Monitoring of no-till fields should begin early to assure proper timing of sprays.

**Soil Moisture and Rainfall**

Weed control with STAMPEDE CM herbicide is strongly influenced by soil moisture. Weed control and crop tolerance is best when adequate moisture is available for active weed and crop growth. The performance of STAMPEDE CM is reduced substantially during periods of prolonged dry weather. STAMPEDE CM IS NOT RECOMMENDED FOR USE WHEN SOIL MOISTURE LEVELS ARE DEEPER THAN 1 - INCHES FROM THE SOIL SURFACE AND ACTIVE PLANT GROWTH STOPS.

STAMPEDE CM herbicide enters the leaf rapidly. Spraying of weeds at the correct stage should not be postponed even if showery weather is expected; however, a heavy rain of 1 inch or more, within 4 hours after application, may reduce control. Fields may be sprayed when the plants are dry or wet with dew or rain.

**TEMPERATURE**

Providing soil moisture is not a limiting factor, the herbicidal activity of STAMPEDE CM herbicide improves as daily maximum temperatures increase. Best weed control occurs when the daily maximum temperatures are 65°F and above. Do not apply STAMPEDE CM when daily maximum temperatures will remain below 50°F or when they are expected to go higher than 85°F.

**COMPATIBILITY WITH OTHER CHEMICALS**

**Foliar Applied Chemicals**

If another herbicide is required, for example, a postemergence wild oat herbicide, a 3 day interval should be allowed between the application of STAMPEDE CM herbicide and the other herbicide.

Severe injury or kill of spring barley, durum or hard red spring wheat plants may result from tank mix or separate applications of STAMPEDE CM herbicide and certain insecticides. Grain crops that have been or will be treated with STAMPEDE CM should not be treated with carbamate insecticides such as carbaryl (Sevin, etc.), methomyl (Lannate®, Nudrin®, etc.) or organophosphorus insecticides such as methyl parathion, Guthion®, etc.

If a foliar insecticide is necessary, malathion or toxaphene can be safely applied 14 days before or after the STAMPEDE CM herbicide treatment.

**Soil Applied Chemicals**

Do not spray spring barley, durum or hard red spring wheat with STAMPEDE CM herbicide if the field was treated with soil-applied systemic organophosphorus insecticides (Disyston®, Thimet®, Counter®, etc.) the previous year as used for potatoes or sugarbeets.

Do not spray STAMPEDE CM herbicide on spring barley, durum or hard red spring wheat that has been treated at planting time with soil-applied systemic insecticides for Hessian Fly control, e.g., Thimet, Disyston, Furadan®, etc.

Grain crops protected with maneb/lindane seed dressings may be treated with STAMPEDE CM herbicide.

**USE RESTRICTIONS**

- Do not apply this product to any crops other than spring barley, durum or hard red spring wheat.
- Do not mix or apply this product with any other pesticide, spray adjuvant or with fertilizer except as specifically recommended on this label.
- Do not apply this product under windy conditions that will allow spray drift to adjacent susceptible crops such as sunflowers, soybeans, sugarbeets, potatoes, forage legumes, gardens, orchards, and shelter belts.
- DO NOT GRAZE TREATED CROP OR CUT FOR GREEN CHOP FEED.

**STORAGE AND DISPOSAL**

**STORAGE** — Do not store at temperatures below 15°F. If stored below 15°F and crystals form, warm to 60°F for 24 hours, periodically shaking container to reconstitute.

**PESTICIDE DISPOSAL** — Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL** — Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED** — Eliminate ignition sources. Ventilate area. Avoid breathing vapors. Use MSHA, NIOSH self-contained breathing apparatus or air mask for large spills in confined areas. Dike the spill with inert material (such as earth, fuller's earth, etc.) and if appropriate, transfer the liquid and solid diking material to separate containers for recovery or disposal. Keep spill out of all sewers and open bodies of water. Refer to Precautionary Statements.

STAMPEDE™ is a trademark of Rohm and Haas Company.  
COUNTER® and THIMET® are registered trademarks of American Cyanamid Co.  
DISYSTON® is a registered trademark of Bayer AG.  
FURADAN® is a registered trademark of FMC Corp.  
GUTHION® is a registered trademark of Mobay Chemical Corp.  
LANNATE® is a registered trademark of E. I. duPont de Nemours & Co.  
NUDRIN® is a registered trademark of Shell Chemical.  
SEVIN® is a registered trademark of Union Carbide Corp.

**BEST AVAILABLE COPY**

10012-B1

