

Please read instructions on reverse before completing form.

Form Approved. OMS No. 2070-0060. Approval expires 11-30-93

(A) EPA	United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460 Application for Pesticide:	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number <div style="border: 1px solid black; padding: 5px; font-size: 1.2em;">197352</div>
------------------------------	---	--	---

Section I

1. Company/Product Number <div style="text-align: center; font-size: 1.2em;">707-238</div>	2. EPA Product Manager <div style="text-align: center;">R. Keigwin</div>	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) CONFIRM 2F Agricultural Insecticide	PM# <div style="text-align: center; font-size: 1.2em;">10</div>	
5. Name and Address of Applicant (Include ZIP Code) Rohm and Haas Company 100 Independence Mall West Phila., PA 19106 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

<input type="checkbox"/> Amendment - Explain below <input type="checkbox"/> Resubmission in response to Agency letter dated _____ <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____ <input type="checkbox"/> "Me Too" Application. <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification per PR Notice 95-5 to reduce Restricted Entry Interval

Section III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes," Unit Package wgt. No. per container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes," Package wgt. No. per container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted.					
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other (_____) <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name <div style="text-align: center; font-size: 1.1em;">Ann M. Tillman, Ph.D</div>	Title <div style="text-align: center;">Product Registration Manager</div>	Telephone No. (Include Area Code) <div style="text-align: center; font-size: 1.1em;">215-592-3102</div>

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received <div style="text-align: center;">(Stamped)</div>
2. Signature 	3. Title <div style="text-align: center;">Product Registration Manager</div>	
4. Typed Name <div style="text-align: center;">Ann M. Tillman</div>	5. Date <div style="text-align: center; font-size: 1.1em;">12-4-95</div>	

CONFIRM[®] 2F AGRICULTURAL INSECTICIDE	ROHM & HAAS PHILADELPHIA, PA 19106 
--	--

ACTIVE INGREDIENT

Tebufenozide

Benzoic acid, 3,5-dimethyl-, 1-(1,1-dimethylethyl)-2-(4-ethylbenzoyl)
hydrazide.....

23.0%*

INERT INGREDIENTS.....

77.0%

TOTAL 100.00%

*Equivalent to 2 lbs. active ingredient per gallon

EPA REG NO. 707-238

EPA EST No. 42761-MS-001

NOTICE: Before using this product, read the entire Precautionary Statements, Notice of Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions.

<p>KEEP OUT OF REACH OF CHILDREN</p> <p>CAUTION</p> <p>STATEMENT OF PRACTICAL TREATMENT</p> <p>IF IN EYES: Flush eyes with plenty of water. Consult a physician if irritation persists.</p> <p>IF INHALED: Move victim to fresh air.</p> <p>IF ON SKIN: Wash affected skin thoroughly with soap and water. Consult a physician if irritation persists.</p> <p>IF SWALLOWED: Dilute by giving two glasses of water to drink and consult a physician. Never give anything by mouth to an unconscious person.</p> <p>NOTE TO PHYSICIAN: Emesis is recommended.</p>

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

May cause eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not swallow, get in eyes, on skin or breathe spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment washwaters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where the water table is shallow may result in groundwater contamination. Application should not be made to field areas which are less than 100 feet from permanent natural bodies of surface water or reservoirs.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of **4 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Coveralls
- Water-proof gloves
- Shoes plus socks

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry well-ventilated area, but not below 32°F.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned stay out of smoke.

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED: Appropriate protective equipment must be worn when handling a spill of this material. Transfer spilled material to suitable containers for recovery or disposal. Contain spills immediately with inert materials (e.g. sand, earth). Transfer liquids and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. **Refer to Precautionary Statements.**

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

GENERAL INFORMATION

CONFIRM 2F Agricultural Insecticide mimics the action of the natural insect hormone 20-hydroxyecdysone, the physiological inducer of the molting and metamorphosis process in insects. CONFIRM 2F is highly active against most lepidopterous larvae while having practically no activity at typical use-rates against other orders of insects. The selectivity of CONFIRM 2F allows for the maintenance of the populations of beneficial and predatory insects which is a key element in integrated pest management programs. CONFIRM 2F controls lepidopterous larvae through a novel mode of action by the induction of a premature lethal molt which initiates within hours of ingestion of treated crop surfaces. Contact activity has also been observed in some insects. Actual death of the larvae will take several days to occur although feeding by the insects generally ceases within 24 hours of ingestion.

USE RATE DETERMINATION

Carefully read, understand and follow label use rates, recommendations and restrictions. Apply the amount specified in the following table with properly calibrated aerial or ground spray equipment. The low rates may be used for light infestations of the target lepidopterous species and the higher rates for moderate to heavy infestations. CONFIRM 2F may be applied in either dilute or concentrate sprays so long as the application equipment is calibrated and adjusted to deliver thorough, uniform coverage. Use the specified amount of CONFIRM 2F per acre regardless of spray volume used.

MIXING AND COMPATIBILITY

Slowly pour CONFIRM 2F into the spray tank as it is being filled. Triple-rinse empty container and add rinsate to spray tank.

CONFIRM 2F Agricultural Insecticide is believed to be compatible with most commonly used agricultural fungicides, insecticides, growth regulators, foliar fertilizers and spray adjuvants. If in doubt, mix proportional amounts of all spray ingredients in a test vessel. Shake the mixture vigorously and allow it to stand for fifteen minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied.

APPLICATION TIMING

The activity of CONFIRM 2F Insecticide is expressed primarily through ingestion by the target larvae. Consequently, the timing of application is dependent on the feeding behavior of the target pest. For cryptic feeding larvae, application must be made prior to the time that surface feeding occurs. For foliar or surface feeding larvae, application made while active feeding is occurring will be effective. For best results, begin applications when first signs of feeding damage or when threshold levels of moths, eggs or larvae occur. Consult the Cooperative Extension Service, or other qualified professional authorities to determine the appropriate threshold for application in your area.

APPLICATION INSTRUCTIONS

Because CONFIRM 2F Insecticide must be ingested, application must be in a manner that assures uniform and thorough coverage. Prepare only the amount of spray solution required to treat the measured acreage. Avoid application under conditions when uniform coverage cannot be assured or when excessive spray drift may occur.

CHEMIGATION

Do not apply this product through any type of irrigation system.

SPRAY ADJUVANTS

The addition of agricultural adjuvants to CONFIRM 2F Agricultural Insecticide sprays will improve initial spray deposits, redistribution and weatherability. The following spray adjuvants have been especially formulated to optimize the performance of foliar-applied agricultural chemicals:

LATRON B-1956TM -A water-dispersible, resin-based nonionic surfactant that resists re-wetting and removal by rain. Effective with dilute sprays applied by ground equipment.

LATRON CS-7TM -A spreader-binder designed specifically for use in concentrate and low volume sprays applied by aircraft or ground equipment.

Place CONFIRM 2F Agricultural Insecticide into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation.

6.36

USE DIRECTIONS FOR WALNUTS

Ground Application: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to walnut trees 4th leaf or younger. For walnut trees 5th leaf or older use a minimum of 100 gallons per acre. Ground speed of the sprayer should not exceed 2 mph.

Aerial Application: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 20 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Split Application: In order to achieve thorough, uniform coverage of extremely tall, dense trees it may be preferable to apply a split application composed of both aerial and ground methods. Both portions of the application must be made within the timing window as described below. The total amount of CONFIRM 2F Agricultural Insecticide applied in a split application cannot exceed 16 ounces per acre.

Target Pests	Application Rate (ounces per acre)	Recommendations	Restrictions
Codling moth	16.0 (0.25lb.ai/acre)	For each codling moth generation, apply at initiation of egg hatch (200-250 DD following biofix). Control of first generation codling moth may require a second application 10-14 days following first application to ensure complete coverage of rapidly expanding foliage and expanding surface area of the walnut. After plant foliage expansion and walnut growth has ceased, multiple applications (every 14-21 days) may be required to provide control of extended codling moth flights.	Do not apply more than 16 oz./application or 64 oz. per season. Allow at least 30 days to elapse between final application and harvest. Allow at least six hours between the completion of insecticide application and the onset of precipitation to assure thorough spray drying. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
Navel Orange Worm		Apply at initiation of egg hatch.	
Fall Webworm		Apply at first sign of larvae appearance.	
Redhumped Caterpillar			

15132-B1b
12/4/95
JL/rdm

CONFIRM, LATRON B-1956 AND CS-7 ARE REGISTERED TRADEMARKS OF ROHM AND HAAS COMPANY