

PM03 707-238

6/29/98

7/6

Revised 6/98

15132-M4a
5/26/98

<h1 style="margin: 0;">CONFIRM[®] 2F</h1> <h2 style="margin: 0;">AGRICULTURAL INSECTICIDE</h2>	 <p style="font-size: small; margin: 0;">ROHM AND HAAS PHILADELPHIA, PA 19106</p>
---------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------

ACTIVE INGREDIENT

Tebufenozide
 Benzoic acid, 3,5-dimethyl-, 1-(1,1-dimethylethyl)-2-(4-ethylbenzoyl) hydrazide.....23.0%*

INERT INGREDIENTS.....77.0%
TOTAL 100.00%

*Equivalent to 2 lbs. active ingredient per gallon

EPA REG NO. 707-238
EPA EST No. 42761-MS-001

NOTICE: Before using this product, read the entire Precautionary Statements, Notice of Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions.


KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT
IF IN EYES: Flush eyes with plenty of water. Consult a physician if irritation persists.
IF INHALED: Move victim to fresh air.
IF ON SKIN: Wash affected skin thoroughly with soap and water. Consult a physician if irritation persists.
IF SWALLOWED: Dilute by giving two glasses of water to drink and consult a physician. Never give anything by mouth to an unconscious person.
NOTE TO PHYSICIAN: Emesis is recommended.

ACCEPTED

JUN 29 1998

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 707-238



**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May cause eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not swallow, get in eyes, on skin or breathe spray mist.

- Personal Protective Equipment (PPE)**
Applicators and other handlers must wear:
- Long-sleeved shirt and long pants
 - Waterproof gloves
 - Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

- Users should:**
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment washwaters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where the water table is shallow may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.
For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Water-proof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry well-ventilated area, but not below 32°F.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned stay out of smoke.

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED: Appropriate protective equipment must be worn when handling a spill of this material. Transfer spilled material to suitable containers for recovery or disposal. Contain spills immediately with inert materials (e.g. sand, earth). Transfer liquids and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Refer to Precautionary Statements.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

GENERAL INFORMATION

CONFIRM 2F Agricultural Insecticide mimics the action of the natural insect hormone 20-hydroxyecdysone, the physiological inducer of the molting and metamorphosis process in insects. CONFIRM 2F is highly active against most lepidopterous larvae while having practically no activity at typical use-rates against other orders of insects. The selectivity of CONFIRM 2F allows for the maintenance of the populations of beneficial and predatory insects which is a key element in integrated pest management programs. CONFIRM 2F controls lepidopterous larvae through a novel mode-of-action by the induction of a premature lethal molt which initiates within hours of ingestion of treated crop surfaces. Contact activity has also been observed in some insects. Actual death of the larvae will take several days to occur although feeding by the insects generally ceases within 24 hours of ingestion.

USE RATE DETERMINATION

Carefully read, understand and follow label use rates, recommendations and restrictions. Apply the amount specified in the following table with properly calibrated aerial or ground spray equipment. The low rates may be used for light infestations of the target lepidopterous species and the higher rates for moderate to heavy infestations. CONFIRM 2F may be applied in either dilute or concentrate sprays so long as the application equipment is calibrated and adjusted to deliver thorough, uniform coverage. Use the specified amount of CONFIRM 2F per acre regardless of spray volume used.

MIXING AND COMPATIBILITY

Slowly pour CONFIRM 2F into the spray tank as it is being filled. Triple-rinse empty container and add rinsate to spray tank.

CONFIRM 2F Agricultural Insecticide is believed to be compatible with most commonly used agricultural fungicides, insecticides, growth regulators, foliar fertilizers and spray adjuvants. If in doubt, mix proportional amounts of all spray ingredients in a test vessel. Shake the mixture vigorously and allow it to stand for fifteen minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied.



4/6

APPLICATION TIMING

The activity of CONFIRM 2F Insecticide is expressed primarily through ingestion by the target larvae. Consequently, the timing of application is dependent on the feeding behavior of the target pest. For cryptic feeding larvae, application must be made prior to the time that surface feeding occurs. For foliar or surface feeding larvae, application made while active feeding is occurring will be effective. For best results, begin applications when first signs of feeding damage or when threshold levels of moths, eggs or larvae occur. Consult the Cooperative Extension Service, or other qualified professional authorities to determine the appropriate threshold for application in your area.

APPLICATION INSTRUCTIONS

Because CONFIRM 2F Insecticide must be ingested, application must be in a manner that assures uniform and thorough coverage. Prepare only the amount of spray solution required to treat the measured acreage. Avoid application under conditions when uniform coverage cannot be assured or when excessive spray drift may occur.

CHEMIGATION

Do not apply this product through any type of irrigation system.

SPRAY ADJUVANTS

The addition of agricultural adjuvants to CONFIRM 2F Agricultural Insecticide sprays will improve initial spray deposits, redistribution and weatherability.


The following spray adjuvants have been especially formulated to optimize the performance of foliar-applied agricultural chemicals and are recommended for use with CONFIRM 2F:

LATRON B-1956® -A water-dispersible, resin-based nonionic surfactant that resists re-wetting and removal by rain. Effective with dilute sprays applied by ground equipment.

LATRON CS-7® -A spreader-binder designed specifically for use in concentrate and low volume sprays applied by aircraft or ground equipment.

Place CONFIRM 2F Agricultural Insecticide into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on both product labels prior to spray preparation.

added for use



USE DIRECTIONS FOR WALNUTS

Ground Application: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to walnut trees 4th leaf or younger. For walnut trees 5th leaf or older use a minimum of 100 gallons per acre. Ground speed of the sprayer should not exceed 2 mph.


Aerial Application: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 20 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Split Application: In order to achieve thorough, uniform coverage of extremely tall, dense trees it may be preferable to apply a split application composed of both aerial and ground methods. Both portions of the application must be made within the timing window as described below. The total amount of CONFIRM 2F Agricultural Insecticide applied in a split application cannot exceed 32 ounces per acre.

Target Pests	Application Rate (ounces per acre)	Recommendations	Restrictions
Codling moth	16.0-32.0 (0.25-0.5 lb.ai/acre)	For each codling moth generation, apply at initiation of egg hatch (200-250 DD following biofix). Control of first generation codling moth may require a second application 10-14 days following first application to ensure complete coverage of rapidly expanding foliage and expanding surface area of the walnut. After plant foliage expansion and walnut growth has ceased, multiple applications (every 14-21 days) may be required to provide control of extended codling moth flights. Higher use rates may also be used for extended residual effectiveness, higher pest infestation levels, larger trees or heavy, dense foliage.	Do not apply more than 32 oz/application or 128 oz. per season. Allow at least 30 days to elapse between final application and harvest. Allow at least six hours between the completion of insecticide application and the onset of precipitation to assure thorough spray drying. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
Navel Orange Worm		Apply at initiation of egg hatch.	
Fall Webworm		Apply at first sign of larvae appearance.	
Redhumped Caterpillar			

15132-M2c
6/25/96
JLJrdm

CONFIRM, LATRON B-1956 AND CS-7 ARE REGISTERED TRADEMARKS OF ROHM AND HAAS COMPANY



6/14

USE DIRECTIONS FOR PECANS

Ground Application: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trees 10 feet tall or less. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre. Operating an airblast sprayer at ground speeds in excess of that which will provide the complete exchange of air volume within the tree canopy or making applications in an alternate row middle pattern, may result in less than satisfactory coverage and poor performance, particularly in conditions of high pest infestation levels, extremely large trees and/or dense foliage.

newly added

Aerial Application: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 20 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Target Pest	Application Rate (Ounces per Acre)	Recommendations	Restrictions
Pecan nut casebearer	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Initiate applications at 1-3% infestation of the cluster. Make subsequent applications at 10 to 14 day intervals under conditions of extended egg lay.	Do not apply more than 80 oz. of CONFIRM 2F per season. Allow at least 14 days to elapse between final application and harvest.
Hickory shuckworm		Initiate applications at half-shell hardening. Make subsequent applications at 14 day intervals to shuck split or while nuts are susceptible to hickory shuckworm for heavy infestations.	Allow at least six hours between the completion of insecticide application and the onset of precipitation to assure thorough spray drying.
Fall webworm		Make applications at the first sign of larval infestation.	Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.

15132-M4a
5/26/98
JL/rdm

CONFIRM, LATRON B-1956 AND CS-7 ARE REGISTERED TRADEMARKS OF ROHM AND HAAS COMPANY

