

PLEASE NOTE

This image contains more than one label approved for this product on this date.

15132-M6a
02/01/00

<h1 style="margin: 0;">CONFIRM[®] 2F</h1> <h2 style="margin: 0;">AGRICULTURAL INSECTICIDE</h2>	 <p style="font-size: small;">PHILADELPHIA, PA 19106</p>
---	--

ACTIVE INGREDIENT

Tebufenozide

Benzoic acid, 3,5-dimethyl-, 1-(1,1-dimethylethyl)-2-(4-ethylbenzoyl)

hydrazide.....23.0%*

INERT INGREDIENTS.....77.0%

TOTAL 100.00%


*Equivalent to 2 lbs. active ingredient per gallon

EPA REG. NO. 707-238

EPA EST. NO. 39578-TX-01

USE DIRECTIONS FOR:

- BUSH BERRIES
- CANE BERRIES
- COLE CROPS, LEAFY VEGETABLES AND TURNIPS
- COTTON
- CRANBERRIES
- FORESTS, TREES AND SHRUBS
- FRUITING VEGETABLES
- MINT
- POME FRUITS
- PECANS
- SUGARCANE
- WALNUTS
- ORNAMENTALS



NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

<p style="text-align: center; font-weight: bold; font-size: 1.2em;">KEEP OUT OF REACH OF CHILDREN</p> <p style="text-align: center; font-weight: bold; font-size: 1.5em;">CAUTION</p> <p>STATEMENT OF PRACTICAL TREATMENT</p> <p>IF IN EYES: Flush eyes with plenty of water. Consult a physician if irritation persists.</p> <p>IF INHALED: Move victim to fresh air.</p> <p>IF ON SKIN: Wash affected skin thoroughly with soap and water. Consult a physician if irritation persists.</p> <p>IF SWALLOWED: Dilute by giving two glasses of water to drink and consult a physician. Never give anything by mouth to an unconscious person.</p> <p>NOTE TO PHYSICIAN: Emesis is recommended.</p>

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May cause eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not swallow, get in eyes, on skin or breathe spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations
Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip.

Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Water-proof gloves
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

- Keep unprotected persons out of treated area until sprays have dried.

added

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry well-ventilated area, but not below 32°F.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned stay out of smoke.

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED: Appropriate protective equipment must be worn when handling a spill of this material. Transfer spilled material to suitable containers for recovery or disposal. Contain spills immediately with inert materials (e.g. sand, earth). Transfer liquids and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Refer to Precautionary Statements.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

GENERAL INFORMATION

CONFIRM 2F Agricultural Insecticide mimics the action of the natural insect hormone 20-hydroxyecdysone, the physiological inducer of the molting and metamorphosis process in insects. CONFIRM 2F is highly active against most lepidopterous larvae while having practically no activity at typical use-rates against other orders of insects. The selectivity of CONFIRM 2F allows for the maintenance of the populations of beneficial and predatory insects which is a key element in integrated pest management programs. CONFIRM 2F controls lepidopterous larvae through a novel mode-of-action by the induction of a premature lethal molt which initiates within hours of ingestion of treated crop surfaces. Contact activity has also been observed in some insects. Actual death of the larvae will take several days to occur although feeding by the insects generally ceases within 24 hours of ingestion.

USE RATE DETERMINATION

Carefully read, understand and follow label use rates, recommendations and restrictions. Apply the amount specified in the tables listed on this label with properly calibrated aerial or ground spray equipment.

The low rates may be used for light infestations of the target lepidopterous species and the higher rates for moderate to heavy infestations. CONFIRM 2F may be applied in either dilute or concentrate sprays so long as the application equipment is calibrated and adjusted to deliver thorough, uniform coverage. Use the specified amount of CONFIRM 2F per acre regardless of spray volume used. Prepare only the amount of spray solution required to treat the measured acreage.

MIXING AND COMPATIBILITY

Fill the spray tank one-third to one-half full of clean water and slowly pour CONFIRM 2F Agricultural Insecticide into the spray tank. Maintain agitation in the spray tank during mixing, loading and application. Triple-rinse empty container and add rinsate to spray tank.

CONFIRM 2F Agricultural Insecticide is believed to be compatible with most commonly used agricultural fungicides, insecticides, growth regulators, foliar fertilizers and spray adjuvants. If in doubt, mix proportional amounts of all spray ingredients in a test vessel. Shake the mixture vigorously and allow it to stand for fifteen minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied.

APPLICATION TIMING

The activity of CONFIRM 2F Agricultural Insecticide is expressed primarily through ingestion by the target larvae. Consequently, the timing of application is dependent on the feeding behavior of the target pest. For internal feeding larvae, application must be made prior to the time that surface feeding occurs. For foliar or surface feeding larvae, application made while active feeding is occurring will be effective.

Re-application may be required to protect new flushes of foliage or rapidly expanding fruit. The re-application interval will vary depending on how rapidly the crop is growing and the generation time of the target pest. While CONFIRM 2F Agricultural Insecticide is essentially equally effective against all instars, it is generally good practice to make applications to early instars to avoid the heavy damage that can be inflicted by later instar larvae.

For best results, begin applications when first signs of feeding damage or when threshold levels of moths, eggs or larvae occur. Consult the Cooperative Extension Service, or other qualified professional authorities to determine the appropriate threshold for application in your area.

APPLICATION INSTRUCTIONS

Because CONFIRM 2F Agricultural Insecticide must be ingested by the larvae, application must be in a manner that assures uniform and thorough coverage. Higher water volume and increased spray pressure generally provide better coverage. Operating an air-blast sprayer at ground speeds greater than 2 mph and making applications in an alternate row middle pattern in tree crops and vines may result in less than satisfactory coverage and poor performance, particularly in conditions of high pest infestation levels, extremely large trees and/or dense foliage. Avoid application under conditions when uniform coverage cannot be assured or when excessive spray drift may occur. A minimum of six hours drying time is required between the completion of application and the onset of precipitation to ensure optimum performance.

CHEMIGATION

Do not apply this product through any type of irrigation system except as specified for use on cranberries.

changed

SPRAY ADJUVANTS

The addition of agricultural adjuvants to CONFIRM 2F Agricultural Insecticide sprays will improve initial spray deposits, redistribution and weatherability. The following spray adjuvants have been especially formulated to optimize the performance of foliar-applied agricultural chemicals and are recommended for use with CONFIRM 2F:

- LATRON B-1956® -A water-dispersible, resin-based nonionic spreader-sticker that resists re-wetting and removal by rain. Effective with dilute sprays applied by ground equipment.
- LATRON CS-7® -A spreader-binder designed specifically for use in concentrate and low volume sprays applied by aircraft or ground equipment.

Place CONFIRM 2F Agricultural Insecticide into suspension prior to adding an adjuvant to the spray mixture. Read and carefully observe the precautionary statements and all other information appearing on all product labels prior to spray preparation.

RESISTANCE MANAGEMENT

Any insect population may contain individuals that are naturally resistant to a specific pesticide, therefore, the use of any one insecticide against many consecutive generations of a pest can result in the development of resistance problems. To prevent or delay the development of resistance, Rohm and Haas Company recommends rotation of CONFIRM 2F with insecticides of alternate modes of action and the utilization of Integrated Pest Management practices such as routine monitoring, the use of treatment thresholds to time applications and cultural and biological controls wherever possible. We further recommend that CONFIRM 2F not be used on more than three consecutive generations of a pest. Since the development of resistance cannot be predicted we suggest that you consult local or State Extension personnel or your local Rohm and Haas Company representative for resistance management guidance appropriate to your crop, locality and production practices.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted at intervals defined below following the final application of CONFIRM 2F Agricultural Insecticide at the recommended rates for a registered use.

CROP	RE-CROPPING INTERVAL
Crops for which CONFIRM 2F use is registered	No restrictions
All other crops	30 days

Note: When using CONFIRM 2F with other registered pesticides, always refer to rotational restrictions and precautions on the other product's label and comply with the most restrictive rotational guidelines.



6/21

USE DIRECTIONS FOR BUSH BERRIES

(Blueberries-high bush and low bush, Currant, Elderberry, Gooseberry and Huckleberry)

GROUND APPLICATION: Make applications by conventional boom or air-blast sprayers which are calibrated to deliver a minimum of 30 gallons per acre.

AERIAL APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 10 gallons per acre.

SPRAY ADJUVANT: A Spreader-Sticker such as LATRON B1956® or similar adjuvant should only be used if recommended by a local expert and if previous experience has been satisfactory. Under certain conditions adjuvant usage can result in blossom and fruit damage.

*wording
 changed*

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Cranberry fruitworm (<i>Acrobasis vaccinii</i>) Cherry fruitworm (<i>Grapholita packardii</i>)	16.0 (0.25 lb. AI/Acre)	Apply at initiation of egg laying (approximately 400 Day Degrees (DD) following biofix*-base 50°F). Make a second application at 100% petal fall (usually 7 to 14 days following the first application). Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	Do not apply more than 64 fluid ounces product per season. Allow at least 14 days to elapse between final application and harvest.
Obliquebanded leafroller (<i>Choristoneura rosaceana</i>)		Spring (overwintering) generation: Make one to two applications at bloom to petal fall to small larvae when threshold levels occur. Summer generation: Begin applications at peak moth flight (200-300 DD) following biofix-base 43°F. Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	
Redbanded leafroller (<i>Argyrotaenia velutinana</i>) Variegated leafroller (<i>Platynota flavedana</i>)		For control of other leafrollers, begin applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10 to 14 days to ensure complete coverage of rapidly expanding fruits or foliage.	
Spanworm		Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	
Green fruitworm (<i>Lithophane antennata</i>)		Begin applications when larvae are first detected in the clusters or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	
Armyworm Cutworm	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	
Gypsy moth (<i>Lymantria dispar</i>)	4.0 to 8.0 (0.06 to 0.12 lb. AI/Acre)	Apply to early instars (1 st , 2 nd , or 3 rd) at first signs of infestation.	

Revised

* First sustained moth catch (biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period

USE DIRECTIONS FOR CANE BERRIES*

GROUND APPLICATION: Make applications by conventional boom or air-blast sprayers which are calibrated to deliver a minimum of 30 gallons per acre.

AERIAL APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 10 gallons per acre.

SPRAY ADJUVANT: A Spreader-Sticker such as LATRON B1956® or similar adjuvant should only be used if recommended by a local expert and if previous experience has been satisfactory. Under certain conditions adjuvant usage can result in blossom and fruit damage. *revised*

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Orange tortrix (<i>Argyrotaenia citrana</i>)	16.0 (0.25 lb. AI/Acre)	Spring generation: Apply at first sign of larval infestation or to small larvae when threshold levels occur. Summer generation: Begin applications at first egg hatch. Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	Do not apply more than 64 fluid ounces product per season. Allow at least 14 days to elapse between final application and harvest.
Obliquebanded leafroller (<i>Choristoneura rosaceana</i>)		Spring (overwintering) generation: Apply at first sign of larval infestation or to small larvae when threshold levels occur. Summer generation: Begin applications at first egg hatch (200-300 DD) following biofix** -base 43°. Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	<i>added</i>
Redbanded leafroller (<i>Argyrotaenia velutinana</i>) Variegated leafroller (<i>Platynota flavedana</i>) Omnivorous leafroller (<i>Platynota stultana</i>)		For control of other leafrollers, begin applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10 to 14 days to ensure complete coverage of rapidly expanding fruits or foliage.	
Alfalfa looper (<i>Autographa californica</i>)	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	
Armyworm Cutworm	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	
Gypsy moth (<i>Lymantria dispar</i>)	4.0 to 8.0 (0.06 to 0.12 lb. AI/Acre)	Apply to early instars (1 st , 2 nd , or 3 rd) at first signs of infestation.	

*The cane berries crop group includes: bingleberry, black satin berry, blackberry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, red and black raspberry, rossberry, Shawnee blackberry, Youngberry and varieties and hybrids of these.

** First sustained moth catch (biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period

USE DIRECTIONS FOR COLE CROPS, LEAFY VEGETABLES* AND TURNIPS (TOPS AND ROOTS)

GROUND APPLICATION: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult-to-cover crops to ensure thorough coverage.

AERIAL APPLICATION: For optimum performance a minimum application volume of 10 gallons per acre is recommended. Lower carrier volumes may result in less uniform coverage and reduced efficacy. *revised*

SPRAY ADJUVANT: One pint of LATRON CS-7 per 100 gallons of spray mixture or a similar spreader-binder is recommended to maximize coverage and distribution of the spray material.

TARGET PESTS	APPLICATION RATE (Fluid Ounces per Acre)	APPLICATION TIMING	RESTRICTIONS
Beet armyworm Cabbage looper Cabbage webworm Cross-striped cabbageworm Fall armyworm Garden webworm Imported cabbageworm Southern armyworm True armyworm Yellowstriped armyworm	6.0 to 8.0 (0.09 to 0.12 lb. AI/Acre)	For early-season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities	Do not apply more than 8 fluid ounces per application and do not exceed 56 ounces product per season. Allow at least 7 days to elapse between final application and harvest See Rotational Crop Restrictions in the body of this label.
	8.0 (0.12 lb. AI/Acre)	For mid- to late-season applications and to heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10- to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	

* The Cole crop grouping (Brassica leafy vegetables) includes broccoli, Chinese broccoli, broccoli raab, Brussels sprouts, cabbage, bok choy, Napa cabbage, Chinese mustard cabbage, cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens. The Leafy Vegetables grouping includes amaranth, arugula, cardoon, celery, Chinese celery, celtuce, chervil, edible-leaved chrysanthemum, garland chrysanthemum, corn salad, garden cress, upland cress, dandelion, dock, endive, Florence fennel, lettuce, orach, parsley, garden purslane, winter purslane, radicchio, rhubarb, spinach, New Zealand spinach, vine spinach, Swiss chard.


USE DIRECTIONS FOR COTTON

GROUND APPLICATION; Make applications by conventional ground sprayers which are calibrated to deliver a minimum of 5 gallons per acre.

AERIAL APPLICATION: Make applications of CONFIRM 2F in a minimum of 2 gallons per acre.

SPRAY ADJUVANTS: Use of LATRON CS-7 or a similar spreader-binder following the manufacturer's labeling is recommended to maximize coverage and distribution of spray mixture.

TARGET PESTS	APPLICATION RATE (Fluid Ounces per Acre)	APPLICATION TIMING	RESTRICTIONS
Beet armyworm	4.0 to 8.0 (0.06 to 0.12 lb. AI/Acre)	For early-season (pre-squaring) applications only. Apply when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities	Do not apply more than 16 fluid ounces per application and do not exceed 64 fluid ounces per season. Allow at least 14 days to elapse between final application and harvest. See Rotational Crop Restrictions in the body of this label
	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	For mid- to late-season applications and to infestations of 2 to 10 hits per 100 row ft. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10- to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	
Cabbage looper Fall armyworm Southern armyworm True armyworm Yellowstriped armyworm	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Apply when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities	



10/21

USE DIRECTIONS FOR CRANBERRIES

GROUND APPLICATION: Make applications by conventional ground sprayers which are calibrated to deliver a minimum of 20 gallons per acre.

AERIAL APPLICATION: Make applications of CONFIRM 2F in a minimum of 10 gallons per acre.

CHEMIGATION APPLICATION: For use only in solid-set sprinkler systems designed specifically for chemigation. Apply through a properly calibrated chemigation system that has the appropriate back flow prevention devices. See the 'MIXING' section of this labeling for specific mixing and dilution instructions. CONFIRM 2F should be applied in dedicated chemigation cycles only, not as a part of a regular irrigation cycle. Do not exceed 900 gallons of water per acre application volume. Minimum volume should be used for flushout to avoid diluting or rinsing off product. Washout time should not exceed six (6) minutes. Sprinkler heads should be set in a spacing not exceeding 50 feet by 60 feet and adjusted to provide 100% overlap. Crop injury, lack of effectiveness, or illegal pesticide residues can result from non-uniform distribution of treated water.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH CHEMIGATION SYSTEMS

- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a positive displacement, metering injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

-CRANBERRIES cont.-

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Blackheaded fireworm	16.0 (0.25 lb. AI/Acre)	First generation: Apply at first sign of larval infestation and make a second application 7 to 10 days following the first application. Second generation: Make the first application at 10 to 20% egg hatch (typically 8 to 12 days following biofix*) followed by a second application 7 to 10 days later.	Do not apply more than 64 fluid ounces of CONFIRM 2F per season Allow at least 30 days to elapse between final application and harvest.
Spotted fireworm	16.0 (0.25 lb. AI/Acre)	First generation: Apply to small larvae before webbing occurs when threshold infestations are detected by sweep net sampling. Make a second application 7 to 10 days following the first application to heavy infestations or sustained moth flight. Second generation: Make the first application at 10 to 20% egg hatch (usually mid to late June) followed by a second application 7 to 10 days later.	
Sparganothis fruitworm	16.0 (0.25 lb. AI/Acre)	First generation: Initiate applications as soon as larvae are detected by sweep net sampling. Make a second application in 7 to 14 days. Summer generation: Make the first application at 5 to 10% egg hatch (usually 10 to 14 days following biofix) followed by a second application 7 to 10 days later (about 60% egg hatch).	
Cranberry fruitworm	16.0 (0.25 lb. AI/Acre)	Apply at initiation of egg laying (approximately 400 Day Degrees (DD) following biofix-base 50°F). Make a second application at 100% petal fall (usually 7 to 14 days after the initial application). Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight. Chemigation application for control of cranberry fruitworm is not recommended.	
Blossom worm False armyworm Gypsy moth	16.0 (0.25 lb. AI/Acre)	Apply when larvae are in the 1 st to 3 rd instar and when action thresholds are reached based on local Extension Service recommendations.	
Spanworms	16.0 (0.25 lb. AI/Acre)	Initiate applications when infestations reach threshold levels based on sweep net sampling. Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	

added

*Biofix is defined as first sustained adult catch in pheromone traps, typically, 5 moths in 3 traps within a seven-day period.

USE DIRECTIONS FOR FORESTS*, TREES** AND SHRUBS***

Because CONFIRM 2F Agricultural Insecticide must be ingested in order to be effective, it is essential that coverage is thorough and uniform. Higher carrier volumes and higher use rates are recommended for very large trees or dense stands and for heavy target pest infestations.

GROUND APPLICATION: Hydraulic ground sprayers should be calibrated to deliver a minimum of 50 gallons per acre. For mist blowers or air blast sprayers, use a minimum of 10 gallons per acre.

AERIAL APPLICATION: Make applications in a minimum of one gallon per acre. Higher carrier volumes are recommended when environmental conditions are less than ideal for aerial applications.

TARGET PESTS	APPLICATION RATE (Fluid Ounces per Acre)	APPLICATION TIMING	RESTRICTIONS
Browntail moth (<i>Euproctis chrysorrhoea</i>) Elm spanworm (<i>Ennomos subsignaria</i>) Fall cankerworm (<i>Alsophila pometaria</i>) Fall webworm (<i>Hyphantria cunea</i>) Forest tent caterpillar (<i>Malecosoma disstria</i>) Gypsy moth (<i>Lymantria dispar</i>)	4.0 to 8.0 (0.06 to 0.12 lb. AI/Acre)	Apply to early instar (1 st , 2 nd , or 3 rd) larvae, in general, foliage development should be a minimum of 20%.	Do not apply more than 16 fl. ounces per year. Uniform coverage of the foliage is essential to provide maximum protection from defoliation and reduction of egg mass deposition.
Pine tip moth (<i>Rhyacionia frustrana</i> , <i>R. neomexicana</i> , <i>R. buoliana</i> , <i>R. rigidana</i> , <i>R. subtropica</i>)	8.0 (0.12 lb. AI/Acre)	Apply to early instar (1 st to 2 nd) larvae after each new foliage flush, in general, at approximately 25% shoot expansion.	
Spruce budworm (<i>Choristoneura fumiferana</i> and <i>C. occidentalis</i>) Tussock moth (<i>Dasychira pinicola</i> , <i>Lophocampa maculata</i> , <i>Orgyia pseudotsugata</i> , <i>O. vetusta</i>)	4.0 to 8.0 (0.06 to 0.12 lb. AI/Acre)	Make application to 4 th to 5 th instar larvae that are actively feeding of foliage or outside candle caps	

Revised

- * Forests include commercial, private and public forest land, conifer release sites, shelterbelts and windbreaks, and forest plantings.
- ** Trees include Christmas Trees, nurseries and plantations, conifer seed orchards, nonresidential ornamental and shade trees.
- *** Shrubs include nonresidential woody shrubs and vines.

USE DIRECTIONS FOR FRUITING VEGETABLES*

GROUND APPLICATION: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult-to-cover crops to ensure thorough coverage.

AERIAL APPLICATION: For optimum performance a minimum application volume of 10 gallons per acre is recommended. Lower carrier volumes may result in less uniform coverage and reduced efficacy.

SPRAY ADJUVANT: One pint of LATRON CS-7 per 100 gallons of spray mixture or a similar spreader-binder is recommended to maximize coverage and distribution of the spray material.

revised

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Alfalfa looper Beet armyworm Black cutworm Cabbage looper European corn borer Fall armyworm Imported cabbageworm Southern armyworm Tobacco hornworm Tomato hornworm True armyworm Yellowstriped armyworm	6.0 to 8.0 (0.09 to 0.12 lb. AI/Acre)	For early-season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities	Do not apply more than 16 fluid ounces per application and do not exceed 64 ounces product per season. Allow at least 7 days to elapse between final application and harvest.
	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	For mid- to late-season applications and to heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10- to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	See Rotational Crop Restrictions in the body of this label.

*The Fruiting vegetable grouping includes eggplant, ground cherry, pepino, pepper (bell, chili, cooking), pimento, tomatillo and tomato.

USE DIRECTIONS FOR MINT

GROUND APPLICATION: Apply a minimum of 8 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 10 gallons per acre to densely foliated or difficult-to-cover crops to ensure thorough coverage.

AERIAL APPLICATION: Make applications of CONFIRM 2F in a minimum of 5 gallons per acre.

SPRAY ADJUVANT: One pint of LATRON CS-7 per 100 gallons of spray mixture or a similar spreader-binder is recommended to maximize coverage and distribution of the spray material.

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Alfalfa looper <i>(Autographa californica)</i> Beet armyworm <i>(Spodoptera exigua)</i> Bertha armyworm <i>(Mamestra configurata)</i>	6.0 to 8.0 (0.09 to 0.12 lb. AI/Acre)	For early-season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities	Do not apply more than 16 fluid ounces per application and do not exceed 64 ounces product per season. Allow at least 14 days to elapse between final application and harvest.
	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre))	For mid- to late-season applications and to heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10- to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	

USE DIRECTIONS FOR POME FRUITS

(Apples, Crabapples, Loquat, Mayhaw, Pears including Oriental, Quince)

GROUND APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trellised trees or trees 10 feet tall or less. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre

AERIAL APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 20 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

SPRAY ADJUVANT: The use of a Spreader-Sticker such as LATRON B-1956 or similar Spreader-Sticker to maximize uniform coverage and distribution of the spray material is recommended.

Revised

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Codling moth (East of the Rockies)	20.0 (0.31 lb. AI/Acre)	For each codling moth generation, apply at initiation of egg hatch [150 to 250 Day Degrees (DD), base 50°F, following biofix*] followed by a second application at 10 to 15 days following the first application (usually 450 to 550 DD). Additional applications at 10 to 15-day intervals may be required under high infestations, sustained moth flight, or to ensure coverage of rapidly expanding fruits or foliage.	Do not apply more than 20 oz. per application or 120 oz. CONFIRM 2F per acre per season. Allow at least 14 days to elapse between final application and harvest.
Codling moth (West of the Rockies) For use against low to moderate infestations in conjunction with alternate control measures such as in established Mating Disruption blocks.	20.0 (0.31 lb. AI/Acre)	For each codling moth generation, apply at initiation of egg hatch [150 to 250 Day Degrees (DD), base 50F, following biofix*] followed by a second application at 10 to 15 days following the first application (usually 450-550 DD). Additional applications at 10- to 15-day intervals may be required under high infestations, sustained moth flight, or to ensure coverage of rapidly expanding fruits or foliage.	Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
Obliquebanded leafroller (West of the Rockies)	20.0 (0.31 lb. AI/Acre)	Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending on infestation level. Summer generation: Begin applications at early egg lay through early egg hatch (usually 200 to 400 DD, base 43°F, following biofix). Make a second application at 10 to 18 days later (usually 650 to 850 DD). A third application 10 to 14 days after the second application may be required under high pressure, sustained moth flight or prolonged shoot growth.	
Obliquebanded leafroller	20.0 (0.31 lb. AI/Acre)	Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending on infestation level. Summer generation: Begin applications at peak moth flight (200-300 DD, base 43°F, following biofix*). Make a second application 7 to 14 days later (usually 500-600 DD). A third application 10 to 12 days after the second application (usually 800 to 900 DD), may be required under high pressure, sustained moth flight or prolonged shoot growth. Fall overwintering generation: Apply to late season larval infestations of overwintering generation to minimize damage to the fruit.	

Revised

Pandemis leafroller	20.0 (0.31 lb. AI/Acre)	<p>Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending on infestation level.</p> <p>Summer generation: Begin applications at early egg lay through early egg hatch (250 to 400 DD, base 41°F, following biofix). Make a second application 10 to 18 days (usually 600-800 DD). Under heavy infestation a third application may be required 10 to 14 days after the second application.</p>
Tufted apple bud moth	12.0 - 20.0 (0.19 - 0.31 lb. AI/Acre)	<p>First generation: Make application at 10-30% egg hatch (600 to 900 DD, base 45°F, after biofix depending on local conditions). A second application at 60 to 90% egg hatch may be required under heavy infestation levels.</p> <p>Second generation: Make the first application at 20 to 30% egg hatch (2300-2500 DD). A second application approximately 14 days later may be required under high pressures or sustained moth flight or late maturing varieties.</p>
Eyespotted bud moth Fruittree leafroller Redbanded leafroller Variegated leafroller	20.0 (0.31 lb. AI/Acre)	<p>For control of other leafrollers, begin applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10 to 14 days to ensure complete coverage of rapidly expanding fruits or foliage.</p>
Lesser appleworm	20.0 (0.31 lb. AI/Acre)	<p>For each generation, apply at initiation of egg hatch before larvae enter the fruit. Make a second application 10 to 14 days following the first to ensure complete coverage of rapidly expanding fruits or foliage or under conditions of high infestation or sustained moth flight.</p>
Green fruitworm	20.0 (0.31 lb. AI/Acre)	<p>Apply at initiation of egg hatch or at the first sign of larval infestation. A second application may be required 10 to 14 days following the first application to ensure complete coverage of rapidly expanding fruits or foliage.</p>

Reviewed

*Biofix is defined as first sustained adult catch in pheromone traps, typically, 5 moths in 3 traps within a seven-day period. Consult State Extension Specialist or other qualified authorities for specific information regarding number, placement and management of pheromone traps.

added

USE DIRECTIONS FOR PECANS

GROUND APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trees 10 feet tall or less. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre.

AERIAL APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 5 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, application by this method may result in reduced efficacy if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

revised

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Pecan nut casebearer	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	For each generation, apply at the initiation of egg hatch (for first generation this is approximately 8 to 15 days following first sustained moth catch*). Control of first generation pecan nut casebearer may require a second application under conditions of extended egg lay or for improved coverage of rapidly expanding nuts and foliage. Use higher rates for extended residual effectiveness, higher pest infestations, low crop load, larger trees or heavy, dense foliage.	Do not apply more than 80 fluid ounces of CONFIRM 2F per season. Allow at least 14 days to elapse between final application and harvest. Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
Hickory shuckworm	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Initiate applications at half-shell hardening. Make subsequent applications at 14-day intervals to shuck split or while nuts are susceptible to hickory shuckworm for heavy infestations.	
Fall webworm Walnut caterpillar	8.0 to 16.0 (0.12 to 0.25 lb. AI/Acre)	Make applications at the first sign of larval infestation.	

* First sustained moth catch (biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period.



USE DIRECTIONS FOR SUGARCANE

GROUND APPLICATION: Make applications by conventional ground sprayers which are calibrated to deliver a minimum of 10 gallons per acre.

AERIAL APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 2 gallons per acre. Higher carrier volume may be required to provide thorough coverage under conditions of high temperatures, low humidity or dense crop canopy.

SPRAY ADJUVANT: Use of LATRON CS-7 or similar spreader-binder following the manufacturer's labeling is recommended to maximize coverage and distribution of spray mixture.

TARGET PEST	APPLICATION RATE (Fluid Ounces per Acre)	APPLICATION TIMING	RESTRICTIONS
Sugarcane borer (<i>Diatrea saccharalis</i>)	6.0 to 8.0 (0.09 to 0.12 lb. AI/Acre)	Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	Do not apply more than 16 fluid ounces per application and do not exceed 64 fluid ounces per season. Allow at least 14 days to elapse between final application and harvest.
Mexican rice borer (<i>Eoreuma lofifini</i>)	16.0 (0.25 lb. AI/Acre)	Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10- to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	See Rotational Crop Restrictions in the body of this label.

11/21

USE DIRECTIONS FOR WALNUTS

GROUND APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to walnut trees 4th leaf or younger. For walnut trees 5th leaf or older use a minimum of 100 gallons per acre. Ground speed of the sprayer should not exceed 2 mph.

AERIAL APPLICATION: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 20 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

SPLIT APPLICATION: In order to achieve thorough uniform coverage of extremely tall, dense trees, it may be preferable to apply a split application composed of both aerial and ground methods. Both portions of the application must be made within the timing window as described below. The total amount of CONFIRM 2F Agricultural Insecticide applied in a split application cannot exceed 32 fluid ounces per acre.

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Codling moth	16.0 to 32.0 (0.25 to 0.5 lb.AI/Acre)	For each codling moth generation, apply at initiation of egg hatch (200 to 250 DD following biofix*). Control of first generation codling moth may require a second application 10 to 14 days following first application to ensure complete coverage of rapidly expanding foliage and expanding surface area of the walnut. After plant foliage expansion and walnut growth has ceased, multiple applications (every 14 to 21 days) may be required to provide control of extended codling moth flights. Higher use rates may also be used for extended residual effectiveness, higher pest infestation levels, larger trees or heavy, dense foliage.	<p>Do not apply more than 32 fluid ounces per application or 128 oz. per season.</p> <p>Allow at least 30 days to elapse between final application and harvest.</p> <p>Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.</p>
Navel orange worm	16.0 to 32.0 (0.25 to 0.5 lb.AI/Acre)	Apply at initiation of egg hatch.	
Fall webworm	16.0 to 32.0 (0.25 to 0.5 lb.AI/Acre)	Apply at first sign of larvae appearance.	

* First sustained moth catch (biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period.

20/21

USE DIRECTIONS FOR ORNAMENTALS

When used as recommended, CONFIRM 2F Agricultural Insecticide will control the designated pests on trees, shrubs, foliage plants and flowers grown in commercial nurseries and greenhouses, in Christmas tree farms, in outdoor landscape areas such as parks, recreational areas, institutional grounds, residential property, etc. and in interior landscapes.

When applied as directed, CONFIRM 2F has shown excellent tolerance on a wide range of ornamental plants. It is impossible, however, to evaluate this product on all ornamentals or under all possible growing conditions. The user should exercise reasonable judgment and caution with this product. Until familiar with results under user growing conditions, a limited number of plants should be treated.

Note: Resistance to pesticides has been shown to develop when a pesticide is used continuously against many generations of a target pest. Rohm and Haas Company encourages the periodic interruption of continuous use by utilization of Integrated Pest Management (IPM) practices or by the periodic use of a product with an alternative mode of action to delay or prevent development of resistance. Since the development of resistance cannot be predicted, we suggest you consult local or State Extension Service personnel for resistance management strategies appropriate to your crop locality and production practices.

Hand Sprayers: Make applications using enough water to thoroughly spray plant foliage until runoff. Refer to the following table for product recommendations when using a hand sprayer.

Label Recommendations CONFIRM 2F Insecticide Fl. Oz. Per Acre	Active Ingredient (LB. AI/Acre)	Equivalent CONFIRM 2F in 1 Gallon of Water (Teaspoon)
4	0.06	1/4
8	0.12	1/2
16	0.25	1

Ground Application: Make applications of Confirm 2F Insecticide by conventional ground or hydraulic sprayers which are calibrated to deliver a minimum of 50 gallons per acre. For mist blowers or air blast sprayers, use a minimum of 10 gallons per acre. Application equipment should be properly calibrated and provide uniform spray coverage throughout the plant canopy.

Aerial Application: Make applications of Confirm 2F Insecticide in a minimum of 20 gallons per acre. Confirm 2F can be applied by aerial application when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits through, uniform coverage of the entire tree canopy.

Spray Adjuvants: A spray adjuvant should be used with Confirm 2F Insecticide applications. The adjuvant will improve initial spray deposits, redistribution and weatherability of Confirm 2F. The adjuvant chosen should be approved for use on the ornamentals being treated. Since some adjuvants can be phytotoxic to certain ornamental plants, the user should have prior experience with the adjuvant before combining it with Confirm 2F.

*Added to
master
label*

TARGET PESTS	APPLICATION RATE Fluid Ounces per Acre	APPLICATION TIMING	RESTRICTIONS
Armyworm <i>Pseudaletia unipuncta</i>	4 to 8 (0.06 to 0.12 lb. AI/Acre)	For best results, begin applications when larvae are observed or at the first sign of feeding damage. Repeat applications on a 10-14 day interval or as necessary based on pest reinfestation.	Allow at least six hours between the completion of insecticide applications and the onset of precipitation to assure thorough spray drying. Uniform coverage of the foliage is essential to provide maximum protection from defoliation and reduction of egg mass deposition.
Beet armyworm <i>Spodoptera exigua</i>			
Elm spanworm <i>Ennomos subsignaria</i>			
Eucalyptus caterpillar <i>Thyrinzeina arnobia</i>			
Fall armyworm <i>Spodoptera frugiperda</i>			
Fall cankerworm <i>Alsophila pometaria</i>			
Fall webworm <i>Hyphantria cunea</i>			
Forest Tent caterpillar <i>Malecosoma disstria</i>			
Gypsy moth <i>Lymantria dispar</i>			
Hemlock looper <i>Lambdina fiscellaria</i>			
Jackpine budworm <i>Choristoneura pinus</i>			
Pine tip moth <i>Rhyacionia frustrana</i> , <i>R. neomexicana</i> <i>R. buoliana</i> , <i>R. rigidana</i> , <i>R. subtropica</i>			
Spruce budworm <i>Choristoneura fumiferana</i> Western Spruce Budworm <i>C. occidentalis</i>			
Tussock moth <i>Dasychira pinicola</i> <i>Lophocampa maculata</i> <i>Orgyia pseudotsugata</i> <i>O. vetusta</i>			
Bagworms <i>Thridopteryx ephemeraeformis</i>	8 to 16 (0.12 to 0.25 lb. AI/Acre)		
Codling moth <i>Cydia pomonella</i>			
Cutworms			
Eastern tent caterpillar <i>Malacosoma americanum</i>			
Processionary caterpillar <i>Thaumtopoea pityocampa</i>			
Yellowneck caterpillar <i>Datana ministra</i>			

15132-M6a
02/01/00

CONFIRM, LATRON B-1956 AND CS-7 ARE REGISTERED TRADEMARKS OF ROHM AND HAAS COMPANY

NEXT

LABEL

CONFIRM 2F**Supplemental
Labeling****ACCEPTED**

MAY 31 2000

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 707-238**SUPPLEMENTAL LABELING FOR USE OF
CONFIRM® 2F AGRICULTURAL INSECTICIDE
IN BUSH BERRIES****EPA Reg. No. 707-238****EPA Est. No. 39578-TX-01****DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10-feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

USE DIRECTIONS FOR BUSH BERRIES (Blueberries - high bush and low bush, Currant, Elderberry, Gooseberry, Huckleberry)

Ground Application: Make applications by conventional boom or airblast sprayers which are calibrated to deliver a minimum of 30 gallons per acre.

Aerial Application: Make applications of CONFIRM 2F in a minimum of 10 gpa.

Adjuvants: A Spreader-Sticker such as LATRON B-1956™ or similar adjuvant should only be used if recommended by a local expert and if previous experience has been satisfactory. Under certain conditions adjuvant usage can result in blossom and fruit damage.

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Obliquebanded leafroller (<i>Choristoneura rosaceana</i>)	16.0 (0.25 lb. AI/acre)	Spring (overwintering) generation: Make one to two applications at bloom to petal fall to small larvae when threshold levels occur. Summer generation: Begin applications at peak moth flight (usually 200-300 DD following biofix-base 43°F). Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	Do not apply more than 64 ounces product per season Allow at least 14 days to elapse between final application and harvest
Redbanded leafroller (<i>Argyrotaenia velutinana</i>) Variegated leafroller (<i>Platynota flavedana</i>)		For control of other leafrollers, began applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10 to 14 days to ensure complete coverage of rapidly expanding fruits or foliage.	
Gypsy moth (<i>Lymantria dispar</i>)	4.0 to 8.0 (0.06 to 0.12 lb. AI/acre)	Apply to early instars (1 st , 2 nd , or 3 rd) at first signs of infestation.	
Spanworm	16.0 (0.25 lb. AI/acre)	Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities	

*First sustained moth catch (Biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period.


CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**



NEXT

LABEL

CONFIRM 2F**Supplemental
Labeling****SUPPLEMENTAL LABELING FOR USE OF
CONFIRM® 2F AGRICULTURAL INSECTICIDE
IN CANEBERRIES****EPA Reg. No. 707-238****EPA Est. No. 39578-TX-01****DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

USE DIRECTIONS FOR CANEBERRIES*

Ground Application: Make applications by conventional boom or airblast sprayers which are calibrated to deliver a minimum of 30 gallons per acre.

Aerial Application: Make applications of CONFIRM 2F in a minimum of 10 gpa.

Adjuvants: A Spreader-Sticker such as LATRON B-1956™ or similar adjuvant should only be used if recommended by a local expert and if previous experience has been satisfactory.

Under certain conditions adjuvant usage can result in blossom and fruit damage.

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Obliquebanded leafroller (<i>Choristoneura rosaceana</i>)	16.0 (0.25 lb. ai/acre)	Spring (overwintering) generation: Apply at first sign of larval infestation or to small larvae when threshold levels occur. Summer generation: Begin applications at early egg lay through early egg hatch (usually 200-400 DD following biofix** -base 43°). Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	Do not apply more than 64 ounces product per season Allow at least 14 days to elapse between final application and harvest
Redbanded leafroller (<i>Argyrotaenia velutinana</i>) Variegated leafroller (<i>Platynota flavedana</i>)		For control of other leafrollers, began applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10 to 14 days to ensure complete coverage of rapidly expanding fruits or foliage.	
Gypsy moth (<i>Lymantria dispar</i>)	4.0 to 8.0 (0.06-0.12 lbs. ai/acre)	Apply to early instars (1 st , 2 nd , or 3 rd) at first signs of infestation.	

*The caneberreries crop group includes bingleberry, black satin berry, blackberry, boysenberry, Cherokee blackberry, Chesterberry, Cheyenne blackberry, coryberry, darrowberry, dewberry, Dirksen thornless berry, Himalayaberry, hullberry, Lavacaberry, loganberry, lowberry, Lucretiaberry, mammoth blackberry, marionberry, nectarberry, olallieberry, Oregon evergreen berry, phenomenalberry, rangeberry, ravenberry, red and black raspberry, rossberry, Shawnee blackberry, youngberry, and varieties and hybrids of these.

**First sustained moth catch (Biofix) is defined as the date on which the total of five moths are captured in three pheromone traps within a seven-day period.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**




NEXT

LABEL

CONFIRM 2F Supplemental Labeling

SUPPLEMENTAL LABELING FOR USE OF CONFIRM® 2F AGRICULTURAL INSECTICIDE IN COLE CROPS, LEAFY VEGETABLES AND TURNIPS (TOPS AND ROOTS)

EPA Reg. No. 707-238
EPA Est. No. 39578-TX-01



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale, and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product, unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted at intervals defined below following the final application of CONFIRM 2F at the recommended rates for a registered use.

Crop	Recropping Interval
Crops for which CONFIRM use is registered.	No restrictions
All other crops	30 days

Note: When using CONFIRM 2F with other registered pesticides, always refer to rotational restrictions and precautions on the other product's label and comply with the most restrictive rotational guidelines.

USE DIRECTIONS FOR COLE CROPS AND LEAFY VEGETABLES* AND TURNIPS (TOPS AND ROOTS)

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult-to-cover crops to insure thorough coverage.

Aerial Application: For optimum performance a minimum application volume of 10 gallons per acre is recommended. Lower carrier volumes may result in less uniform coverage and reduced efficacy.

Spray Adjuvant: One pint of LATRON CS-7 per 100 gallons of spray mixture or similar spreader-binder is recommended to maximize coverage and distribution of the spray material.

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Beet armyworm Cabbage looper Cabbage webworm Cross-striped cabbageworm Fall armyworm	6.0 to 8.0 (0.09 to 0.12 lb. ai/acre)	For early season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	Do not apply more than 8 fluid ounces per application and do not exceed 56 fluid ounces per season.
Garden webworm Imported cabbageworm Southern armyworm True armyworm Yellowstriped armyworm	8.0 (0.12 lb. ai/acre)	For mid- to late-season applications and to heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10 to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	Allow at least 7 days to elapse between final application and harvest. See Rotational Crop Restrictions in the body of this label.

*The Cole crop grouping (Brassica leafy vegetables) includes broccoli, Chinese broccoli, broccoli raab, Brussels sprouts, cabbage, bok choy, Napa cabbage, Chinese mustard cabbage, cauliflower, cavalo broccolo, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens. The Leafy Vegetables grouping includes amaranth, arugula, cardoon, celery, Chinese celery, celtuce, chervil, edible-leaved chrysanthemum, garland chrysanthemum, corn salad, garden cress, upland cress, dandelion, dock, endive, Florence fennel, lettuce, orach, parsley, garden purslane, winter purslane, radicchio, rhubarb, Spinach, New Zealand spinach, vine spinach, Swiss chard.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**




NEXT

LABEL

CONFIRM 2F Supplemental Labeling

SUPPLEMENTAL LABELING FOR USE OF CONFIRM® 2F AGRICULTURAL INSECTICIDE IN CRANBERRIES



EPA Reg. No. 707-238
EPA Est. No. 39578-TX-01

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

USE DIRECTIONS FOR CRANBERRY

Ground Application: Make applications by conventional ground sprayers which are calibrated to deliver a minimum of 20 gallons per acre.

Aerial Application: Make applications of CONFIRM 2F in a minimum of 10 gpa.

Chemigation Application: For use only in solid-set sprinkler systems designed specifically for chemigation. Apply through a properly calibrated chemigation system that has the appropriate back flow prevention devices. See the 'MIXING' section of this labeling for specific mixing and dilution instructions. CONFIRM 2F should be applied in dedicated chemigation cycles only, not as a part of a regular irrigation cycle. Do not exceed 900 gallons of water per acre application volume. Minimum volume should be used for flush out to avoid diluting or rinsing off product. Washout time should not exceed six (6) minutes. Sprinkler heads should be set in a spacing not exceeding 50 feet by 60 feet and adjusted to provide 100% overlap. Crop injury, lack of effectiveness, or illegal pesticide residues can result from nonuniform distribution of treated water.

General Precautions for Applications Through Chemigation Systems

- Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- Chemigation systems connected to public water systems must contain a functional reduced pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a positive displacement, metering injection pump (e.g. diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Do not apply when wind speed favors drift beyond the area intended for treatment.
- If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

TARGET PESTS	APPLICATION RATE Fluid Ounces per acre	APPLICATION TIMING	RESTRICTIONS
Blackheaded fireworm	16.0 (0.25 lb. ai/acre)	First generation: Apply at first sign of larval infestation and make a second application 7 to 10 days following the first application. Second generation: Make the first application at 10-20% egg hatch (typically 8 to 12 days following biofix*) followed by a second application 7 to 10 days later.	Do not apply more than 64 oz of CONFIRM 2F per season Allow at least 30 days to elapse between final application and harvest.
Spotted fireworm	16.0 (0.25 lb. ai/acre)	First generation: Apply to small larvae before webbing occurs when threshold infestations are detected by sweep net sampling. Make a second application 7 to 10 days following the first application to heavy infestations or sustained moth flight. Second generation: Make the first application at 10-20% egg hatch (usually mid to late June) followed by a second application 7 to 10 days later.	
Sparganothis fruitworm	16.0 (0.25 lb. ai/acre)	First generation: Initiate applications as soon as larvae are detected by sweep net sampling. Make a second application in 7 to 14 days. Summer generation: Make the first application at 5-10% egg hatch (usually 10 to 14 days following biofix) followed by a second application 7 to 10 days later (about 60% egg hatch).	
Blossom worm False armyworm Gypsy moth	16.0 (0.25 lb. ai/acre)	Apply when larvae are in the 1 st to 3 rd instar and when action thresholds are reached based on local Extension Service recommendations.	
Spanworms	16.0 (0.25 lb. ai/acre)	Initiate applications when infestations reach threshold levels based on sweep net sampling. Additional applications at 10- to 14-day intervals may be required under high pressure or sustained moth flight.	

*Biofix is defined as first sustained adult catch in pheromone traps, typically, 5 moths in 3 traps within a seven-day period.


CONDITIONS OF SALE AND WARRANTY
 Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

© 2000 Rohm and Haas Company
 01/24/00



NEXT

LABEL

CONFIRM 2F**Supplemental
Labeling**

**SUPPLEMENTAL LABELING FOR USE OF
CONFIRM® 2F AGRICULTURAL INSECTICIDE
IN FORESTRY**

**EPA Reg. No 707-238
EPA Est. No. 39578-TX-01**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

USE DIRECTIONS FOR FORESTS*, TREES AND SHRUBS*****

Because CONFIRM 2F Agricultural Insecticide must be ingested in order to be effective, it is essential that coverage is thorough and uniform. Higher carrier volumes and higher use rates are recommended for very large trees or dense stands and for heavy target pest infestations.

Ground Application: Hydraulic ground sprayers should be calibrated to deliver a minimum of 50 gallons per acre. For mist blowers or air blast sprayers, use a minimum of 10 gallons per acre.

Aerial Application: Make applications of CONFIRM 2F in a minimum of 1 gallon per acre. Higher carrier volumes are recommended when environmental conditions are less than ideal for aerial applications.

Target pests	Application Rate (fluid ounces per acre)	Application Timing	Restrictions
Browntail moth (<i>Euproctis chrysorrhoea</i>) Elm spanworm (<i>Ennomos subsignaria</i>) Fall cankerworm (<i>Alsophila pometaria</i>) Fall webworm (<i>Hyphantria cunea</i>) Forest tent caterpillar (<i>Malacosoma disstria</i>) Gypsy moth (<i>Lymantria dispar</i>)	4.0 to 8.0 (0.06 to 0.12 lb. ai/acre)	Apply to early instar (1st, 2nd, or 3rd) larvae. In general, foliage development should be a minimum of 20%.	Do not apply more than 16 fl. Ounces per year. Uniform coverage of the foliage is essential to provide maximum protection from defoliation and reduction of egg mass deposition.
Pine tip moth (<i>Rhyacionia frustrana</i> , <i>R. neomexicana</i> , <i>R. buoliana</i> , <i>R. rigidana</i> , <i>R. subtropica</i>)	8.0 (0.12 lb. ai/acre)	Apply to early instar (1st to 2nd) larvae after each new foliage flush, in general, at approximately 25% shoot expansion.	
Spruce budworm (<i>Choristoneura fumiferana</i> and <i>C. occidentalis</i>) Tussock moth (<i>Dasychira pinicola</i> , <i>Lophocampa maculata</i> , <i>Orgyia pseudotsugata</i> , <i>O. vetusta</i>)	4.0 to 8.0 (0.06 to 0.12 lb. ai/acre)	Make application to 4th to 5th instar larvae that are actively feeding on foliage or are feeding outside the candle caps.	

*Forests include commercial, private and public forest land, conifer release sites, shelterbelts and windbreaks, and forest plantings.

**Trees include Christmas trees, nurseries and plantations, conifer seed orchards, nonresidential ornamental and shade trees.

***Shrubs include nonresidential woody shrubs and vines.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**




NEXT

LABEL

CONFIRM 2F Supplemental Labeling

SUPPLEMENTAL LABELING FOR USE OF CONFIRM® 2F AGRICULTURAL INSECTICIDE IN FRUITING VEGETABLES

EPA Reg. No 707-238
EPA Est. No. 39578-TX-01



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale, and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product, unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment washwaters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

ROTATIONAL CROP RESTRICTIONS

The following rotational crops may be planted at intervals defined below following the final application of CONFIRM 2F at the recommended rates for a registered use.

Crop	Recropping Interval
Crops for which CONFIRM use is registered.	No restrictions
All other crops	30 days

Note: When using CONFIRM 2F with other registered pesticides, always refer to rotational restrictions and precautions on the other product's label and comply with the most restrictive rotational guidelines.

USE DIRECTIONS FOR FRUITING VEGETABLES*

Ground Application: Apply a minimum of 10 gallons per acre by conventional ground equipment to young crop and small plants. Apply a minimum of 20 gallons per acre to densely foliated or difficult-to-cover crops to insure thorough coverage.

Aerial Application: For optimum performance a minimum application volume of 10 gallons per acre is recommended. Lower carrier volumes may result in less uniform coverage and reduced efficacy.

Spray Adjuvant: One pint of LATRON CS-7 per 100 gallons of spray mixture or similar spreader-binder is recommended to maximize coverage and distribution of the spray material.

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Alfalfa looper Beet armyworm Black Cutworm Cabbage looper European corn borer Fall armyworm Imported cabbageworm	6.0 to 8.0 (0.09 to 0.12 lb. ai/acre)	For early season applications only to young crop and small plants. Begin applications when first signs of feeding damage appear or when infestations reach threshold levels as defined by Cooperative Extension Service or other qualified professional authorities.	Do not apply more than 16 fluid ounces per application and do not exceed 64 fluid ounces per season. Allow at least 7 days
Southern armyworm Tobacco hornworm Tomato hornworm True armyworm Yellowstriped armyworm	8.0 to 16.0 (0.12 to 0.25 lb. ai/acre)	For mid- to late-season applications and to heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10 to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	to elapse between final application and harvest. See Rotational Crop Restrictions in the body of this labeling.

*The fruiting vegetable grouping includes eggplant, ground cherry, pepino, pepper (bell, chili, cooking), pimento, tomatillo and tomato.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**



© 2000 Rohm and Haas Company
* 02/24/00

NEXT

LABEL

CONFIRM 2F**Supplemental
Labeling****SUPPLEMENTAL LABELING FOR USE OF
CONFIRM® 2F AGRICULTURAL INSECTICIDE
IN POME FRUIT**

EPA Reg. No 707-238

EPA Est. No. 39578-TX-01

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.


USE DIRECTIONS FOR POME FRUITS

(Apples, Crabapples, Loquat, Mayhaw, Pears including oriental, Quince)

Ground Application: Make applications of CONFIRM 2F Agricultural Insecticide by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trellised trees or trees 10 feet tall or less. For trees greater than 10 feet tall, use a minimum of 100 gallons per acre.

Aerial Application: Make applications of CONFIRM 2F Agricultural Insecticide in a minimum of 20 gallons per acre. CONFIRM 2F can be applied by aerial applications when situations warrant. However, this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

Spray Adjuvant: The use of a Spreader-Sticker such as Latron B-1956 or similar Spreader-Sticker to maximize uniform coverage and distribution of the spray material is recommended.



USE DIRECTIONS FOR POME FRUITS (CONTINUED)

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Codling moth (East of the Rockies)	20.0 (0.31 lb. ai/acre)	For each codling moth generation, apply at initiation of egg hatch [150-250 Day Degrees (DD), base 50°F, following biofix*] followed by a second application 10-15 days following the first application (usually 450-550 DD). Additional applications at 10-15 day intervals may be required under high infestations, sustained moth flight, or to ensure coverage of rapidly expanding fruits or foliage.	Do not apply more than 20 oz per acre per application or 120 oz CONFIRM 2F per acre per season. Allow at least 14 days to elapse between final application and harvest.
Codling moth (West of the Rockies) For use against low to moderate infestations in conjunction with alternate control measures such as in established Mating Disruption blocks.	20.0 (0.31 lb. ai/acre)	For each codling moth generation, apply at initiation of egg hatch [150-250 Day Degrees (DD), base 50°F, following biofix*] followed by a second application 10-15 days following the first application (usually 450-550 DD). Additional applications at 10-15 day intervals may be required under high infestations, sustained moth flight, or to ensure coverage of rapidly expanding fruits or foliage.	Do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
Obliquebanded leafroller (West of the Rockies)	20.0 (0.31 lb. ai/acre)	Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending on infestation level. Summer generation: Begin applications at early egg lay through early egg hatch (200-400 DD, base 43°F, following biofix). Make a second application 10-18 days later (usually 650-850 DD). A third application 10-14 days after the second application may be required under high pressure, sustained moth flight or prolonged shoot growth.	
Obliquebanded leafroller	20.0 (0.31 lb. ai/acre)	Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending on infestation level. Summer generation: Begin applications at peak moth flight (200-300 DD, base 43°F, following biofix). Make a second application 7-14 days later (usually 500-600 DD). A third application 10-12 days after the second application (usually 800-900 DD), may be required under high pressure, sustained moth flight or prolonged shoot growth. Fall overwintering generation: Apply to late season larval infestations of overwintering generation to minimize damage to the fruit.	

USE DIRECTIONS FOR POME FRUITS (CONTINUED)

TARGET PESTS	APPLICATION RATE Fluid ounces per acre	APPLICATION TIMING	RESTRICTIONS
Pandemis leafroller	20.0 (0.31 lb. ai/acre)	Spring (overwintering) generation: Make 1 to 2 applications during the pink to petal fall period depending on infestation level. Summer generation: Begin applications at early egg lay through early egg hatch (250-400 DD, base 41°F, following biofix). Make a second application 10 to 18 days later (usually 600-800 DD). Under heavy infestation a third application may be required 10-14 days after the second application.	
Tufted apple bud moth	12.0 - 20.0 (0.19 - 0.31 lb. ai/acre)	First generation: Make an application at 10-30% egg hatch (600-900 DD, base 45°F, after biofix depending on local conditions). A second application at 60-90% egg hatch may be required under heavy infestation levels. Second generation: Make the first application at 20-30% egg hatch (2300-2500 DD). A second application approximately 14 days later may be required under high pressures or sustained moth flight or late maturing varieties.	
Eyespotted bud moth Fruittree leafroller Redbanded leafroller Variegated leafroller	20.0 (0.31 lb. ai/acre)	For control of other leafrollers, begin applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10-14 days to ensure complete coverage of rapidly expanding fruits or foliage.	
Lesser appleworm	20.0 (0.31 lb. ai/acre)	For each generation, apply at initiation of egg hatch before larvae enter the fruit. Make a second application 10-14 days following the first to ensure complete coverage of rapidly expanding fruits or foliage or under conditions of high infestation or sustained moth flight.	
Green fruitworm	20.0 (0.31 lb. ai/acre)	Apply at initiation of egg hatch or at the first sign of larval infestation. A second application may be required 10-14 days following the first application to ensure complete coverage of rapidly expanding fruits or foliage.	

*Biofix is defined as first sustained adult catch in pheromone traps, typically, 5 moths in 3 traps within a seven-day period. Consult State Extension specialists or other qualified authorities for specific information regarding number, placement and management of pheromone traps.

CONDITIONS OF SALE AND WARRANTY


Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**



© 2000 Rohm and Haas Company
02/24/00

NEXT

LABEL

CONFIRM 2F**Supplemental
Labeling**

**SUPPLEMENTAL LABELING FOR USE OF
CONFIRM® 2F AGRICULTURAL INSECTICIDE
IN CHRISTMAS TREES**

**EPA Reg. No 707-238
EPA Est. No. 39578-TX-01**

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions on the container labeling. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemical may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip. Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to inter-tidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment wash-waters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

USE DIRECTIONS FOR CHRISTMAS TREES

Because CONFIRM 2F Agricultural Insecticide must be ingested in order to be effective, it is essential that coverage is thorough and uniform. Higher carrier volumes and higher use rates are recommended for large trees or dense stands and for heavy target pest infestations.

Ground Application: Hydraulic ground sprayers and air blast sprayers should be calibrated to deliver a minimum of 50 gallons per acre. For mist blowers use a minimum of 10 gallons per acre.

Aerial Application: Make applications of CONFIRM 2F in a minimum of 5 gallons per acre. Higher carrier volumes are recommended for dense tree stands, high pest infestations or when environmental conditions are less than ideal for aerial applications.

Target pests	Application Rate (fluid ounces per acre)	Application Timing	Restrictions
(Gypsy moth (<i>Lymantria dispar</i>))	4.0 to 8.0 (0.06 to 0.12 lb. ai/acre)	Apply to early instar (1st, 2nd, or 3rd) larvae. In general, foliage development should be a minimum of 20%.	Do not apply more than 16 fl. Ounces per year. Uniform coverage of the foliage is essential to provide maximum protection from defoliation and reduction of egg mass deposition.
Pine tip moth (<i>Rhyacionia frustrana</i> , <i>R. neomexicana</i> , <i>R. buoliana</i> , <i>R. rigidana</i> , <i>R. subtropica</i>)	8.0 (0.12 lb. ai/acre)	Apply to early instar (1st to 2nd) larvae after each new foliage flush, in general, at approximately 25% shoot expansion.	
Spruce budworm (<i>Choristoneura fumiferana</i> and <i>C. occidentalis</i>) Tussock moth (<i>Dasychira pinicola</i> , <i>Lophocampa maculata</i> , <i>Orgyia pseudotsugata</i> , <i>O. vetusta</i>)	4.0 to 8.0 (0.06 to 0.12 lb. ai/acre)	Make application to 4th to 5th instar larvae that are actively feeding on foliage or are feeding outside the candle caps.	

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

