

PM 22

707-239

11-2-98

1 of 5

8221-B3a
10/13/97

INDAR® 75 WSP

AGRICULTURAL FUNGICIDE

In Water-Soluble Pouches

ACTIVE INGREDIENT

Fenbuconazole

a-[2-(4-chlorophenyl)ethyl]-a-phenyl-1H-1,2,4-triazole-1-propanenitrile.....75%

INERT INGREDIENTS.....25%

TOTAL 100.0%

EPA REG. NO. 707-239

U.S. Patent No. 4,366,165

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with a large amount of water for at least 15 minutes. Consult a physician if irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NET CONTENTS: 1 LB. (0.454 KG.) 8 X 2 OZ. WATER SOLUBLE POUCHES

ACCEPTED

NOV 2 1998

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 707-239

2 of 5

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

In addition to the required PPE, the following is recommended

- Protective Eyewear
- Chemical-resistant headgear for overhead exposure.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high-water mark. Drift or runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of

12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Shoes plus socks
- Waterproof gloves

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry area above freezing. The water-soluble pouch may become brittle at storage temperatures below 32°F but the fungicide is not affected. Do not remove the water-soluble pouches from container except for immediate use.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environment Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty overwrap into application equipment then dispose of overwrap in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Keep dust to a minimum. Appropriate protective equipment must be worn when handling a spill of this material. Remove contaminated clothing promptly. Wash exposed skin areas with soap and water immediately after exposure.

Contain spills immediately with inert materials (e.g. sand, earth). Transfer spilled material and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Refer to Material Safety Data Sheet.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.

GENERAL INFORMATION

INDAR 75WSP fungicide is a protectant fungicide for the control of specific diseases mentioned on this label. INDAR 75WSP fungicide should be used in a regularly scheduled preventive spray program.

A wetting agent such as LATRON B-1956, LATRON CS-7 or other non-polymer containing spray adjuvants approved for use in registered pesticide products should be added to spray solutions according to manufacturers' use instructions to achieve optimum disease control.

USE RATE DETERMINATION- Carefully read, understand and follow label use rates and restrictions. Under low disease conditions, minimum suggested use rates can be used, while maximum suggested label rates and shortened spray schedules are recommended for severe or threatening disease conditions. For proper application, determine the number of acres to be treated, the suggested label use rate and the gallonage to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

MIXING- Always place INDAR 75WSP fungicide into solution prior to adding co-applied agricultural chemicals registered for use on specific crop or crops being treated. Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately ten minutes from the time they are added to the water.

NOTE: Do not use water containing suspended soil particles is used, such as water from ponds, streams or irrigation canals.

HANDLING - The enclosed pouches of INDAR 75WSP fungicide are water-soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

APPLICATION

Do not make ground or aerial applications within 75 feet of streams, rivers, ponds, lakes or reservoirs.

Ground- Thorough coverage sprays generally result in optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and tractor speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

Aerial - For those crops where applications are not limited to ground equipment, apply in a minimum of 10 gallons of water per acre. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to human or animals. Flagmen and loaders should wash thoroughly before eating, drinking or smoking and at the end of each days spray operation.

Aerial Spray Drift Management- AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

Chemigation - Do not apply this product through any type of irrigation system.

COMPATIBILITY - INDAR 75WSP fungicide is compatible with most commonly used agriculture fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

OTE: INDAR 75WSP fungicide is compatible with boron and spray oils; however, the water-soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray-mixtures.

USE DIRECTIONS FOR PECANS

INDAR 75WSP is a protectant fungicide. Best disease control is achieved on a protectant application schedule using thorough coverage sprays. When necessary, add LATRON B-1956, LATRON CS-7 or other non-polymer containing spray adjuvants approved for use on registered pesticide products should be added to spray solutions according to manufacturers' use instructions to achieve optimum disease control.

DISEASE	RATE OF INDAR 75WSP PER ACRE	USE RECOMMENDATIONS	RESTRICTIONS
Dormant Sprout Blight (<i>Dothiorella</i>) Gnomonia Leafspot (<i>Gnomonia</i> sp.) Leaf Spot (<i>Mycophtheria</i>) Leaf Spot (<i>Microsporia</i>) Pecan Blight (<i>Microsporia</i>) Scab (<i>Cladosporium</i> sp.) Vein Spot (<i>Gnomonia</i> sp.)	2 to 2.5 ounces (0.09 to 0.12 lb. active)	Begin applications at bud break and continue at 10 to 14-day intervals through pollination. Apply at 14- to 21-day intervals through the cover sprays.	Do not apply after shuck split or within 28 days of harvest. Do not apply more than 1 pound (0.75 lb. active) per acre per season. Do not graze livestock in treated areas or feed crops grown in treated areas to livestock.

5 of 5

USE DIRECTIONS FOR STONEFRUITS

Use one 2-ounce pouch of INDAR 75WSP per acre in a minimum of 50 gallons by ground or 10 gallons by air. INDAR 75WSP fungicide may be applied up to the day of harvest.

INDAR 75WSP is a protectant fungicide. Best disease control is achieved when a protectant application schedule is followed.

A wetting agent such as LATRON B-1956, LATRON CS-7 or other non-polymer containing spray adjuvant approved for use in registered pesticides products should be added to spray solutions according to manufacturers' use instructions to achieve optimum disease control.

CROP	DISEASE	USE RECOMMENDATIONS	RESTRICTIONS
Apricots	Blossom Blight (<i>Monilinia spp.</i>)	Begin applications at early red bud stage before infection occurs. If conditions are favorable for disease development, apply again at full bloom and at petal-fall.	For all crops, do not apply more than 1 lb. (0.75 lbs. active) INDAR 75WSP fungicide per acre per season. For all crops, do not graze livestock in treated areas or feed cover crops grown in treated areas to livestock.
	Fruit Brown Rot (<i>Monilinia spp.</i>)	Begin applications 2 to 3 weeks before harvest using a 7 to 10 day spray interval.	
Cherries	Blossom Blight Fruit Brown Rot	Refer to apricots.	
	Leaf Spot (<i>Blumeriella sp.</i>)	Follow blossom blight schedule and continue application at 10 to 14 day intervals. Additional foliar applications may be made after harvest.	
Nectarines	Blossom Blight Fruit Brown Rot	Refer to apricots.	
Peaches	Blossom Blight Fruit Brown Rot	Refer to apricots.	
	Scab (<i>Cladosporium sp.</i>)	Begin applications at shuck split. Make 2 to 3 subsequent thorough coverage applications at 10 to 14 day intervals.	

8221-B3a
10/13/97
EC/rdm