



United States Environmental Protection Agency
Office of Pesticide Programs (H7505C)
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number
197395

Section I

| | | |
|--|--|--|
| 1. Company/Product Number ██████████ 707-253 | 2. EPA Product Manager C. Welch - 21 | 3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted |
| 4. Company/Product (Name) Eagle WSP Fungicide/ Systhane WSP Ornamental Fungicide | PM# 21 | |
| 5. Name and Address of Applicant (Include ZIP Code) Rohm and Haas Company 100 Independence Mall West Philadelphia, PA 19106-2399 <input type="checkbox"/> Check if this is a new address | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____ | |

Section II

| | |
|--|--|
| <input type="checkbox"/> Amendment - Explain below | <input type="checkbox"/> Final printed labels in response to Agency letter dated _____ |
| <input type="checkbox"/> Resubmission in response to Agency letter dated _____ | <input type="checkbox"/> "Me Too" Application. |
| <input checked="" type="checkbox"/> Notification - Explain below. | <input type="checkbox"/> Other - explain below. |

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Changes included on these labels are those we discussed at the meeting in your office on 04APR96.

NOTIFICATION OKAY Advt 8/16/96
AUG 26 1996

Section III

| | | | | | |
|---|---|--|---|------------------------|-------------------|
| 1. Material This Product Will Be Packaged In: | | | | 2. Type of Container | |
| Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No | Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No | Metal Plastic Glass Paper Other (Specify) | | |
| * Certification must be submitted. | | If "Yes," Unit Package wgt. | No. per container | If "Yes," Package wgt. | No. per container |
| 3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container | 4. Size(s) of Retail Container | | 5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product | | |
| 6. Manner In Which Label Is Affixed To Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled | | | <input type="checkbox"/> Other (_____) | | |

Section IV

| | | |
|---|------------------------------------|---|
| 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.) | | |
| Name Richard D. Costlow, Ph.D., DABT | Title Product Registration Mgr. | Telephone No. (Include Area Code) 215-592-3581 |

Certification

| | | |
|--|---------------------------------------|---|
| I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. | | 6. Date Application Received (Stamped) |
| 2. Signature | 3. Title Product Registration Mgr. | |
| 4. Typed Name Richard D. Costlow | 5. Date May 24, 1996 | |

2911

RECD EPA/OPP/DPD1

8777-BIW
5/16/96

'96 JUN -4 P2 :23

SYSTHANE® WSP



ORNAMENTAL FUNGICIDE
-In Water-Soluble Pouches-

NOT FOR USE IN NASSAU OR SUFFOLK COUNTIES, NEW YORK.

ACTIVE INGREDIENT
MYCLOBUTANIL

a-butyl-a-(chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile

INERT INGREDIENTS

TOTAL 100%

NOTIFICATION
AUG 26 1996

NOTIFICATION
AUG 26 1996

EPA REGISTRATION NO. 707-253

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use. Use Restrictions and Storage and Disposal instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with a large amount of water for at least 15 minutes. Call a physician if irritation persists.

IF INHALED: Move victim to fresh air.

IF ON SKIN: Wash affected skin thoroughly with soap and water.

IF SWALLOWED: Dilute by giving two glasses of water to drink and call a physician. Never give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not swallow, get in eyes, on skin or breathe spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards(WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24** hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls •Waterproof gloves •Shoes plus socks •Protective eyewear
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Seed treatments and professional applications to lawn grasses, golf courses, industrial (office park), municipal and residential lawns are not within the scope of the Worker Protection Standard.

- Keep unprotected persons out of treated area until sprays have dried.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry area above freezing. The water-soluble pouch may become brittle at storage temperatures below 32°F. but the fungicide is not affected. Do not remove the water-soluble pouches from the container except for immediate use.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Completely empty container into application equipment. Dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Wear eye protection. Wear protective clothing. Spray water on powder and dust. Scoop or shovel solid material into a suitable container for recovery or disposal. Keep dust to a minimum. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit. Remove contaminated clothing promptly and wash affected skin areas with soap and water. Thoroughly launder clothing before reuse. Refer to Precautionary Statements.

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

GENERAL INFORMATION

SYSTHANE WSP fungicide is a systemic, protectant and curative fungicide recommended for the control of specific diseases mentioned on this label. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled preventative spray program.

USE RATE DETERMINATION - Carefully read, understand and follow label use rates and restrictions.

Under low disease conditions, minimum label use rates per application can be used while maximum label rates and shortened spray schedules are recommended for severe or threatening disease conditions.

For proper application, determine the number of acres to be treated, the recommended label use rate and the gallonage to be applied per acre. Prepare only the amount of spray solution required to treat the measured acreage. Careful calibration of spray equipment is recommended prior to use.

HANDLING - The enclosed pouches of SYSTHANE WSP fungicide are water-soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal overwrap bag to protect remaining unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

MIXING- Always place SYSTHANE WSP fungicide into solution prior to adding co-applied materials registered for use on specific crops. Add the required number of unopened pouches as determined by the dosage recommendations into the spray tank with agitation. Depending on the water temperature and the degree of agitation, the pouches should dissolve completely within approximately ten minutes from the time they are added to the water.

COMPATIBILITY - SYSTHANE WSP fungicide is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients and spray adjuvants. When preparing tank mixes, user should consult spray compatibility charts or State Cooperative Extension Service Specialists prior to actual use.

NOTE: SYSTHANE WSP fungicide is compatible with boron and spray oils, however, the water soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures.

5911

APPLICATION PROCEDURES

Ground Equipment: Application equipment should be properly calibrated and provide a uniform spray coverage throughout the plant canopy.

Handgun or Pressurized Sprayers: Thoroughly spray plant foliage until run-off.

Sprinkler Irrigation: SYSTHANE WSP fungicide for ornamental plants must be applied on a regular protectant fungicide schedule, *not an irrigation schedule*. Apply only through solid set or hand move sprinkler irrigation systems. Do not apply product through any other type of irrigation system.

Lack of fungicidal effectiveness can result from nonuniform distribution of treated water. Greatest efficacy is achieved when the application of treated water does not exceed 1/4 inch per acre per application.

If you have questions about calibration, you should contact State Extension Service specialists or equipment manufacturers.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Before applying SYSTHANE fungicide to ornamental plants through sprinkler irrigation equipment, the chemigation system must meet the following specifications:

- Public water system means a system for the provision to the public of piped water for human consumption is such system that has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

- Chemigation systems connected to public water systems must contain a functional reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the pipe fill and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.

- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

- Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Solid-set and Hand Move Irrigation Equipment:

- Determine area covered by sprinkler.

- Fill injector solution tank with water and adjust flow rate to use the contents over a 10 to 30 minute interval.

- Determine the amount of SYSTHANE WSP fungicide required to treat area.

- Add the required amount of SYSTHANE WSP fungicide into the same quantity of water used to calibrate the injection equipment.

- Maintain constant solution tank agitation during the injection period.

- Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.

- Inject SYSTHANE WSP fungicide at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention.

- Stop injection equipment after treatment is completed. Continue to operate the system until the SYSTHANE WSP fungicide solution has cleared the last sprinkler head.

6911
D

USE DIRECTIONS FOR ORNAMENTALS

SYSTHANE WSP is a systemic fungicide having protectant and curative properties. Best control of labeled diseases is achieved in thorough coverage sprays, applied to runoff on a protectant application schedule. For greenhouse or field-grown ornamentals plants, SYSTHANE WSP fungicide should be used at 2.0 ounces (1 pouch) product per 50 gallons spray solution, unless otherwise directed, and applied on a 10 to 14 day application schedule. For concentrate sprays adjust amount accordingly. The addition of a nonphytotoxic spray adjuvant will improve spray coverage and fungicidal performance. Treated plants should be maintained in a vigorous growing condition. Plants under nutritional or water stress will not respond as well to treatment as well-maintained plants. Overdosage of SYSTHANE WSP fungicide can result in observable foliar greening and shortened internodes. If this condition is observed, reduce the fungicide use rate but do not extend the indicated application schedule.

CROP TOLERANCE

Treated plants should be maintained in a vigorous growing condition. Plants under nutritional or water stress will not respond as well to treatment as well-maintained plants. Overdosage can result in observable foliar greening and shortened internodes. If this condition is observed, reduce the use rate but do not extend the application schedule.

Plant tolerances are acceptable in the specific plants listed on this label. It is not possible to evaluate all ornamental plant species or varieties for tolerance to SYSTHANE WSP fungicide. The user should test for possible phytotoxic responses by treating a limited number of plants at recommended use rates prior to initiating large scale use.

The effects of spraying SYSTHANE WSP fungicide in combination with plant growth regulators are not fully understood at this time. If the use of a plant growth regulator is planned in an area being treated, the user should test for possible enhanced growth regulatory effects by treating a small number of plants at the recommended use rates of all products prior to initiating large scale use. Since the effectiveness of such products depends not just on plant species or cultivar but also weather and seasonal differences (e.g., daylight hours) it is recommended that tests be repeated on previously tested varieties as environmental factors change and that observations for growth regulator responses be made at regular intervals.

SPECIFIC USE DIRECTIONS FOR CHRYSANTHEMUM

FOLIAR SPRAYS: Best control is achieved by thorough coverage sprays, applied to runoff on a protectant application schedule. Use SYSTHANE WSP fungicide at a rate of 2 ounces per 50 gallons of spray solution (do not apply more than 6 ounces per acre per application or more than 5 pounds per acre per year). Applications should be made on a 10 to 14-day schedule (not to exceed 21 days).

PRESTICK DIP TREATMENT: Chrysanthemum cuttings may be treated by a dip procedure prior to planting as follows. Prepare a dip suspension at a concentration equivalent to 2 ounces SYSTHANE WSP fungicide per 50 gallons of water. Cuttings must be fully submerged in the dip suspension until wet throughout. (Cuttings should not remain submerged longer than 2 minutes). If cuttings are dipped, this procedure will represent the first spray under the quarantine program. Used dip suspension should be disposed of if it becomes contaminated with soil, plant debris or other foreign matter. Used dip suspension can be disposed of by spraying onto registered crops (but not onto previously dipped cuttings) after filtering, or in a manner consistent with local, state, and federal guidelines.

NOTE: All infected plant material must be destroyed if your state is under quarantine directive.

RESTRICTIONS

- Do not apply more than 6 ounces (2.4 ounces active) per acre per application.
- Do not apply more than 5 pounds (2 pounds active) per acre per year.
- Do not use treated plant materials for food or feed.

79 11

ORNAMENTALS -cont.-

| CROP | DISEASES CONTROLLED | REMARKS | CROP SPECIFIC RESTRICTIONS |
|--------------------------------------|---|--|----------------------------|
| Abelia | Cercospora Leaf Spot Powdery Mildew | | |
| Abutilon (Flowering Maple) | Rust | | |
| Acalypha (Copper-Leaf) | Cercospora Leaf Spot Powdery Mildew | | |
| Achillea (Yarrow) | Powdery Mildew Rust | | |
| African Violet | Powdery Mildew | | |
| Ageratum | Rust | | |
| Alder | Powdery Mildew Rust | | |
| Almond, Flowering | Blossom Blight (monilinia spp.) | Apply prebloom, 50% bloom and at petal fall | |
| Amelanchier (Juneberry, Shadbush) | Fabraea Leaf Spot Powdery Mildew Rust | | |
| Amorpha (False Indigo) | Cercospora Leaf Spot Powdery Mildew Rust | | |
| Anemone | Rust | | |
| Angelica | Cercospora Leaf Spot Rust | | |
| Ash | Rust | | |
| Aster | Rust | | |
| Azalea | Petal Blight (Ovulinia spp.) Powdery Mildew | Begin applications when flowers start to exhibit color. | |
| Barberry | Powdery Mildew Rust | | |
| Begonia | Powdery Mildew | | |
| Bellflower | Cercospora Leaf Spot Powdery Mildew Rust | | |
| Birch | Rust | | |
| Bittersweet | Powdery Mildew | | |
| Buckeye | Powdery Mildew | | |

ORNAMENTALS -cont.-

| CROP | DISEASES CONTROLLED | REMARKS | CROP SPECIFIC RESTRICTIONS |
|-----------------------|--|---|---|
| Buttonbush | Cercospora Leaf Blight Powdery Mildew Rust | | |
| Calendula | Cercospora Leaf Spot | | |
| California Poppy | Powdery Mildew | | |
| Carnation | Powdery Mildew Rust | | |
| Catalpa | Cercospora Leaf Spot Powdery Mildew | | |
| Cherry, Flowering | Leaf Spot Powdery Mildew | | |
| Chestnut, Horse | Powdery Mildew | | |
| China Aster | Rust | | |
| Chokeberry | Rust Twig and Fruit blight | | Fruit may not be used for food or feed. |
| Chrysanthemum | Rust White Rust | See specific use directions. | |
| Columbine | Rust | | |
| Comflower | Rust | | |
| Cosmos | Powdery Mildew | | |
| Crabapple, Flowering | Powdery Mildew Rust Scab | | |
| Crepe-Myrtle | Powdery Mildew | | |
| Dahlia | Powdery Mildew | | |
| Delphinium | Powdery Mildew Rust | | |
| Dogwood | Anthracnose Powdery Mildew | | |
| Douglas Fir (Nursery) | Needle Rust | Apply 6 to 10 ounces per acre starting early spring. Continue applications at 2 to 3 week intervals until the threat of infection has passed. Spray adjuvants must be added to spray solutions to obtain good spray coverage and disease control. | |
| Dianthus | Rust | | |
| Elm | Powdery Mildew | | |
| Euonymus | Powdery Mildew | | |
| Fleabane | Cercospora Leaf Spot Powdery Mildew Rust | | |
| Four O'Clock | Rust | | |

ORNAMENTALS -cont-

| CROP | DISEASES CONTROLLED | REMARKS | CROP SPECIFIC RESTRICTIONS |
|-------------------------|---|--|---|
| Fuchsia | Rust | | |
| Gaillardia | Powdery Mildew Rust | | |
| Gardenia | Powdery Mildew Rust | | |
| Geranium | Powdery Mildew Rust | | |
| Gourd, Ornamental | Powdery Mildew | | |
| Grape Leaf Ivy | Powdery Mildew | | |
| Hackberry | Cercospora Leaf Spot Powdery Mildew | | |
| Hawthorn | Fabraea Leaf Spot Powdery Mildew Rust Scab | | |
| Hollyhock | Powdery Mildew Rust | | |
| Honeysuckle | Cercospora Leaf Spot Powdery Mildew | | |
| Hydrangea | Cercospora Leaf Spot | | |
| Iris | Didymellina Leaf Spot Rust | Apply 4 oz. per 50 gallons spray solution. | |
| Leucothoe | Cercospora Leaf Spot | | |
| Lilac | Powdery Mildew | | |
| Loblolly Pine (Nursery) | Fusiform Rust | Refer to Douglas Fir | |
| Locust | Powdery Mildew | | |
| Maple | Powdery Mildew | | Treated trees may not be used for syrup production. |
| Marigold | Cercospora Leaf Spot Rust | | |
| Mock-Orange | Powdery Mildew Rust | | |
| Moonflower | Rust | | |
| Mountain Laurel | Cercospora Leaf Spot Ovulinia Petal Blight Powdery Mildew | Refer to Azalea | |
| Nephtytis | Cephalosporium Leaf Spot | | |
| Ninebark | Rust | | |
| Oak | Powdery Mildew | | |
| Pansy | Powdery Mildew Rust | | |

ORNAMENTALS -cont.-

| CROP | DISEASES CONTROLLED | REMARKS | CROP SPECIFIC RESTRICTIONS |
|------------------------|--|--|---|
| Pear, Flowering | Powdery Mildew Rust Scab | | |
| Petunia | Powdery Mildew Rust | | |
| Phlox | Cercospora Leaf Spot Powdery Mildew Rust | | |
| Photinia | Entomosporium Leaf Spot Powdery Mildew Rust | | |
| Poinsettia | Powdery Mildew | | |
| Privet | Cercospora Leaf Spot Powdery Mildew | | |
| Pyracantha (Firethorn) | Fusicladium Scab | | |
| Quince, Flowering | Blossom and Twig Blight Cercospora Leaf Spot Fabraea Leaf Spot Rust | | |
| Rhododendron | Cercospora Leaf Spot Ovulinia Petal Blight Powdery Mildew | Refer to Azalea | |
| Rose | Black Spot Powdery Mildew Rust | Apply on a 7 to 10 day schedule. In areas where black spot is not a problem, spray intervals may be increased to a maximum of 14 days. | |
| Russian Olive | Cercospora Leaf Spot Rust | | |
| Salvia | Powdery Mildew Rust | | |
| Slash Pine (Nursery) | Fusiform Rust | Refer to Douglas Fir | |
| Smoke-Tree (Cotinus) | Cercospora Leaf Spot Rust | | |
| Snapdragon | Powdery Mildew Rust | | |
| Spirea | Powdery Mildew | | |
| Sunflower | Cercospora Leaf Spot Powdery Mildew Rust | | Seeds from treated plants may not be used for food or feed. |
| Sycamore | Powdery Mildew | | |

11 2 11

ORNAMENTALS -cont.-

| GROP | DISEASES CONTROLLED | REMARKS | CROP SPECIFIC RESTRICTIONS |
|-----------------|--|---------|--|
| Trumpet-Creeper | Cercospora Leaf Blight Powdery Mildew | | |
| Viburnum | Powdery Mildew Rust | | |
| Walnut | Powdery Mildew | | Nuts from treated trees may not be used for food purposes. |
| Willow | Powdery Mildew | | |
| Zinnia | Cercospora Leaf Spot Powdery Mildew | | |

SYSTHANE IS A TRADEMARK OF ROHM AND HAAS COMPANY

8777-B1W
5/16/96