

PM 01

707-261

8/26/97

1/7

	<p align="center">U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460</p>	<p>EPA Reg. Number: 707-261</p>	<p>Date of Issuance: AUG 26 1997</p>
		<p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p align="center">(under FIFRA, as amended)</p>	
		<p>Name of Pesticide Product: RH-144228</p>	
<p>Name and Address of Registrant (include ZIP Code): Rohm and Haas Company 100 Independence Mall West Philadelphia, PA 19106-2399</p>			
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p>			
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p align="center">This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the following labeling change before you release the product for shipment: Add the phrase "EPA Registration No. 707-261". 3. Submit one copy of the final printed label for the record. <p>If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e).</p>			
<p>Signature of Approving Official: <i>Mary J. Waller</i></p>		<p>Date: 8/26/97</p>	

Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Mary L. Waller

Mary L. Waller
Acting Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosure

7505C:C.Grable:cg:8/20/97



RH-144228

Fungicide

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

ACTIVE INGREDIENT	
SULFUR.....	96.75%
MYCLOBUTANIL	
a-butyl-a-(4-chlorophenyl)-1H-1,2,4-triazole-1-propanenitrile...	0.50%
INERT INGREDIENTS	2.75%
TOTAL	100.00%

ACCEPTED
with **COMMENTS**
In EPA Letter Dated
AUG 26 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 707-561

EPA REG NO 707-

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Flush eyes with a large amount of water for at least 15 minutes. Call a physician if irritation persists.
- IF INHALED:** Move victim to fresh air.
- IF ON SKIN:** Wash affected skin thoroughly with soap and water.
- IF SWALLOWED:** Dilute by giving two glasses of water to drink, induce vomiting and call a physician. Never give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Do not swallow, get in eyes, on skin or breathe spray mist.

Personal Protective Equipment (PPE)

- Applicators and other handlers must wear:
- o Long-sleeved shirt and long pants
 - o Waterproof gloves
 - o Shoes plus socks
 - o Chemical-resistant headgear for overhead exposure

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/ maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards(WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

- Users should:
- o Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - o Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - o Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

4/7

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

PHYSICAL HAZARD

Sulfur dust suspended in air ignites easily. Keep away from heat, sparks or flame. Do not smoke while applying this product.

GENERAL INFORMATION

RH-144228 fungicide is recommended for the control of specific diseases mentioned on this label. Optimum disease control is achieved when the fungicide is applied in a regularly scheduled program.

USE RATE DETERMINATION

Carefully read, understand and follow label use rates and restrictions. Under low disease conditions minimum label use rates per application can be used, while maximum label rates and shortened application schedules are recommended for severe or threatening disease conditions and sensitive varieties.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

CHEMIGATION

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- o Long-sleeved shirt and long pants
- o Waterproof gloves
- o Shoes plus socks
- o Chemical-resistant headgear for overhead exposure

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only and keep sealed. Use caution when moving, opening, closing or pouring.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty container into application equipment. Dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Wear eye protection. Wear protective clothing. Spray water on powder and dust. Scoop or shovel solid material into a suitable container for recovery or disposal. Keep dust to a minimum. Flush contaminated area with a large amount of water to a chemical or sanitary sewer containing a settling pit. Remove contaminated clothing promptly and wash affected skin areas with soap and water. Thoroughly launder clothing before reuse. Refer to Precautionary Statements.

6/7

USE RECOMMENDATIONS

This material may be applied with ground equipment or aircraft depending upon local conditions.

IMPORTANT: Applications of sulfur should not be made just prior to or during periods of high summer temperatures. Do not make applications within 2 weeks of an oil spray. Do not use or allow this material to drift onto sulfur sensitive Melons, Squash, Cucumbers, Apricots or other plants which may be injured by sulfur. When growing crops for processing, consult the processor before applying sulfur..

Crop	Disease	Use Rate per Acre	Use Recommendations	Restrictions
Almonds	Brown Rot Blossom Blight (<i>Monilinia</i> spp.)	40 lb	Begin applications at prebloom and continue on a 7 to 10 day interval	Do not make more than 3 applications per season or apply more than 120 lbs. per acre. Do not apply within 90 days of harvest.
Apples	Scab (<i>Venturia</i> sp.) Powdery Mildew (<i>Podosphaera</i> sp.)	50	Begin applications at green tip and continue applications on a 7 to 10 day schedule.	Do not apply more than 400 lbs. per acre per season. Do not apply within 14 days of harvest.
Cherries	Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Cherry Leaf Spot (<i>Blumeriella</i> sp.) Powdery Mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)	30	Begin applications at prebloom and continue on a 7 to 10 day interval	Do not apply more than 240 lbs. per acre per season. Applications may be made to within 24 hours of harvest.
Grapes	Powdery Mildew (<i>Uncinula</i> sp.)	10 to 20	Make the first application when shoots are 6 to 8 in. long. Make a second application when shoots are 12 to 15 in. long and a third about 14 days later. Repeat applications at 7 to 20 day intervals to protect new growth. Avoid over-applications in hot weather. Concord and other <i>Vitis labrusca</i> type grapes may be injured by sulfur.	Do not apply more than 120 lbs. per acre per season. Do not apply within 14 days of harvest.
Nectarines	Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Powdery Mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)	30	Begin applications at prebloom and continue on a 7 to 10 day interval	Do not apply more than 240 lbs. per acre per season. Applications may be made to within 24 hours of harvest.
Peaches	Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Powdery Mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)	30	Begin applications at prebloom and continue on a 7 to 10 day interval	Do not apply more than 240 lbs. per acre per season. Applications may be made to within 24 hours of harvest.
Plums and Prunes	Brown Rot Blossom Blight (<i>Monilinia</i> spp.) Rust (<i>Tranzschelia</i> sp.)	30	Begin applications at prebloom and continue on a 7 to 10 day interval	Do not apply more than 210 lbs. per acre per season. Applications may be made to within 24 hours of harvest.

NOTE: If fungi displaying reduced sensitivity to the DMI fungicides are known or suspected to be in the crop being treated, do not exceed a 7-day application interval.

GENERAL USE RESTRICTIONS:

Ten (10) pounds of this product contains 0.05 lb. active myclobutanil fungicide. If other myclobutanil fungicide applications are also made to treated crops, do not exceed the total seasonal pounds active myclobutanil per acre, per season.

R2b
EC/rdm
8/20/97
7-7700

7/7

