

707-269

5/4/2001

1/8

8359-M2b  
3/26/01



**INTREPID™ 80WSP**  
**AGRICULTURAL INSECTICIDE In Water-Soluble Pouch**

**ACTIVE INGREDIENT**

Methoxyfenozide

Benzoic acid, 3-methoxy-2-methyl-, 2-(3,5-dimethylbenzoyl)-2-(1,1-dimethylethyl) hydrazide ..... 30%

INERT INGREDIENTS..... 20%

TOTAL 100%

EPA REG. NO. 707-269

EPA EST. NO.

NOTICE: Before using this product, read the entire Precautionary Statements, Conditions of Sale and Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions. If the Conditions of Sale and Warranty are not acceptable, return the product unopened within thirty days of purchase to the place of purchase.

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**

FIRST AID	
IF IN EYES	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
IF INHALED	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
IF SWALLOWED	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>HOT LINE NUMBER</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 215-592-3000 day or night, for emergency treatment information	

**ACCEPTED**  
 MAY 4 2001  
 Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 707-269

NET CONTENTS: 1.5 LBS. (6-4 OUNCE WATER-SOLUBLE POUCHES)

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6), the handler PPE requirements may be reduced or modified as specified in the WPS.

Water-soluble pouches when used correctly, qualify as a closed loading system under the WPS. Handlers handling this product while it is enclosed in intact water-soluble pouches may elect to wear reduced PPE of long-sleeved shirt, long pants, shoes and socks.

**IMPORTANT:** When reduced PPE is worn because of a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as spill or equipment break-down.

<p><b>USER SAFETY RECOMMENDATIONS</b></p> <p>Users should:</p> <ul style="list-style-type: none"> <li>• Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing immediately if pesticide gets inside. Then wash thoroughly with soap and water and put on clean clothing.</li> </ul>
--

**ENVIRONMENTAL HAZARDS**

Drift and runoff from applications of this product may be hazardous to sensitive aquatic invertebrates in water bodies adjacent to the treatment area. For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Methoxyfenozide can contaminate surface water through spray drift. Under some conditions, methoxyfenozide may also have a high potential for runoff into surface water (primarily via dissolution in runoff water), for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas over-laying extremely shallow ground water, areas with in-field canals or ditches that drain to overlying tile drainage systems that drain to surface water.

Do not cultivate within 10 feet of aquatic areas to allow growth of a vegetative filter strip.

Do not apply by ground within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds; estuaries and commercial fish farm ponds.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift.

Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

<p><b>AGRICULTURAL USE REQUIREMENTS</b></p> <p>Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.</p> <p>Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.</p> <p>PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:</p> <ul style="list-style-type: none"> <li>• Coveralls</li> <li>• Water-proof gloves</li> <li>• Shoes plus socks</li> </ul>
--

**STORAGE AND DISPOSAL**

**STORAGE:** Store in a cool, dry area above freezing. Do not store in sunlight. The water-soluble pouch may become brittle at storage temperatures below 32°F. but the insecticide is not affected. Do not remove the water-soluble pouches from the container except for immediate use.

**PESTICIDE DISPOSAL:** Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

**CONTAINER DISPOSAL:** Completely empty container into application equipment. Dispose of in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED:**

Evacuate the spill area. Wear full protective equipment. Transfer spilled material into a suitable container for recovery or disposal. Keep dust to a minimum. . Keep spills and cleaning runoff out of municipal sewers and open bodies of water.

Remove contaminated clothing and wash affected skin areas with soap and water. Refer to Precautionary Statements.

**CONDITIONS OF SALE AND WARRANTY**

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

**GENERAL INFORMATION**

INTREPID 80 WSP Agricultural Insecticide belongs to the diacylhydrazine class of insecticides and has a novel mode of action that mimics the action of the molting hormone of Lepidopterous (moths, butterflies) larvae.

Upon ingestion, larval stages of the order Lepidoptera undergo an incomplete and developmentally premature molt which is ultimately lethal. This process interrupts and rapidly halts their feeding. Feeding typically ceases within hours of ingestion although complete mortality of the larvae may take several days. Affected larvae often become lethargic and develop discolored areas or bands between segments.

INTREPID 80 WSP has virtually no effect on any order of insects or Arthropods except the Lepidoptera, making it an ideal tool for Integrated Pest Management. This selectivity allows beneficial insects (including bees) and other Arthropods to function unimpeded in the management of secondary pests while INTREPID 80 WSP provides control of troublesome Lepidoptera pests.

**USE RATE DETERMINATION**

Carefully read, understand and follow label use rates, recommendations and restrictions. Apply the amount specified in the following table with properly calibrated aerial or ground spray equipment. Prepare only the amount of spray solution required to treat the measured acreage. The low rates may be used for light infestations of the target lepidopterous species and the higher rates for moderate to heavy infestations. INTREPID 80 WSP Agricultural Insecticide may be applied in either dilute or concentrate sprays so long as the application equipment is calibrated and adjusted to deliver thorough, uniform coverage. Use the specified amount of INTREPID 80 WSP per acre regardless of the spray volume used.

**COMPATIBILITY**

INTREPID 80 WSP Agricultural Insecticide is believed to be compatible with most commonly used agricultural fungicides, insecticides, growth regulators, foliar fertilizers and spray adjuvants. If in doubt, the user should consult spray compatibility charts or State Cooperative Extension Specialists prior to preparing tank mixtures.

**NOTE:** INTREPID 80 WSP Agricultural Insecticide is compatible with boron and spray oils; however, the water soluble pouches must be completely dissolved before adding spray oils or products containing boron to spray mixtures

4/8

## SPRAY ADJUVANTS

The addition of agricultural adjuvants to INTREPID 80 WSP Agricultural Insecticide sprays will improve initial spray deposits, redistribution and weatherability. The following spray adjuvants have been especially formulated to optimize the performance of foliar-applied agricultural chemicals:

- LATRON B-1956® -A water-dispersible, resin-based nonionic spreader-sticker that resists re-wetting and removal by rain. Effective with dilute sprays applied by ground equipment.
- LATRON CS-7® -A spreader-binder designed specifically for use in concentrate and low volume sprays applied by aircraft or ground equipment.

## MIXING

Fill spray tank at least one-third or one-half full before adding water soluble pouches to the spray tank. Always place the INTREPID 80 WSP pouches in the spray tank and insure that the pouches are completely dissolved before adding other co-applied products. Add the required number of pouches as determined by the dosage recommendations into the tank with continuous agitation. Pouches should completely dissolve within approximately five minutes depending on the water temperature and degree of agitation.

## HANDLING

The enclosed pouches of INTREPID 80 WSP Agricultural Insecticide are water soluble. Do not allow pouches to become wet prior to adding to the spray tank. Do not handle the pouches with wet hands or wet gloves. Always reseal the overwrap bag to protect unused pouches. Do not remove water-soluble pouches from overwrap except to add directly to the spray tank.

## APPLICATION TIMING

The activity of INTREPID 80 WSP Agricultural Insecticide is expressed primarily through ingestion by the target larvae. Consequently, the timing of application is dependent on the feeding behavior of the target pest. For cryptic (internal) feeding larvae, application must be made prior to the time that surface feeding occurs i.e., just prior to initiation of egg hatch. For foliar or surface feeding larvae, application may be made while active feeding is occurring.

Re-application may be required to protect new flushes of foliage, rapidly expanding fruit or extended infestations. The re-application interval will vary depending on how rapidly the crop is growing, the generation time of the target pest and the duration of the infestation.

INTREPID 80 WSP Agricultural Insecticide is effective against all larval instars; however, it is good practice to make applications to early instars to minimize feeding damage. For best results, begin applications when threshold levels of moths, eggs or larvae occur. Consult the Cooperative Extension Service, or other qualified professional authorities to determine the appropriate threshold and timing for application in your area.

## APPLICATION INSTRUCTIONS

INTREPID 80 WSP Agricultural Insecticide must be ingested by insect larvae to be fully effective. Applications must be in a manner that assures uniform and thorough coverage. Higher water volume and increased spray pressure generally provide better coverage.

When using an airblast sprayer, coverage is also improved by operation of the sprayer at ground speeds that assure that the air volume within the tree canopy is completely replaced by the output from the airblast sprayer. Making applications in an alternate row middle pattern may result in less than satisfactory coverage and poor performance in conditions of high pest infestation levels, extremely large trees and/or dense foliage.

To avoid drift and achieve maximum performance of this product, make ground applications when the wind velocity favors on-target product depositions (3-10 mph). Do not apply when wind velocity exceeds 10 mph. Shut off the sprayer when turning at row ends. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind directions are toward the aquatic area.

A minimum of six hours drying time is required between the completion of application and the onset of precipitation to ensure retention of the spray deposit.

**RESISTANCE MANAGEMENT**

Any insect population may contain individuals that are naturally resistant to a specific pesticide, therefore, the use of any one insecticide against many consecutive generations of pest can result in the development of resistance problems. To prevent or delay the development of resistance, Rohm and Haas Company recommends rotation of INTREPID 80 WSP with insecticides of alternate modes of action and the utilization of Integrated Pest Management practices such as routine monitoring, the use of treatment thresholds to time applications and cultural and biological controls wherever possible. We further recommend that INTREPID 80 WSP not be used on more than three consecutive generations of a pest. Since the development of resistance cannot be predicted, we suggest that you consult local or State Extension personnel or your local Rohm and Haas Company representative for resistance management guidance appropriate to your crop, locality and production practices.

**CHEMIGATION**

Do not apply this product through any type of irrigation system.

**ENDANGERED SPECIES**

Do not use this product in Door County in the State of Wisconsin.

Do not apply this product within One mile of sandy habitats that support wild lupine plants in the following States/Counties:

MICHIGAN

Allegan  
Monroe  
Montcalm  
Muskegon  
Newaygo  
Oceana

WISCONSIN

Adams	Juneau	Sauk
Burnett	Marquette	Shawano
Chippewa	Menominee	Trempeleau
Clark	Monroe	Waupaca
Dunn	Oconto	Waushara
Eau Claire	Outagamie	Wood
Green Lake	Polk	
Jackson	Portage	

**ROTATIONAL CROP RESTRICTIONS**

The following rotational crops may be planted at intervals defined below following the final application of INTREPID 80 WSP at recommended rates for a registered use.

CROP	RE-CROPPING INTERVALS (DAYS)
Cotton	No restrictions
Leafy vegetables (except Brassica vegetables) and Root and Tuber vegetables	365 days
All other crops	Prohibited

Note: When using INTREPID 80 WSP with other registered pesticides, always refer to rotational restrictions and precautions on the other product label(s) and comply with the most restrictive rotational guidelines.

6/8

**USE DIRECTIONS FOR POME FRUIT** (Apples, Crabapples, Loquat, Mayhaw, Pears including Oriental, Quince)

**Ground Application:** Make applications by conventional ground sprayers which are calibrated to deliver a minimum of 50 gallons per acre to trellised trees or trees 10 feet tall or less. For trees greater than 10 feet tall use a minimum of 100 gallons per acre.

**Aerial Application:** Make applications of INTREPID 80 WSP Agricultural Insecticide in a minimum of 20 gallons per acre. INTREPID 80 WSP can be applied by aerial applications when situations warrant. However this method should not be used if the size of the tree or density of the foliage prohibits thorough, uniform coverage of the entire tree canopy.

**Spray Adjuvants:** The use of Spreader-Sticker such as LATRON B-1956 or similar spreader-sticker is highly recommended to maximize coverage and retention of the spray material.

TARGET PESTS	APPLICATION RATE		APPLICATION TIMING	RESTRICTIONS
	Acres per Pouch *	Ounces Product Per Acre		
Codling moth	2/3 to 1-1/3	3 to 6 (0.15 to 0.3 lb. AI/Acre)	For each codling moth generation, apply at peak egg lay (150-250 Day Degrees (DD) following biofix) followed by a second application in 10-15 days following the first application (usually 450 to 550 DD). Additional applications at 10 to 18 day intervals may be required for heavy infestations or sustained moth flight.	Do not apply more than 20 ounces product per acre per season.  Allow at least 14 days to elapse between final application and harvest.  Aerial application is allowed only for the last two applications prior to harvest.
Lesser appleworm Oriental fruit moth			For other internal feeding Lepidoptera larvae, apply at initiation of egg hatch. A second application may be required 10-15 days following the first application to ensure complete coverage of rapidly expanding fruits or foliage.	
Obliquebanded leafroller			<b>Spring (overwintering) generation:</b> Make 1 to 2 applications during the pink to petal fall period depending on infestation level. <b>Summer generation:</b> Begin applications at peak egg lay through early egg hatch (200-400 DD, base 43F°, following biofix). Make a second application 10 to 18 days later (usually 650-850 DD). A third application 10 to 15 days after the second application may be required under high pressure or sustained moth flight.	
Pandemis leafroller			<b>Spring (overwintering) generation:</b> Make 1 to 2 applications during the pink to petal fall period depending on infestation level. <b>Summer generation:</b> Begin applications at early egg lay through early egg hatch (250-400 DD, base 41F°, following biofix). Make a second application 10 to 18 days later (usually 600-800 DD). Under heavy infestation a third application may be required 10-14 days after the second application.	
Fruittree leafroller Redbanded leafroller Variegated leafroller			For control of other leafrollers, begin applications at early egg hatch for each generation. Make the first application before webbing and sheltering begins. Make a second application in 10-18 days to ensure complete coverage of rapidly expanding fruits or foliage.	

7/8

Tufted apple bud moth	1 to 2	2 to 4 (0.1 to 0.2 lb. AI/Acre)	<p><b>First generation:</b> Make application at 600-900 DD, base 45F°, following biofix, depending on local conditions. A second application at 60-90% egg hatch may be required under heavy infestation levels.</p> <p><b>Second generation:</b> Make the first application at 2300-2500 DD (20-30% egg hatch). A second application approximately 14 days later may be required under high pressure or sustained moth flight or on late maturing varieties.</p>
Spotted tentiform leafminer Western tentiform leafminer			<p><b>First generation:</b> Apply at pink to petal fall.</p> <p><b>Second, third generation:</b> Begin applications at early egg hatch for each generation.</p>

*\*INTREPID 80 WSP Agricultural Insecticide is supplied in water-soluble pouches containing 4 ounces of 80% AI wettable powder.*

8/8

**USE DIRECTIONS FOR COTTON**

**Ground Application:** Make applications by conventional ground sprayers which are calibrated to deliver a minimum of 5 gallons per acre.

**Aerial Application:** Make applications in a minimum of 2 gallons per acre.

**Spray Adjuvants:** The use of LATRON CS-7 spreader-binder or similar spreader-binder following the manufacturer's labeling is recommended to maximize coverage and retention of the spray material.

TARGET PESTS	APPLICATION RATE		APPLICATION TIMING	RESTRICTIONS
	Acres Per Pouch * (LBS. AI/Acre)	Ounces Product Per Acre		
Cotton Bollworm Tobacco Budworm	0.5 to 0.8	5 to 8 (0.25 to 0.4 lb. AI/Acre)	For bollworms, budworms and other internal feeding larvae, initiate applications when larval infestations on the terminals reaches 2 to 4 %. Retreat when larval reinfestation reaches 4 % or 2 worms per 25 plants.	For resistance management purposes do not make applications of INTREPID 80 WSP to more than two consecutive generations of cotton bollworm or tobacco budworm in a single season.
Beet Armyworm Cabbage Looper Cotton Leafworm Cotton leaf perforator Fall Armyworm Saltmarsh caterpillar Southern Armyworm Soybean Looper True Armyworm Yellowstriped Armyworm	1.33 to 4.0	1 to 3 (0.05 to 0.15 lb. AI/Acre)	Apply at egg hatch or when first signs of feeding occur. Use the higher rates for heavier infestations and under conditions in which thorough coverage is more difficult. Under heavy infestations, continuous moth flights and/or egg masses and larvae in all stages of development, reapplication on a 10- to 14-day schedule will be required to protect new growth until moth flights and/or hits subside.	Do not apply more than 20 ounces. product per season.  Allow at least 14 days to elapse between final application and harvest.  See <b>ROTATIONAL CROP RESTRICTIONS</b> in the body of this label.

*\*INTREPID 80 WSP Agricultural Insecticide is supplied in water-soluble pouches containing 4 ounces of 80% AI wettable powder.*

**INTREPID, LATRON B-1956 & CS-7 ARE TRADEMARKS OF ROHM AND HAAS COMPANY**

**ROHM AND HAAS COMPANY, 100 INDEPENDENCE MALL WEST, PHILADELPHIA, PA. 19106, TELEPHONE 215-592-3000 5-0165**

8359-M2b  
JL  
3/26/01