



000707-00270-061699

Printed: 09:58:05 Wednesday, 21 Jul, 1999 # 19 / 3647

Systems Integration Group, Inc.



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:
707-270

Date of Issuance:
JUN 16 1999

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Unconditional

Name of Pesticide Product: MIMIC
2LV

Name and Address of Registrant (include ZIP Code):

Rohm and Haas Company
100 Independence Mall West
Philadelphia, PA 191062399

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5).

Make the following label changes before releasing the product for shipping:

- a. Revise the EPA Registration Number to read, "EPA Reg. No. 707-270".
- b. Add the following statements to the Environmental Hazards section of the label:

"This product is toxic to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Under some conditions, this chemica may also have a high potential for runoff into surface water for several weeks or months after application. Do not cultivate within 10 feet of aquatic areas so as to allow growth of a vegetative filter strip".

- c. A Spray Drift precaution, "Drift from applications of this pesticide is likely to result in damage to sensitive aquatic invertebrates in water bodies adjacent to the treatment area.", must be added to the label.

Signature of Approving Official:

See page 2

Date:

June 16, 1999

- d. Change the Ground Water Statement to read; "This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination."
- e. In the Statement Of Practical Treatment for If Inhaled add, "If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention."
- f. In the Precautionary labeling change the statement, " May cause eye irritation." to read "causes moderate eye irritation.". Also change "Harmful if absorbed through the skin, swallowed, inhaled" to read "Harmful if inhaled.". Add the statement, "Wear long-sleeved shirt and long pants, socks and shoes."

) A copy of the acute toxicity and product chemistry reviews are enclosed for your files.

Submit two (2) copies of the revised final printed label before you release the product for shipment.

If the final printed labeling is not received prior to shipment of this product, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these requirements.

A stamped copy of the label is enclosed for your records.

) Sincerely,

Joseph M. Johnson
)
 Marion J. Johnson, Jr.
 Product Manager 10
 Insecticide Branch
 Registration Division (7505C)
gjt

MO
12/21/98
Forestry

<p>MIMIC[®] 2LV</p> <p>AGRICULTURAL INSECTICIDE</p>	 <p>ROHM AND HAAS PHILADELPHIA, PA 19106</p>
--	---

ACTIVE INGREDIENT

Tebufenozide

Benzoic acid, 3,5-dimethyl-, 1-(1,1-dimethylethyl)-2-(4-ethylbenzoyl)hydrazide.....

24.0%*

INERT INGREDIENTS.....

76.0%

TOTAL

100.00%

*Equivalent to 2 lbs. active ingredient per gallon

EPA REG NO. 707-

EPA EST. NO. 39578-TX-01

NOTICE: Before using this product, read the entire Precautionary Statements, Notice of Warranty, Directions for Use, Use Restrictions and Storage and Disposal Instructions.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Consult a physician if irritation persists.

IF INHALED: Move victim to fresh air.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF SWALLOWED: Call a physician or poison control center. Drink one or two glasses of water. If person is unconscious, do not give anything by mouth and do not induce vomiting.

ACCEPTED
with COMMENTS
In EPA Letter Dated:
JUN 16 1999

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
707-270

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May cause eye irritation. Harmful if absorbed through the skin, swallowed, inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations
Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark, except under forest canopy when aerially applied to control forest pests. Do not contaminate water when disposing of equipment washwaters and rinsate. Do not apply when weather conditions favor drift or runoff from areas treated.

This pesticide demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where the water table is shallow may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Water-proof gloves
- Shoes plus socks

STORAGE AND DISPOSAL

STORAGE: Store in a cool dry well-ventilated area, but not below 32°F.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA regional office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, by incineration or if allowed by State and local authorities, by burning. If burned stay out of smoke.

STEPS TO BE TAKEN IF MATERIAL IS RELEASED OR SPILLED: Appropriate protective equipment must be worn when handling a spill of this material. Transfer spilled material to suitable containers for recovery or disposal. Contain spills immediately with inert materials (e.g. sand, earth). Transfer liquids and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. **Refer to Precautionary Statements.**

CONDITIONS OF SALE AND WARRANTY

Rohm and Haas warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. **ROHM AND HAAS MAKES NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE.** Handling, storage and use of the product by Buyer or User are beyond the control of Rohm and Haas and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to, weather or soil conditions, presence of other materials, disease, pest, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. **IN NO CASE WILL ROHM AND HAAS OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT.**

GENERAL INFORMATION

MIMIC 2LV Agricultural Insecticide mimics the action of the natural insect hormone 20-hydroxyecdysone, the physiological inducer of the molting and metamorphosis process in insects. MIMIC 2LV is highly active against most lepidopterous larvae while having practically no activity at typical use-rates against other orders of insects. The selectivity of MIMIC 2LV allows for the maintenance of the populations of beneficial and predatory insects which is a key element in integrated pest management programs. MIMIC 2LV controls lepidopterous larvae through a novel mode-of-action by the induction of a premature lethal molt which initiates within hours of ingestion of treated crop surfaces. Contact activity has also been observed in some insects. Actual death of the larvae will take several days to occur although feeding by the insects generally ceases within 24 hours of ingestion.

USE RATE DETERMINATION

Carefully read, understand and follow label use rates, recommendations and restrictions. Apply the amount specified in the following table with properly calibrated aerial or ground spray equipment.

The low rates may be used for light infestations of the target lepidopterous species and the higher rates for moderate to heavy infestations. MIMIC 2LV may be applied in either dilute or concentrate sprays so long as the application equipment is calibrated and adjusted to deliver thorough, uniform coverage. Use the specified amount of MIMIC 2LV per acre regardless of spray volume used. Prepare only the amount of spray solution required to treat the measured acreage.

MIXING AND COMPATIBILITY

Fill the spray tank one-third to one-half full of clean water and slowly pour MIMIC 2LV Agricultural Insecticide into the spray tank. Maintain agitation in the spray tank during mixing, loading and application. Triple-rinse empty container and add rinsate to spray tank.

MIMIC 2LV Agricultural Insecticide is believed to be compatible with most commonly used agricultural fungicides, insecticides, growth regulators, foliar fertilizers and spray adjuvants. If in doubt, mix proportional amounts of all spray ingredients in a test vessel. Shake the mixture vigorously and allow it to stand for fifteen minutes. Rapid precipitation of the ingredients and failure to re-suspend when shaken indicates that the mixture is incompatible and should not be applied.

APPLICATION TIMING

The activity of MIMIC 2LV Insecticide is expressed primarily through ingestion by the target larvae. Consequently, the timing of application is dependent on the feeding behavior of the target pest. For internal feeding larvae, application must be made prior to the time that surface feeding occurs. For foliar or surface feeding larvae, application made while active feeding is occurring will be effective.

Re-application may be required to protect new flushes of foliage or rapidly expanding fruit. The re-application interval will vary depending on how rapidly the crop is growing and the generation time of the target pest. While MIMIC 2LV Agricultural Insecticide is essentially equally effective against all instars, it is generally good practice to make applications to early instars to avoid the heavy damage that can be inflicted by later instar larvae.

For best results, begin applications when first signs of feeding damage or when threshold levels of moths, eggs or larvae occur. Consult the Cooperative Extension Service, or other qualified professional authorities to determine the appropriate threshold for application in your area.

APPLICATION INSTRUCTIONS

Because MIMIC 2LV Insecticide must be ingested by the larvae, application must be in a manner that assures uniform and thorough coverage. Higher water volume and increased spray pressure generally provide better coverage. Operating an air-blast sprayer at ground speeds greater than 2 mph and making applications in an alternate row middle pattern in tree crops and vines may result in less than satisfactory coverage and poor performance, particularly in conditions of high pest infestation levels, extremely large trees and/or dense foliage. Avoid application under conditions when uniform coverage cannot be assured or when excessive spray drift may occur. A minimum of six hours drying time is required between the completion of application and the onset of precipitation to ensure optimum performance.

CHEMIGATION

Do not apply this product through any type of irrigation system.

SPRAY ADJUVANTS

The addition of agricultural adjuvants is not required to maximize the performance, coverage or weatherability of MIMIC 2LV. The addition of spray adjuvants is not recommended.

RESISTANCE MANAGEMENT

Any insect population may contain individuals that are naturally resistant to a specific pesticide, therefore, the use of any one insecticide against many consecutive generations of a pest can result in the development of resistance problems. To prevent or delay the development of resistance, Rohm and Haas Company recommends rotation of MIMIC 2LV with insecticides of alternate modes of action and the utilization of Integrated Pest Management practices such as routine monitoring, the use of treatment thresholds to time applications and cultural and biological controls wherever possible. We further recommend that MIMIC 2LV not be used on more than three consecutive generations of a pest. Since the development of resistance cannot be predicted we suggest that you consult local or State Extension personnel or your local Rohm and Haas Company representative for resistance management guidance appropriate to your crop, locality and production practices.

USE DIRECTIONS FOR FORESTS, TREES AND SHRUBS

Because MIMIC 2LV Insecticide must be ingested in order to be effective, it is essential that coverage is thorough and uniform. Higher carrier volumes and higher use rates are recommended for very large trees or dense stands and for heavy target pest infestations.

GROUND APPLICATION: Hydraulic ground sprayers should be calibrated to deliver a minimum of 50 gallons per acre. For mist blowers or air blast sprayers, use a minimum of 10 gallons per acre.

AERIAL APPLICATION: Make applications of MIMIC 2LV Insecticide in a minimum of one gallon per acre. Higher carrier volumes are recommended when environmental conditions are less than ideal for aerial applications.

Target Pests	Application Rate Ounces per Acre (lbs. AI/Acre)	Recommendations	Restrictions
Browntail Moth (<i>Euproctis chryssorrhoea</i>) Elm spanworm (<i>Ennomos subsignaria</i>) Fall Cankerworm (<i>Alsophila pometaria</i>) Fall webworm (<i>Hyphantria cunea</i>) Forest tent caterpillar (<i>Malacosoma disstria</i>) Gypsy moth (<i>Lymantria dispar</i>)	2.0 to 8.0 (0.03 to 0.12 lb. AI/Acre)	Apply to early instar (1 st , 2 nd , or 3 rd) larvae. In general, foliage development should be a minimum of 20%.	Do not apply more than 16 Ounces per year. Uniform coverage of the foliage is essential to provide maximum protection from defoliation and reduction of egg mass deposition.
Pine tip moth (<i>Rhyacionia frustrana</i> , <i>R. neomexicana</i> , <i>R. buoliana</i> , <i>R. rigidana</i> , <i>R. subtropica</i>)	8.0 (0.12 lb. AI/Acre)	Apply to early instar (1 st to 2 nd) larvae after each new foliage flush, in general, at approximately 25% shoot expansion.	
Spruce budworm (<i>Choristoneura fumiferana</i> and <i>C. occidentalis</i>) Tussock moth (<i>Dasychira pinicola</i> , <i>Lophocampa maculata</i> , <i>Orgyia pseudotsugata</i> , <i>O. vetusta</i>)	4.0 to 8.0 (0.06 to 0.12 lb. AI/Acre)	Make applications to 4 th to 5 th instar larvae that are actively feeding on foliage or are feeding outside the candle caps.	

M0
12/21/98
JL/rdm

MIMIC IS A REGISTERED TRADEMARKS OF ROHM AND HAAS COMPANY