


Facilities and equipment in federally inspected meat and poultry plants must be washed with an effective cleaning compound followed by a potable water rinse.

BEST AVAILABLE COPY



MAY 1 1985

Mr. Keith L. Raddick
Atlas Manufacturing Company
P.O. Box 224968
2321 Benetrix Street
Dallas, TX 75264-0968

Dear Mr. Raddick:

Subject: Revised Labeling for TOX Industrial Roach & Fly Spray
EPA Registration No. 713-5

The revised label submitted with your application of March 20, 1985, is acceptable provided that you delete the signal word "Warning" in the statement of flammability and replace it with the signal word "Caution."

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Timothy A. Gardner
Product Manager (17)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

RD/IRB:JOB: 51497:R.King:RD: 85:21:reg:Kandrick: 898-1270:4/18/85:Del.4/24/85

BEST AVAILABLE COPY

CONCURRENCES							
SYMBOL	B-767						
SURNAME	RK						
DATE	4/30/85						

2012 #713-5 PM 17

BEST AVAILABLE COPY

TOX INDUSTRIAL ROACH & FLY SPRAY

ACTIVE INGREDIENTS 100%:

Aromatic Petroleum Distillate	99.34%
N Octyl Bicycloheptene Dicarboximide	0.35%
Technical Piperonyl Butoxide*	0.21%
Pyrethrins	0.10%
	<u>100.00 %</u>

*Consists of 0.17% (butylcarbityl) (6-propylpiperonyl) ether and 0.04% other related compounds.

CAUTION: KEEP OUT OF THE REACH OF CHILDREN.
(See Side Panel for Further Cautions, & Disposal)

FPA Reg. No. 713-5

EPA Est. No. 713-TX-1

CONTENTS: () GALLONS

DIRECTIONS FOR USE

ROACHES AND WATERBUGS: Apply by spraying heavily in all cracks, crevices in woodwork, underneath sinks, dark places behind pipes, and in all infested areas, contacting as many insects as possible. Repeat the treatment as often as necessary.

FLIES, MOSQUITOES AND GAAITS: Close room and windows, and discharge spray mist vigorously in all parts of the room, using about 2 or 3 ounces for each 1,000 cubic feet of space. Remove pets, birds and cover fish aquariums before spraying. Leave room closed for at least 10 to 20 minutes, ventilate before re-entering. Sweep up all fallen insects and decay. Repeat treatment when necessary.

FLEAS (INFESTING PREMISES): Remove and destroy animal's bedding and spray sleeping quarters thoroughly, forcing the spray into all cracks and crevices in the floor and about Repeat treatment as necessary.

CLOTHES MOTHS, CARPET BEETLES (Exposed Stage only): Spray rugs, carpets, and all infested articles thoroughly, giving particular attention to folds and seams. Interior surface of storage containers such as closets, chests, and trunks should be sprayed regularly. Treatment should be repeated every 30 days, unless the articles are stored in a moth-tight container.

Grassary weevil, rice weevil, confound flour beetle, red flour beetle, sawtooth grain beetle, rusty grain beetle, spider beetle, angoumois grain moth, Mediterranean flour beetle, cheese skipper cheese mites, store tick mites: Discharge spray vigorously to all infested surfaces, insect hiding places and runways, contacting as many insects as possible for best results, using spray at the rate of 1 gallon TOX INDUSTRIAL ROACH AND FLY SPRAY to 750 square feet of area, equivalent to pint of run-off. Repeat the treatment when necessary.

700

ATLAS MANUFACTURING CO
P O BOX 22898 DALLAS, TEXAS 75221
PHONE (214) 551-0611

CHEMICAL
CLEANING
COMPOSITIONS

