

FRONT PANEL

1 of 3

PEARSON'S
GREEN - DEVIL DUST
Containing
5% Malathion


For many insects attacking vegetables, field and ornamental
crops, orchards, cattle, hogs, pets and poultry.

CAUTION: Keep out of reach of children
(See back panel for additional cautions.)

ACTIVE INGREDIENTS:

Malathion*.....5%
Inert Ingredients.....95%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

Manufactured by
PEARSON & COMPANY, Mobile, Ala. U.S.A.
Net Weight 1 pound



Back Panel

PEARSON'S GREEN DEVIL DUST
containing 5% Malathion

NOTICE: Read entire label and use in accordance with applications and
cautions. Thorough, full coverage application should be made as
necessary.

DIRECTIONS FOR USE

VEGETABLES: Beans: Mexican Bean Beetle, Leafhopper, Spider Mites, Aphids,
apply 30 to 35 lbs. per acre or 1 lb. per 1350 sq.ft. Make 2 or more
applications as needed. Do not apply within 1 day of harvest. BROCCOLI,
BRUSSELS SPROUTS, CABBAGE, KALE, TURNIP, MUSTARD GREENS: Aphids, imported
Cabbage worms, Cabbage Looper, apply 30 lbs. per acre or 1 lb. per 1350
sq. ft. Do not apply within 3 days of harvest on Broccoli. Do not apply
within seven days of harvest on Brussels Sprouts, Cabbage, Kale, Turnip,
Mustard Greens. Apply as needed.

CAROTS, RADISHES, PARSNIP, SALSIFY, HORSE RADISH: Aphids, apply 30 to 35
lbs. per acre or 1 lb. per 1350 sq.ft. Do not apply within 7 days of harvest.

CAULIFLOWER: Aphids, apply 50 lbs per acre or 1 lb. per 875 sq.ft. Do not
apply within 7 days of harvest.

COLLARDS, DANDELION, WATERCRESS, SWISS CHARD, PARSLEY: Aphids, apply 30 to
35 lbs. per acre or 1 lb. per 1350 sq. ft. Do not apply within 21 days of
harvest.

CORN (Grain form & Forage). Sap Beetle, Corn Earworm, apply 40 lbs. per
acre or 1 lb. per 1100 sq. ft. For Corn Earworm, begin treatments when 10%
of ears show silk. Make 4 to 5 applications at 3 to 5 day intervals, apply
by ground on hand dusting equipment or apply to individual ears by using a
brush. Injury may occur in the Whorl stage. Do not apply within 5 days of
harvest or forage use.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites. (Leafhopper on melons)
apply 30 to 35 lbs. per acre, or 1 lb per 1350 sq. ft. Do not apply Malathion

to Cucurbits unless plants are dry. Do not apply within 3 days of harvest.

ENDIVE: Aphids, Mites, apply 30 to 40 lbs. per acre or 1 lb. per 1250 sq. ft. Do not apply within 7 days of harvest.

KOHLRABI: Aphids, apply 30 lbs. per acre or 1 lb. per 1450 sq. ft. Do Not apply within 7 days of harvest.

LETTUCE: Aphids, Mites, Cabbage Looper, Leafhoppers, apply 30 to 40 lbs. per acre or 1 lb. per 1250 sq. ft. Do not apply within 14 days of harvest for leaf or 7 days of harvest for head.

MUSHROOMS: Phorid and Sclerid Flies, apply 10 to 15 lbs. per acre or 2 lbs. per 60 ft. single house. Make thorough application as soon after picking as possible. Repeat applications as necessary, usually twice a week. Do not apply within 1 day of harvest.

ONIONS: Thrips, apply 30 to 40 lbs per acre or 1 lb. per 1250 sq. ft. Do not apply within 3 days of harvest.

PEAS: Pea Aphid, apply 25 lbs. per acre or 1 lb. per 1750 sq. ft. Do not apply within 3 days of harvest.

POTATOES: Aphids, Leafhopper, apply 25 to 30 lbs. per acre or 1 lb. per 1575 sq. ft. Do not apply within 3 days of harvest.

PUMPKINS: Aphids, Mites, Leafhoppers, apply 30 to 35 lbs. per acre or 1 lb. per 1350 sq. ft. Do not apply to cucurbits unless plants are dry. Do Not apply within 3 days of harvest.

SPINACH: Aphids, apply 30 to 35 lbs. per acre or 1 lb. per 1350 sq. ft. Do not apply within 7 days of harvest.

TOMATOES: Spider Mites, Aphids, Tomato Russet Mite, apply 35 to 40 lbs. per acre or 1 lb per 1140 sq. ft. Do not apply within 3 days of harvest.

FRUITS & NUTS: Apples; Woolly Apply Aphid, Green Apple Aphid, Rosy Apple Aphid, European Red Mite, Clover Mite-(Applications as needed), Williamette Mite, Two Spotted Mite, make at least two applications 10 to 12 days apart in summer months, apply 30 to 40 lbs. per acre or 1 lb. per 1250 sq. ft. Do not apply within 3 days of harvest.

DATES: Nitidulid Beetles, apply 35 to 40 lbs. per acre, apply thoroughly to each cluster. Do not apply within 7 days of harvest.

PEARS: Pear Psylla, Aphids, Woolly Aphids, apply 30 to 40 lbs. per acre as needed, or 1 lb. per 1250 sq. ft. Do not apply within 3 days of harvest.

PINEAPPLE: Mealybug, apply 100 lbs per acre or 1 lb. per 435 sq. ft. Do not apply within seven days of harvest.

BLUEBERRIES: Blueberry Maggot, apply 25 lbs. per acre or 1 lb per 1750 sq. ft. Do not apply within 1 day of harvest.

CRANBERRIES: Leafhoppers, Black-headed Fireworms, apply 50 lbs. by airplane, 30 lbs. by ground machine per acre or 1 lb. per 1450 sq. ft. Do not apply within 3 days of harvest.

GRAPES: Leafhopper, Spider Mites, apply 25 to 30 lbs. per acre or 1 lb per 1575 sq. ft. Make 2 or more applications as needed. Do not apply within 3 days of harvest.

STRAWBERRIES: Aphids, Spider Mites apply 40 lbs. per acre or 1 lb. per 1100 sq. ft. Do not apply within 3 days of harvest.

FIELD CROPS: Alfalfa: Aphids, Potato Leafhopper, Spider Mites, Alfalfa Weevil Larvae, Spittle Bug, Vetch Bruchid, apply 30 lbs per acre or 1 lb. per 1450 sq. ft. For Spotted Alfalfa Aphid apply 12 to 15 lbs. per acre or 1 lb. per 3375 sq. ft. Do not apply to Alfalfa in bloom. Make repeated applications as needed. Do not apply within 7 days of harvest or pasturing. Clover: Aphids, Spider Mite, Leafhopper, apply 25 lbs. per acre or 1 lb. per 1750 sq. ft. Do not apply to clover in bloom. Do not apply within 7 days of pasturing.

COTTON: Boll Weevil, apply 25 to 50 lbs. per acre. For Desert Spider Mite, Cotton Aphid, Leafhopper, White Flies, Brown Cotton Leafworm, Cotton Leaf Perforator, Cotton Leafworm, Thrips, Lygus Bug, apply 10 to 25 lbs. per acre.

TOBACCO: Aphids, apply 20 to 30 lbs. per acre or 1 lb. per 1750 sq.ft. Do not apply within 7 days of harvest.

ORNAMENTALS: Aphids, Spider Mites, Whitefly, Fourlined Leaf Bug, Thrips, Rose Leafhopper, Bagworms, apply 25 to 30 lbs. per acre or 1 lb. per 1600 sq.ft. Injury may occur on certain ferns and on some species of crassula and canaerti juniper.

FLEAS: On dogs and cats apply sufficient 4% Malathion dust to treat animal thoroughly. For control of fleas in kennels, pens, yards, lawns and under houses. Remove manure or debris. Dust area at rate of 1 to 2 lbs. per 1000 sq.ft., using hand duster. If necessary, repeat applications limited to ONCE weekly.

DIRECTIONS for Making 4% Malathion Dust: Mix 4 parts 5% Malathion Green Devil Dust with 1 part talc (Ordinary talcum powder).

POULTRY: Northern Fowl Mite, Poultry Lice, Chicken Red Mite, dust individual birds with 4% Malathion dust by means of a shaker can or rotary hand duster. Poultry premise treatments: Northern Fowl Mite, Chicken Red Mite, Poultry Lice, apply 1 lb. per 50 to 60 sq.ft. of litter and floor space and to nest, roosts and adjacent areas. Apply with a rotary hand duster, puff duster or by sprinkling from a can or other container. Repeat application as necessary.

BEEF CATTLE: Flies, apply 5% dust direct to coat of animal, rub in. May be repeated at 2 week intervals. Do not use on dairy cattle.

HOGS: (Swine): Flies, apply 5% dust to animal as needed.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid getting in eyes, on skin or on clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs. Keep away from children and domestic animals. Malathion, upon repeated or prolonged use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine (1/100 grain tablets) is antidotal.

NOTICE:

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material.

CAUTION: This product is poisonous to fish. Keep out of lakes, streams and ponds.

U.S.D.A. Reg. No. 728-36

Manufactured by
PEARSON & COMPANY
Mobile, Alabama, U.S.A.

NOTE: All Precautionary Labeling will be printed in accordance with Type Size Requirements upon acceptance of this label.