

728-37

PM-16

10/26/87

19 3



**GREEN DEVIL SPRAY
CONTAINS 50% MALATHION**

Controls Scale, Aphids, Red Spider Mites on Camellias, Azaleas and other flowers and shrubs.

ACTIVE INGREDIENTS:

Malathion*	50.0%
INERT INGREDIENTS:	50.0%
TOTAL	100.0%

*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate.
This product contains 4.35 pounds of active ingredient per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes and get medical attention.
IF ON SKIN: Immediately remove contaminated clothing, INCLUDING SHOES, and wash skin with soap and plenty of water. If irritation develops, send for a physician.

NOTE TO PHYSICIAN: Malathion is a cholinesterase inhibitor. Atropine is antidotal.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

EPA REG. NO. 728-37	MANUFACTURED BY:
EPA EST. NO. 19713-MS-1	SOUTHLAND PEARSON & COMPANY
	BOX 7344 * MOBILE, AL 36607

**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: HARMFUL IF SWALLOWED. Avoid breathing of spray. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Read entire label. Observe all cautions and warnings. Spray infested plants thoroughly; be sure to wet the underside of leaves. Repeat as often as necessary for good control.

ORNAMENTALS: Roses, Camelias, Azaleas, Stocks, Chrysanthemum, Evergreens, and many other flowers and shrubs: Aphids - 1 1/2 to 2 tsp./gal. of water. Red Spider Mites, Mealybugs, Woolly Aphids, White Flies, Thrips, Bagworms, Rose Leafhoppers, Japanese Beetle Adults - 1 tbsp./gal. of water. Black Scale, Soft Brown Scale, Oyster Shell Scale, and certain others - 1 1/2 to 2 tbsp./gal. of water (*See Caution below).

(*Do not use on Boston, Maidenhair, and Pteris ferns and some speices of Crassula).

VEGETABLES: Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Beans, Peas, Potatoes: Aphids - 1 1/2 to 2 ts./gal. of water. Red Spider Mites, Mealybugs, White Flies, Thrips, Bean Leafhopper, Tomato Leafhopper, Japanese Beetle Adults - 1 tbsp./gal. of water. Do not apply to beans within 1 day of harvest. Do not apply to broccoli and peas within 3 days of harvest, and to brussels sprouts, cabbage, cauliflower, or kale within 7 days of harvest.

FLEAS ON DOGS AND CATS: 2 tbsp./gal. of water. Wet animal thoroughly. Do not apply more than once a week. ANIMAL QUARTERS: 10 tbsp./gal. of water. Spray kennels, pens, yards, lawns and under houses at the rate of 1 gal...diluted... spray per 1000 sq. ft. Remove animals before treatment. Put in fresh bedding after treatment.

HOUSEHOLD PESTS: Roaches, Ants, Spiders, Mosquitoes, Silverfish, Crickets, Clover Mites, Earwigs: 8 tbsp./gal. of water. Using a coarse, wet spray or a paint brush apply to window sills and screens, baseboards, under sinks (especially where pipes come through walls and floors) and dark places where pests hide. Do not contaminate food or utensils.

Use only with adequate ventilation. After using this product indoors, ventilate thoroughly before occupying enclosed spaces. Do not allow contact with treated surfaces until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities.

WARRANTY AND CONDITION OF SALE

OUR RECOMMENDATIONS FOR USE of this product are based upon tests believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall Southland Pearson or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by Southland Pearson & Company and is accepted as such by the Buyer.

