

PEARSON'S
5%
Sevin Dust
BETTER
KILLER

controls
PICKLEWORM INJURY
MEXICAN BEAN BEETLE
TOMATO FRUIT WORM
COLORADO POTATO BEETLE

*and dozens of other
garden pests as
listed on back panel*

Manufactured by
PEARSON & COMPANY
Mobile, Ala.

PEARSON'S



5% Sevin Dust
BETTER
KILLER

THIS PACKAGE CONTAINS

ACTIVE INGREDIENT	
1-Naphthyl N-Methylcarbamate	5.00%
INERT INGREDIENTS	95.00%
TOTAL	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs, troughs and watering receptacles.

© Trademark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate
Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application. If possible avoid use when honeybees are visiting crop. If necessary to use during this period warn beekeepers in advance of application to move hives from vicinity for one week after application.

U.S.D.A. Reg. No. 728-58

NET WEIGHT 4 POUNDS

Manufactured by
PEARSON & COMPANY • MOBILE, ALABAMA

PEARSON'S
5%
Sevin Dust
BETTER
KILLER

controls
PICKLEWORM INJURY
MEXICAN BEAN BEETLE
TOMATO FRUIT WORM
COLORADO POTATO BEETLE

*and dozens of other
garden pests as
listed on back panel*

Manufactured by
PEARSON & COMPANY
Mobile, Ala.

ACCEPTED WITH COMMENTS

PEARSON'S



5% Sevin Dust BEETLE KILLER

THIS PACKAGE CONTAINS

ACTIVE INGREDIENT	
(1-Naphthyl N-Methylcarbamate).....	5.00%
INERT INGREDIENTS	95.00%
TOTAL	100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and foodstuffs, troughs and watering receptacles.

© Trademark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate

Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application. If possible avoid use when honeybees are visiting crop. If necessary to use during this period warn beekeepers in advance of application to move hives from vicinity for one week after application.

U.S.D.A. Reg. No. 728-58

NET WEIGHT 4 POUNDS

Manufactured by
PEARSON & COMPANY • MOBILE, ALABAMA

PEARSON'S 5% Sevin Dust BEETLE KILLER

controls

- PICKLEWORM INJURY
- MEXICAN BEAN BEETLE
- TOMATO FRUIT WORM
- COLORADO POTATO BEETLE

*and dozens of other
garden pests as
listed on back panel*

Manufactured by
PEARSON & COMPANY
Mobile, Ala.

begin applications when insect damage is first noticed. Repeat every 7 to 10 days or as necessary for good control.

VEGETABLES

BEANS: Apply 1 to 2 ounces per 50 feet of row to control Mexican Bean Beetle, Bean Leaf Beetle, Cucumber Beetle, Leafhopper, Flea Beetle, Japanese Beetle, Lima Bean Pod Borer, Lygus Bug, and Corn Earworm.

CUCUMBER and SUMMER SQUASH: Apply 1 ounce per 50 feet of row to control Pickleworm, Melonworm, Cucumber Beetle, Squash Bug, and Flea Beetle.

TOMATO, EGGPLANT, and PEPPER: Apply 1 to 2 ounces per 50 feet of row to control Tomato Fruitworm, Fall Armyworm, Flea Beetle, Leafhopper, Tomato Hornworm, European Corn Borer and Stinkbug.

POTATOES: Apply 1 to 2 ounces per 50 feet of row to control Flea Beetle, Colorado Potato Beetle, Leafhopper, and Lygus Bug.

On all vegetables listed above Pearson's 5% Sevin Dust may be applied up to the day of harvest.

ORNAMENTALS

Dust infested plants thoroughly when damage is first noticed. Repeat every 7 days or as often as necessary for good

POULTRY (Chickens, Partridges, Pheasants, Pigeons): Northern Fowl Mite Lice and as a supplement to premise treatments for Chicken Mite, Fleas—1 lb. per 100 birds. Apply thoroughly by means of shaker can or hand duster. Repeat in 4 weeks if necessary. Do not use on poultry or game birds within 7 days of slaughter. **IN OR AROUND POULTRY HOUSES:** Chicken Mite, Fleas, Bed Bugs—1 lb. per 40 sq. ft. of litter or floor space, and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nest litter. Avoid direct contamination of eggs. Do not use in premises within 7 days of slaughter.

DOGS, CATS: Brown Dog Ticks, Fleas—Dust liberally over the animal and rub thoroughly into the skin. Also apply to the feet between the toes and on the legs. Apply at weekly intervals if needed. Avoid treatment of kittens less than 4 weeks of age.

IN OR AROUND DOG AND CAT QUARTERS: Ticks, Fleas—Dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary.

Manufactured by
PEARSON & COMPANY, Mobile, Ala.

control of Apple Aphid, Bagworm, Birch Leaf Miner, Blister Beetle, Boxelder Bug, Boxwood Leaf Miner, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Japanese Beetle, Lace Bug, Leafhopper, Leaf Roller, Oak Leaf Miner, Orange Tortrix, Periodical Cicada, Psyllids, Roselug, Rose Aphid, Scale Insects in the Crawler Stage, Tent Caterpillars, Willow Leaf Beetle, and exposed Thrips.

For Scale control apply in Spring and early Summer when Scale Crawlers are present.

LAWNS

Apply 1 pound per 250 square feet (10x25) to control Fall Army Worm, Sod Webworm, and Leafhopper. For best results apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

NOTICE: Pearson & Company makes no warranty, expressed or implied, concerning this material except that it conforms to the chemical description on the label. Neither Pearson & Company nor the seller shall be responsible in any way for injury to person, damage to property, or other loss resulting from the handling, storing, or use of this material. The buyer assumes all risk and liability and accepts and uses this material on these conditions.

DIRECTIONS: Use 2 level teaspoons of Pearson's 50% Sevin in 1 gallon of water. For best results, apply with a spraying apparatus. To use mix designated amount of Pearson's 50% Sevin in a little water to make a thin paste, then add required amounts of water. Do not store diluted spray. Make new solution for each use.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS AND SHADE TREES: Japanese Beetles, Aphids, Bagworms, Elm Leaf Beetles, Certain Leafminers, Rose Slugs, Periodical Cicadas (17-year Locust), Scale Insects, Tent Caterpillars, Blister Beetles, Leaf Rollers and Exposed Thrips.

Begin spraying when insects or their damage appear. Repeat applications weekly or as necessary. Do not use in Boston Bay.

FRUIT TREES (APPLE, PEARS AND PEACHES): Apple Maggots, Plum Curculion, Apple Aphids, Leaf Roller, Peach tree Borers, Oriental Fruit Moths and Codling Moths -- Make one application between bud break and bloom, a second application at petal fall and then at 3 to 4 week intervals for 3 to 4 months as necessary. Apple Maggots -- For best control, apply at 10 day intervals. Rosy Apple Aphids -- Apply before leaves are curled. May be applied within 1 day of harvest.

VEGETABLES (TOMATOES, EGGPLANT, PEPPERS, CUCUMBERS AND SUMMER SQUASH): Pickle Worms, Fruitworms, Melon Worms, Horn Worms, Fall Armyworms and Corn Borers. Apply when insects first appear. Repeat applications at 7 to 10 day intervals. Do not apply within 1 day of harvest.

NET WEIGHT SIX OUNCES

LAWN INSECTS (CHINCH BUGS, SOD WEBWORMS, FALL ARMYWORMS, FLEAS AND ANTS): Mix 6 ounces in 15 gallons of water and cover 850 sq. ft. of established lawn. For best results, use a spraying apparatus. Mow lawn first, and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary.

NOTICE: Follow directions carefully. Time and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

CAUTION: HARMFUL IF SWALLOWED OR INHALED. DO NOT TAKE INTERNALLY. AVOID BREATHING DUST. AVOID CONTACT WITH SKIN. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING. CHANGE CONTAMINATED CLOTHING DAILY. AVOID CONTAMINATION OF FEED AND FOODSTUFFS, TROUGHS AND WATERING RECEPTACLES.

* Trademark of Union Carbide Corp. for 1-Naphthyl N-Methylcarbamate.

EPA REG. NO. 7289-4

PEARSON'S 50% SEVIN*

Wettable Powder — Garden Spray
For Flowers, Ornamentals,
Lawns, Vegetables
and Fruits

KILLS: Chinch Bugs, Japanese Beetles, Fleas and Ants. See back panel for more complete list.

CAUTION: Keep out of reach of Children

Pearson & Company, Mobile, Ala. 36607

ACCEPTED WITH COMMENTS

ACCEPTED

**CHINCH BUGS, SOD WEBWORMS,
ANTS, FLEAS AND ANTS):** Mix 6 ounces
per and cover 850 sq. ft. of established
ults, use a spraying apparatus. Mow
ply after rain or watering. Do not
two days after application. Repeat
as necessary.

Directions carefully. Timing and method
rather and crop conditions, mixtures
not specifically recommended, and
factors in the use of this product are
of the seller. Buyer assumes all risks
and handling of this material not in
with directions given herewith.

**DANGEROUS IF SWALLOWED OR INHALED.
TOXIC INTERNALLY. AVOID BREATHING
CONTACT WITH SKIN. WASH THOR-
OUGHLY WITH SOAP AND WATER AFTER HANDLING.
WASH CONTAMINATED CLOTHING DAILY. AVOID
USE OF FEED AND FOODSTUFFS.
WATERING RECEPTACLES.**

© 1964 Union Carbide Corp. for 1-Naphthyl

ce

P.A. Reg. No. 728-64

PEARSON'S

50% SEVIN*

**Wettable Powder — Garden Spray
For Flowers, Ornamentals,
Lawns, Vegetables
and Fruits**

**KILLS: Chinch Bugs, Japanese
Beetles, Fleas and Ants. See
back panel for more complete
list.**

**CAUTION: Keep out of reach of
Children**

(See back panel for additional cautions)

CONTENTS

ACTIVE INGREDIENT	BY WEIGHT
Carbaryl (1-Naphthyl N- Methylcarbamate)	50.00%
INERT INGREDIENTS	50.00%
Total	100.00%

Manufactured By

Pearson & Company • Mobile, Ala. 36607