

DIRECTIONS FOR USE

FOR ESTABLISHED PLANTS: For trees and large plants, Punch holes 6" to 8" deep, 12" apart in area covered by foliage of plant. Put 1/2 teaspoonful of Pearson's 10% Nema-Kill in each hole and cover with soil. Slowly sprinkle with water to seal if possible.

FOR ESTABLISHED SMALL PLANTS: Side dress by making trench on both sides of the row, 6" from plants and 4" to 6" deep. Apply Pearson's 10% Nema-Kill to each side. Cover with soil and, if possible, sprinkle with water to seal.

FOR TREATMENT OF SEED BED: Prepare bed in usual manner. Apply Pearson's 10% Nema-Kill at rate shown in Supplemental Direction Sheet. Sprinkle with water to seal the surface. If soil temperature is around 80°F., seeding or transplanting may be made 10 days after treatment, at 70°F. after 14 days, at 65°F. after 21 days. Pearson's 10% Nema-Kill is also effective at lower temperatures, but longer waiting is necessary before planting.

FOR LAWN AND TURF: Prepare lawn or turf by making incisions with garden fork or aerator 2" to 4" deep, not more than 1 foot apart. Apply evenly, 1/2—1 1/2 lbs. of Pearson's 10% Nema-Kill per 100 sq. ft. Lightly sprinkle lawn or turf with water so that the fumigant may be washed from plants onto the soil. After 24 hours, water lawn or turf heavily, but do not wash the fumigant away by flooding.

PREPLANTING APPLICATIONS: Apply evenly in rows 12" apart at a depth of 6" to 8", then seal the chisel channels immediately with pressure roller or flood with water.

AT TIME OF PLANTING APPLICATIONS: Prepare bed in usual manner, taking precautions to avoid contact of seed or plants with the fumigant.

CAUTION: Harmful if swallowed. Avoid breathing of vapor. Do not get in eyes or on skin. Wash thoroughly with soap and water after handling. Do not contaminate feed and foodstuffs. Keep out of reach of children, pets and irresponsible persons. Keep children and pets off treated areas until fumigant has been washed into soil and surface is dry. Do not reuse container. Destroy when empty by burning. Stay away from smoke or fumes.

Use only for plants listed; do not use more than recommended dosage and treat seed beds as far in advance of planting as possible.

Do not feed forage crops grown on soils treated with Nemagon to dairy animals or to livestock finished for slaughter.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from area treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

USE PEARSON'S 10% NEMA-KILL FOR CONTROL OF NEMATODES IN MINERAL SOILS FOR:

Established Plants	Seed, Transplants or Bulbs
Lawn and Turf grasses	Tomatoes
Peach trees	Carrots
Fig trees	Snap Beans
English Walnut trees	Summer Squash
Grape vines	Radishes
Hibiscus	Cotton
Rose	Croft Lily
Azalea	Grape cuttings
Camellia	
Gardenia	
Strawberry	

Application Rates and Hundreds of Additional Uses shown in SUPPLEMENTAL DIRECTION SHEET INSIDE.

For Early and Late Blight, Anthracnose, Septoria Leaf Spot, Tomato Fruit Worm, Tomato Horn Worm, Flea Beetle and Pin Worm, use Pearson's Tomato Dust.

U.S.D.A. Reg. No. 728-66

Manufactured by
PEARSON & COMPANY
MOBILE, ALABAMA

PEARSON'S

10%

Nema-Kill

Kills Nematodes

- GRANULES
- EFFECTIVE
- EASY TO USE

ACTIVE INGREDIENTS

1, 2-dibromo-3-chloropropane	9.70%
other Halogenated C3 Compounds	20%
INERT INGREDIENTS	90.10%

TOTAL 100.00%

CAUTION: Keep out of reach of Children
(See back panel for additional cautions.)

FARM • GARDEN • LAWN

NET WEIGHT 50 POUNDS

SUPPLEMENTAL DIRECTION SHEET FOR PEARSON'S 5% NEMA-KILL

*p — Planting Time
*PP — Post Planting

NOTE: When using Pearson's 10% Nema-Kill, apply at 1/2 the rates shown below.

CROP	lbs/acre	lbs/100 ft. row	DIRECTIONS
<u>Cotton</u>	520 173 (36" rows)	1 1/4 1 1/4	preplant, over-all application p, pp, row application
FRUIT & NUT CROPS			
<u>grapes, peaches, apricots, & nectarines</u>	1720	4	preplant, p, pp
<u>raspberries, black, dew, logan & boysen berries</u>	1720	4	preplant, p, pp allow at least 48 days between treatment and harvest
<u>strawberries</u>	525 270 (24" rows)	1 1/4 1 1/4	preplant overall application preplant, p, pp, row application allow 55 days between treatment & harvest
<u>strawberries</u>	525	1 1/4	preplanting, overall application, allow 55 days from treating to harvest
<u>figs</u>	1200	2 3/4	pp, allow at least 30 days between treatment and harvest
<u>ENGLISH WALNUTS</u>	690	1 1/2	pp, allow at least 30 days between treatment and harvest
VEGETABLES			
<u>snap beans, cucumbers, summer squash carrots, celery, parsnip, radishes turnips, cabbage, cauliflower, broccoli brussels sprouts, okra</u>	480	1	preplant, p, pp, row application do not feed plant parts of parsnips, carrots, turnips or vines of beans to livestock
	480	1	preplant, p, pp, over-all application
<u>lettuce, endive (escarole)</u>	350	3/4	preplant, over-all application
<u>melons</u>	520 108 (5' rows)	1 1/4 1 1/4	preplant, over-all application preplant, p, row application
<u>tomato, eggplant, pepper (except bell peppers) direct seeded</u>	525 171 (36" rows)	1 1/4 1 1/4	preplant, overall application preplant, p, pp, row application
<u>transplant</u>	260	2/3	preplant, over-all application

CROP	lbs/acre	lbs/100 ft. row	DIRECTIONS
ROSE new to 6 mo old established, 6-12 established over 12 mo	175	1/2 1 2	preplant, pp row application row application
flowering cherry, chokeberry, deutzia dogwood, gladiola, hydrangea, phila- delphus, snowberry, spirea, visburnum	350	3/4	preplant, p, pp
TURF & LAWN bent grass, bluegrass, bermuda grass, st. augustine	700-1300	1-3	preplant, pp
ORNAMENTAL & NURSERY PLANTS african violet, ageratum, aloe	350-1000	1-1 1/3	preplant, over-all application
aralia, baby tears, begonia cactus caladium croton, dieffenbachia, dracaena, godseffiana, haworthia, ivy, jadeplant, leuodendron, nautilogalyx, nephthytis, palm, neantha belle, panda ears, sarcocca, schefflera, spathiphyllum, tradescantia,	2 lbs per cubic yard potting soil	1-1 1/3	row application
achillea, agladnema, ajuga, anchusa arbor vitae, ardisia, asparagus fern, aster, astilbe, azalea, barberry, bleeding heart, (dicentra), boxwood, camellia, campanula, caryopteris, catananohe, chrysanthemum (hardy), coreopsis, shasta daisy, delphinium, dianthus, american elm, euonymous, fatsia, forsythia, funkia, gardenia, gypsophila, heliopsis, helleborus, hibiscus, hypericum, illex (holly), juniper, lavender, ligustrum (privet), lilac, croft lily, lily of the valley, lythrum, maple, maranta, monarda, monstera, oak, pachysandra, penstemon, peony, peperomia, peri- winkle (vinca), philodendron, philox, pine, pittosporum, poppy, pothos, privet, flowering quince, rose, rudbeckia, salvia, sedum, spruce, stokesia, synogonium, taxus, tritoma, trollius, veronica, vinca, violet, weigelia	860-1730		preplant, over-all application 1-2 1/2 per 100 sq ft row application, pp on established plants

