


Revised labelling for Pearson's Borer Spray & Stock Dip U.S.D.A. Reg. No. 728-70  
July 19, 1967

FRONT PANEL

PEARSON'S  
BORER SPRAY & STOCK DIP  
Contains 20% Lindane E.C., 1.65 lbs. Lindane per gallon  
KILLS

Pine Bark and Turpentine Beetles, Flat Headed Borers, Peach Tree Borers, Leaf Miners.  
Stock dip for Beef Cattle, Swine, Sheep & Goats. Spray for Dogs.

Active ingredients:  
Lindane-Gamma isomer of Benzene Hexachloride ..... 20%  
Xylene ..... 77%  
Inert ingredients ..... 3%  
100%



CAUTION: Keep out of reach of children.  
(See back panel for additional cautions)

Pearson & Company  
Mobile, Alabama 36601

Contents: 1 Pint

U.S.D.A. Reg. No. 728-70

BACK PANEL

DIRECTIONS

ORNAMENTALS

Aphids, Thrips, Leaf Miners, Cucumber Beetles: On Azaleas, Boxwood, Chrysanthemum, Larkspur, Rose and other flowers, both ornamental shrubs and trees, use one pint per 100 gallons of water, or 1 teaspoonful per 1 gallon of water.

Lacebug: On Azaleas, Pyracantha and Rhododendron, use 1½ pints per 100 gallons of water, or 1½ teaspoonfuls per 1 gallon of water. Apply in spring when immature insects first appear, and repeat as necessary.

Pine Bark Beetles, Turpentine Beetle: Use 1 pint per 5 gallons of water. Wet trunks thoroughly to runoff to height of 6 feet. Apply at first sign of attack, and repeat in 3 to 6 months if necessary.

Flat Headed Borers, Round Headed Borers, Peach Tree Borers: On Apples, Cherries, Peaches, use 3 pints per 100 gallons of water, or 1 tablespoonful per 1 gallon of water. Apply to trunk, crotch to ground. Soak bark to runoff and around base of tree. For best results the first trunk spray should be made between July 10 and 15, and two subsequent applications at 3 to 4 week intervals. DO NOT APPLY TO FRUIT OR FOLIAGE.

Dogwood Borer, Rhododendron Borer, Lilac Borer: On Dogwood, Rhododendron, Lilac, use 3 pints per 100 gallons of water, or 1 tablespoonful per 1 gallon of water. Make 2 to 3 applications 7 to 10 days apart in May and early June. Apply to trunk and to lower limbs, particularly to injured areas and old borer wounds.

LAWNS

Ticks, Chiggers, Fleas, Leafhoppers: In lawns, use 1 pint per 10,000 square feet, or 3 tablespoonfuls per 1,000 square feet in sufficient water for good coverage. Repeat application if necessary. CAUTION: Keep children and pets off treated areas until the insecticide has been washed into the soil and treated areas have dried completely.

BACK PANEL, DIRECTIONS CONTINUED

LIVESTOCK

CAUTION: Do not use on dairy animals. Do not use on emaciated, sick or lactating animals. Do not spray animals within 30 days or dip animals within 60 days of slaughter.

Thoroughly stir dipping vats before using. If any separation of the emulsion occurs, do not use the dip. Do not treat animals under three months of age. If young animals are to be dipped, individual dipping is recommended to avoid the possible swallowing of the dipping solution by the animals. Do not treat sow within 2 weeks of farrowing, nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather. Avoid treating overheated or sick animals.

Beef Cattle: Lice and Ticks. Use  $1\frac{1}{2}$  pints per 100 gallons of water. Thoroughly spray entire animal. Repeat treatment in 2 to 3 weeks if necessary. Sarcoptic Mange and Scab Mites: Spray using 2 pints per 100 gallons of water. Make only 2 applications, 7 days apart. Do not apply to calves under 6 months of age at this dosage.

Swine: Lice and Sarcoptic Mange Mites. Use 2 pints per 100 gallons of water and apply as a spray or dip. For Mange Mites, repeat treatment in 2 weeks if necessary.

Sheep and Goats: Ticks, Keds, Lice. Use 2 pints per 100 gallons of water as a spray, or use 1 pint per 100 gallons of water in a dip. Repeat treatment in 2 to 3 weeks if necessary.

Dogs: Fleas, Lice, Ticks and Sarcoptic Mange. Use 1 teaspoonful per 1 gallon of water and apply as a spray or wash by ruffing the coat to insure penetration of the solution down to the skin, paying particular attention to ticks. This application should be repeated as necessary. OR use  $1-\frac{1}{3}$  teaspoonfuls per 1 gallon of water and apply as a spray or wash after wetting the coat with water. Do not reapply this application more often than at weekly intervals.

RESIDUAL SPRAYS

Stable Fly, Mosquitoes, Roaches, Ants: Inside and outside of barns, hog pens, stables and other buildings, use  $1\frac{1}{2}$  to 2 pints per 10 gallons of water and apply at the rate of 1 gallon per 1000 square feet of surface. Thoroughly cover resting surfaces for flies and mosquitoes. Apply to cracks and crevices, and other hiding places for control of roaches and ants. Repeat applications as necessary. Do not apply in milk rooms or poultry houses. Do not contaminate animal food or water.

Ticks and Chiggers: Use 2 quarts per 100 gallons of water and spray weeds and underbrush harboring insects. Repeat applications as necessary. DO NOT use on food crops or grazing areas.

CAUTION: Keep out of reach of children. Harmful if swallowed. Avoid breathing spray mist. May be absorbed through skin. In case of contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets other than dogs. Do not use on humans. Do not use or store near heat or open flame. Do not apply to foliage of white or sweet potatoes. Destroy container when empty.

WARNING: Keep out of lakes, ponds and streams. Do not use for any purpose not specified on the label. This product is poisonous to fish and wildlife.

NOTICE: This material is sold without warranty as to hazards or results.