



ACCEPTED

BORER SPRAY and STOCK DIP

(Contains 20% LINDBAN® I.C.)
1.65 lbs. Lindane per gallon.

KILLS Pine Bark and Turpentine Beetles, Flat Headed Borers, Peach Tree Borers, Leaf Miners.

Stock Dip for Beef Cattle, Swine, Sheep, and Goats.

Active Ingredients:

KEEP OUT OF REACH OF CHILDREN

CAUTION

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center immediately.

If swallowed, call a physician or poison control center. Gastric lavage is indicated if material was taken internally. Do not induce vomiting.

GASTRIC TUMOR: IS INFLUENCE OF H. PYLORI ON THE CANCER RISK RELATED TO INDUCE VOMITING? NORBERTO M. MARCHETTI AND CARLOS R. PIZZOLI

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing, and wash affected area with soap and water.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

See [Sect. 1.4.1](#) for Prerequisites.

Southland Lures n & Company
Box 7344
Mobile, Alabama 36654

EIA Ref. 17-5-1
EIA Est. 17-5-11-1

Net Content

PRECAUTIONARY STATEMENT

Hazards to Humans and Domestic Animals.

CAUTION: May be harmful if swallowed. Avoid breathing spray mist. May be absorbed through skin. In case of contact, wash immediately with soap and water. Wash thoroughly after handling and before eating or smoking. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not use on nursing pups or on bitches nursing pups. Do not use or store near heat or open flame. Do not apply to foliage of white or sweet potatoes.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds and other wildlife feeding in treated areas may be killed. Do not apply directly to water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on treated area. Do not apply when bees are actively visiting the crop, over crop, or weeds blending in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARD

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ORNAMENTALS

APHIDS, THrips, LEAFMINERS, CUCUMBER BEETLES: On Azaleas, Boxwood, Chrysanthemum, Larkspur, Rose and other flowers, both ornamental shrubs and trees, use one pint per 10 gallons, or 1 teaspoonful to 1 gallon of water.

LACEWING: On Azalea, Pyracantha, and Rhododendron, use 1½ pints per 100 gallons, or 1½ teaspoonful to 1 gallon of water. Apply in spray when immature insects first appear, and repeat as necessary.

FINE BARK EELMITE, TURFINGE EELMITE: Use 1 pint in 10 gallons of water. Wet trunks thoroughly to run off to height of 6 feet. Apply at first signs of attack, and repeat if necessary if necessary.

FLAT-HEADED BORER, ROUND-HEADED BORER, FRACH TREE BORER: On Apples, Cherries, Peaches, use 3 pints per 100 gallons, or 1 tablespoonful to 1 gallon of water. Apply to trunk, crevices to ground. Soak bark to run-off and around base of tree. For best results the first trunk spray should be made between July 10 and 15, and two subsequent applications at 3 to 4 week intervals. DO NOT APPLY TO FRUIT OR FOLIAGE.

DOGWOOD BORER, RHODODENDRON BORER, LILAC BORER: On Dogwood, Rhododendron, Lilac, use 3 pints per 100 gallons, or 1 tablespoonful to 1 gallon of water. Make 2 to 3 applications 7 to 10 days apart in May and early June. Apply to trunk and to lower limbs, particularly to injured areas and old borer wounds.

LAWNS: TICKS, CHIGGERS, FLEAS, LEAFHOPPERS in lawns. Use 1 pint to 10,000 square feet, or 3 tablespoonful to 1,000 square feet in sufficient water for good coverage. Repeat application if necessary. Caution: Keep children and pets off treated areas until the insecticide has been washed into the soil and treated areas have dried completely.

LIVESTOCK

CAUTION: Do not use on dairy animals. Do not use on emaciated, sick or lactating animals. Do not spray animals within 30 days or dip animals within 60 days of slaughter. Thoroughly stir dipping vats before using. If any separation of the emulsion occurs, do not use the dip. Do not treat animals under 3 months of age. If young animals are to be dipped individual dipping is recommended to avoid the possible swallowing of the dipping solution by the animals. Do not treat sow within 2 weeks of farrowing nor for at least 3 weeks thereafter. Avoid treating animals in cold stormy weather. Avoid treating over-heated or sick animals.

BEEF CATTLE: Lice and Ticks. Use 1½ pints per 100 gallons of water. Thoroughly spray entire animal. Repeat treatment in 2 to 3 weeks if necessary. **SARCOPTIC MANGE AND SCAB MITE:** Spray using 2 pints per 100 gallons of water. Make only 2 applications, 7 days apart. Do not apply to calves under 6 months of age at this dosage.

SWINE: Lice and Sarcoptic Mange Mites. Use 2 pints per 100 gallons of water and apply as a spray or dip. For mange mites, repeat treatment in 2 weeks if necessary.

SHEEP AND GOATS: Ticks, Keds, Lice. Use 2 pints per 100 gallons of water as a spray, or use 1 pint per 1½ gallons in a dip. Repeat treatment in 2 to 3 weeks if necessary.

DOGS: Fleas, Lice, Ticks, and Sarcoptic Mange. Use 1 teaspoonful per 1 gallon of water and apply as a spray or wash by rinsing the coat to insure penetration of the solution down to the skin, pay particular attention to ticks. This application should be repeated as necessary. Use 1 ½ teaspoonful per 1 gallon of water and apply as a spray or wash after wetting the coat with water. Do not apply this application more often than at weekly intervals.

RESIDUAL SPRAYS

STABLE FLY, HORSES, ROACHES, AND ANTS: Inside and outside of barns, hog pens, stables and other buildings, use 1½ to 2 pints in 10 gallons of water and apply at the rate of 1 gallon to 1000 square feet of surface. Thoroughly cover resting surfaces for flies and mosquitoes. Apply to cracks, crevices, and other hiding places for control of roaches and ants. Repeat applications as necessary. Do not apply in milking or poultry houses. Do not contaminate animal food or water.

TICKS AND CHIGGERS: Use 2 quarts per 100 gallons of water and spray woods and underbrush harboring pests. Repeat applications as necessary. Do NOT use on food crops or grazing areas.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Do not store near heat or open flame.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved State procedures under Subtitle C of the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then dispose of in a sanitary landfill, or by other approved State and local procedures.

WARRANTY - CONDITION OF SALE:

DIRECTIONS FOR USE of this product are based on field use and tests believed reliable and should be followed carefully. It is however impossible to eliminate all risks associated with use of this product. Because such factors as weather conditions, foreign material and manner of use for application are all beyond the control of the Seller of this product, such things as crop injury, ineffectiveness or other unintended consequences may result.

ALL SUCH RISKS ARE ASSUMED BY THE BUYER

Seller warrants that the product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the direction for use. Seller makes no other warranties, express or implied, including FITNESS OR MERCHANTABILITY. In no case shall the Seller be liable for consequential, special or indirect damage resulting from the use of the "fix" of this product. The statement is a condition of sale by Seller and is accepted as such by the Buyer.