

ACCEPTED  
 May 29-1969  
 UNDER THE  
 FEDERAL  
 PURCHASE  
 REGULATIONS  
 728-81

**ORNAMENTALS — Azaleas and Camellias** (Azalea petal blight and Camellia flower blight): apply as a drench under plants at a rate of 1½ quarts in 30 gallons of water per 150 square feet. Begin application prior to opening of buds and repeat every 3 to 4 weeks through the bloom period.

**NOTICE TO BUYER:** Seller makes no warranty, express or implied, by operation of law or otherwise, except that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. This warranty shall not extend to the use of this product contrary to label instructions, or under abnormal conditions, or in combination with other materials, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any injury or damage resulting from such use. Seller shall not be responsible for special, consequential or contingent damages arising from a breach of this warranty.

Manufactured by  
**PEARSON & COMPANY**  
 MOBILE, ALABAMA 36601  
 U.S.D.A. Reg. No. 728-81

# PEARSON'S

## GREEN LAWN FUNGICIDE

Contains  
**Terrachlor\***

**For control of BROWN PATCH of certain lawns and certain soil borne diseases of ornamentals.**

Active Ingredient:	Pentachloronitrobenzene .....	24%
Inert Ingredients:	.....	76%
		<u>100%</u>

Contains 2 lbs. of Pentachloronitrobenzene per U.S. gallon at 68°F.

\* ® Terrachlor is an OMCC Trademark.

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 Harmful if swallowed.

Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using. Do not use or store near heat or open flame. Use adequate ventilation. Keep container closed. Avoid contamination with feed or foodstuffs.

## PEARSON'S GREEN LAWN FUNGICIDE

**DIRECTIONS FOR USE:** For control of Brown Patch (*Rhizoctonia solani*) of Southern or warm season lawns such as St. Augustine, Bermuda and similar grasses: Make one application in the Fall or Spring at first signs of infection. Mix one quart in sufficient water (10 to 15 gallons) to cover 1,000 square feet. Apply with sprinkler can, hand sprayer, or any other convenient method of obtaining even coverage. Then lightly water the treated area to insure movement of the chemical through the grass to the soil level. Retreat the area 3-4 weeks later if it has been subjected to unusually heavy rain-fall, or is flooded, or if the disease is severe or re-appears. Under certain growing conditions, a temporary discoloration of the grass may appear occasionally. This causes no harm and disappears shortly.

**NET CONTENTS 5 GALLONS**