

PEARSON'S ROSE DUST

DIRECTIONS FOR USE:

For best results start treatment early and use at weekly intervals throughout growing season. More frequent applications should be made during rainy periods.

FOR USE AS A DUST:

Punch holes in lid of can and sift directly on plants; OR BETTER - apply with dust gun. Dust lightly but thoroughly cover foliage.

FOR USE AS A SPRAY:

Use 1 pound of Pearson's ROSE DUST to 12 gallons of water. To make 1 gallon of spray use 5 level tablespoonsful of dust. Mix well, then spray. Shake sprayer occasionally to keep powder in suspension. Be sure to wet under side of leaves.

CAUTION:

Harmful if swallowed. Avoid breathing dust or spray mist. May be absorbed through skin; wash after handling. Avoid prolonged or repeated contact. Keep away from all contact with food or feed.

PROPER USE OF PEARSON'S ROSE DUST USUALLY EFFECTIVELY CONTROLS THE FOLLOWING:

Diseases:

Roses: Black Spot, powdery mildew, rust. Snapdragon rust, Carnation leaf spot.

Insect Pests:

Aphids, canker worm, cucumber beetle, flea beetle, flea hopper, Japanese beetle, rose chafer, thrips (certain species), Red Spider. For Hard-To-Kill garden insect pests use Red Devil Dust (Containing 20% specially prepared Sabadilla Seed). Controls Squash Bugs, Stink Bugs, and Harlequin Bugs.

U.S.D.A. Reg. No. 728-86

Manufactured by

PEARSON & COMPANY

MOBILE, ALABAMA 36601



MAY ALSO BE USED AS A ROSE SPRAY INSECTICIDE - FUNGICIDE

For Control of Many Plant Diseases and Insect Pests

Active ingredients:

Sulphur (98% through 325 mesh).....	20.0%
Copper, as metallic (derived from basic copper sulphate).....	3.7%
Zineb (zinc ethylene bisdithiocarbamate).....	3.5%
Carbaryl (1-Naphthyl N-Methylcarbamate).....	5.0%
Lindane (gamma isomer of benzene hexachloride).....	1.0%
Inert ingredients:	66.8%
Total:	100.0%

*(metallic zinc equivalent .829%)

CAUTION: Keep Out of Reach of Children.
(SEE BACK PANEL FOR ADDITIONAL CAUTIONS)

NET WEIGHT 1 POUND

ACCEPTED

June 17-1970

728-86