

CHILD-RESISTANT PACKAGE FOR BOMBOWER USES



LINDANE 20% BORER SPRAY

For Home & Garden Insect Control

KILLS Pine Bark and Turpentine Beetles, Flat Headed Borers, Peach Tree Borers, Leaf Miners.

ACTIVE INGREDIENTS:

Lindane (Gamma Isomer of Benzene Hexachloride).....	20.0%
INERT INGREDIENTS:.....	80.0%
TOTAL.....	100.0%

1.65 lbs. Lindane per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove victim to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected area with soap and water.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: This product contains a light hydrocarbon solvent and an aspiration hazard may exist. Gastric lavage may be indicated if material was taken internally.

See Side Panel for Precautionary Statements.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

EPA REG. NO. 728-123
EPA EST. NO. 19713-MS-1

NET CONTENTS:

Manufactured for:
Southland Pearson & Company

P. O. Box 7344

Mobile, Alabama 36607

(205) 456-1400

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. May be absorbed through skin. Wash thoroughly after handling and before eating, drinking, or smoking. Avoid contamination of feed and foodstuffs. Kitchen utensils such as teaspoons or tablespoons must not be used for food purposes after use in measuring pesticides. Do not use on humans. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, birds, and other wildlife. Do not apply directly to water or wetlands. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

This pesticide is highly toxic to bees exposed to direct application. Applications should be timed to coincide with periods of minimum bee activity, usually between late evening and early morning.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Avoid exposure to children. Do not allow children to apply or handle this product. Do not allow children or pets in treated area until surfaces are dry.

NOTE: For treatment of ornamentals and lawns the applicators must wear the following protective clothing during the application process: longsleeved shirt, long pants, waterproof gloves, full foot covering (such as shoes or boots) and a head covering (such as a hat).

ORNAMENTALS

APHIDS, THRIPS, LEAFMINERS, CUCUMBER BEETLES: On Azaleas, Boxwood, Chrysanthemum, Larkspur, Rose and other flowers, both ornamental shrubs and trees, use 1 teaspoonful to 1 gallon of water.

LACEBUG: On Azalea, Pyracantha, and Rhododendron, use 1½ teaspoonful to 1 gallon of water. Apply in spring when immature insects first appear, and repeat as necessary.

PINE BARK BEETLES, TURPENTINE BEETLE: Use 6 tablespoonfuls in 1 gallon of water. Wet trunks thoroughly to runoff to height of 6 feet. Apply at first sign of attack, and repeat in 3 to 6 months if necessary.

BEST AVAILABLE COPY

7 3
FLAT-HEADED BORERS, ROUND-HEADED BORERS, PEACH TREE BORER: On Apples, Cherries, Peaches, use 1 tablespoonful to 1 gallon of water. Apply to trunk, crotch to ground. Soak bark to runoff and around base of tree. **DO NOT APPLY TO FRUIT OR FOILAGE.** Pre-harvest intervals for apples and peaches is a 60 day.

DOGWOOD BORER, RHODODENDRON BORER, LILAC BORER: On Dogwood, Rhododendron, Lilac, use 1 tablespoonful to 1 gallon of water. Make 2 to 3 applications 7 to 10 days apart in May and early June. Apply to trunk and to lower limbs, particularly to injured areas and old borer wounds.

LAWNS

TICKS, CHIGGERS, FLEAS, LEAFHOPPERS in lawns. Use 3 tablespoonfuls to 1,000 square feet in 3 gallons of water for good coverage. Repeat application if necessary. Caution: Keep children and pets off treated areas until the insecticide has been washed into the soil and treated areas have dried completely.

Do not apply to food crops.

STORAGE AND DISPOSAL

Storage: Store in original container, preferably in a locked storage area, away from food or feed.

Disposal: Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

WARRANTY - CONDITION OF SALE:

OUR RECOMMENDATIONS FOR USE of this product are based upon test believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

In no case shall the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

BEST AVAILABLE COPY