

11474
 ONE-BOTTLE
 PRE-MEASURED TREATMENTS

Sudbury SEVIN

LIQUID CONCENTRATE
 INSECT SPRAY



ACTIVE INGREDIENTS: Cyfluthrin (100% active ingredient) 0.5%
 INERT INGREDIENTS 99.5%
 TOTAL 100%

KEEP OUT OF REACH OF CHILDREN
 CAUTION
 SEE BACK FOR ADDITIONAL PRECAUTIONS

USE ONE
 BOTTLE
 EACH TIME
 YOU SPRAY.

EACH
 ONE-BOTTLE
 DILUTES
 TO MAKE
 ENOUGH
 SPRAY
 FOR ONE
 AVERAGE
 APPLICATION.

BUY
 AND MIX
 EXACTLY
 WHAT YOU
 NEED.

NO
 MEASURING.
 ADDITIVE
 AND EASY
 TO USE.

CONVENIENT
 NO STORAGE
 OF PARTIALLY
 USED
 BOTTLES

NOCE
 with Cyfluthrin
 In EPA Letter Dated
 OCT 28 1982
 Under the Federal Insecticide,
 Fungicide and Rodenticide Act
 as amended, for the 25th year
 registered under EPA 10-1-1

BEST DOCUMENT AVAILABLE

Refer to back for complete use instructions. Remove bottles as needed from stock.
 DO NOT DESTROY THIS CARD UNTIL ALL UNITS HAVE BEEN USED

#48142
Net Contents: 8 bottles,
0.8 fl. oz. each

EPA REG. NO. 731-31
EPA EST. NO. 731-MA

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS AND DOMESTIC ANIMALS.**

CAUTION:

Irritant if swallowed. Do not breathe spray or dust. Avoid contact with skin. Wear gloves and long-sleeved work clothing. Wash hands thoroughly after use.

STATEMENT OF PRACTICAL TREATMENT

If swallowed — Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by sticking the back of the tongue with finger. Do not induce vomiting if you have swallowed less than 1 ounce.

If inhaled — Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin — Remove contaminated clothing and wash affected areas with soap and water.

If in eyes — Flush eyes with plenty of water. Call a physician immediately.

First Aid Note — Call a physician if severe effects develop. Atropine is antidote. Do not use 2-PAM, epinephrine or other respiratory stimulants.

ENVIRONMENTAL HAZARDS

This product is highly toxic to fish. Do not apply directly to water. Do not apply within 100 feet of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BEE CAUTION: MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Before applying, consult beekeepers to locate hives to avoid bee flight range one week after application or to take other equally effective precautions.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Protect from freezing. Do not reuse empty container bottle. Wrap container and put in trash collection.

Use any style clean sprayer. Shake Use-Berries according to rates below. Where a dosage of 1 to 2 bottles is indicated, use 1 bottle on young plants and 2 on mature plants. Apply when insects or other damage first appear. Repeat as necessary. For best results, apply full coverage to all exposed parts of plants, especially tops and under leaf surfaces. Do not store diluted spray. Unless otherwise specified, application can be made on day of harvest.

VEGETABLE CROPS

Asparagus

Asparagus Beetle on seedlings or spears. Use one or two bottles per 1 quart (32 oz.).

Application: Do not apply more often than once every 3 days. Do not apply within 1 day of harvest.

Beans (Green) (Vine Crop)

Mexican Bean Beetle. Use 1 bottle per 1/2 gallon (64 oz.) Bean Beetle, Spider Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Vein Beet, Caterpillars, Western Bean Cutworm. Use 1 bottle per 1 quart (32 oz.).

Armyworms, Colorado Corn Earworm, Stink Bugs, Tarnished Plant Bugs. Use 1 bottle per 1 quart (32 oz.).

Cauli (Chinese Cabbage, Bok Choy, Broccoli, Cauliflower, Chinese Broccoli, Kohlrabi, Kale, Lettuce, Mustard Greens, Peas, Snap Beans, Radishes, Rutabaga, Swiss Chard, Turnips, Spinach)

Flea Beetles, Harlequin Bug, Leafhoppers. Use 1 or 2 bottles per 1/2 gallon (64 oz.).

Spotted Leafhopper (Asian Yellow), Flea Beetles. Use 1 bottle per quart (32 oz.).

Armyworms, Imported Cabbageworm, Diamond Back Moth Larvae and Pupae, Corn Earworm, Tarnished Plant Bug, Stink Bugs. Use 1 or 2 bottles per 1 quart (32 oz.).

Application: For optimum control of Corn Earworm, when larvae heads begin to form, use 1 bottle per 1 quart (32 oz.), at 5 to 7 day intervals. To avoid possible injury to leaves, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Do not apply within 3 days of harvest of heads. Lettuce, Cauli, Beans (broccoli), Kohlrabi, Radishes, Rutabaga, Swiss Chard, and Turnips (broccoli). Do not apply within 14 days of harvest of Chinese Cabbage, Cauli, Broccoli, Endive, Escarole, Garden Beets (broccoli), Kale, Leaf Lettuce, Mustard Greens, Parsley, Savoy, Spinach, Swiss Chard and Turnips (broccoli).

Corn Crops (Sweet) — Broccoli Sprouts, Cabbage, Cauliflower, Potatoes

Flea Beetles, Harlequin Bug. Use 1 or 2 bottles in 1/2 gallon (64 oz.).

Armyworms, Imported Cabbageworm, Diamond Back Moth Larvae and Pupae, Corn Earworm. Use 1 or 2 bottles in 1 quart (32 oz.). Do not apply when 3 days of harvest.

Corn (Sweet)

Corn Earworm, Corn Rootworm, Aphids, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhoppers, Sap Beetles. Use 1 or 2 bottles per 1 quart (32 oz.).

Application: For control of corn rootworm damage, and to control fleas, apply to entire plant, repeat as necessary. For control of European corn borer and fall armyworm, apply at 2 to 3 day intervals starting when first signs begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

Curculionids — Use 1 bottle in 16 oz.

Application: Application to the base, region of the corn plant near the soil surface will seriously reduce flea populations.

Cucumbers, Melons, Pumpkins, Squash

Melonworm, Flea Beetles. Use 1 bottle in 1/2 gallon (64 oz.).

Cucumber Beetles, Flea Beetles, Leafhoppers, Squash Bugs. Use 1 bottle in 1 quart (32 oz.).

Application: Apply immediate application. To avoid possible injury to tender tissue, do not apply when humidity is expected during the next 2 days.

Garden Peas

Colorado Potato Beetle, Leafhoppers. Use 1 bottle in 1 quart (32 oz.).

Peas, Potatoes, Eggplants, Peppers

Colorado Potato Beetle, Flea Beetles, Leafhoppers. Use 1 or 2 bottles in 1/2 gallon (64 oz.).

European Corn Borer, Fall Armyworm, Lace Bugs, Stink Bugs, Tarnished Plant Bugs, Tomato Pinworm, Tomato Hornworm. Use 1 or 2 bottles in 1 quart (32 oz.).

Curculionids (on Potato, Tomato, Eggplant). Use 1 bottle in 16 oz.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS, AND SHADE TREES

Rose, Camellia, Gardenia, Zinnia, Chrysanthemum, Lilac, Azalea, Forsythia, Hydrangea, Azalea, Elm, Maple, Oak, Dogwood, Birch and Pine (do not use on Red-tailed).

Application: Apply 1 bottle per 1 quart (32 oz.) of water when insects or other damage appear. Repeat weekly or as needed to control Apple Aphid, Bagworm, Birch Leaf Miner, Bristle Beetle, Box Elder Bug, Browned Leaf Miner, Elm Leaf Apterous, Elm Leaf Beetle, Elm Bark Beetle, Elm Bark Beetle, Gypsy Moth, Japanese Beetle, June Beetles, Lace Bugs, Leafhoppers, Leaf Rollers, Mealy Bugs, Minute Weevil, Oak Leaf Miner, Orange Tortrix, Periodical Cicada, Plant Bugs, Red Tailed Saw, Pear Saw, Rose Apterous, Rosehug, Scale insects, Tent Caterpillars, Thorn Bugs, Thrips (various), White Leaf Beetles.

LIMITED WARRANTY: Sudbury and its subsidiaries warrant the storage and handling of this material, not in strict accordance with directions.

Sudbury. SUDBURY LABORATORY, INC., 6 OCTOBER HILL ROAD, HOLLISTON, MA 01746

BEST DOCUMENT AVAILABLE



