



50%

# METHOXYCHLOR

(50% Methoxychlor Wettable Powder)

## ACTIVE INGREDIENTS:

|                           |             |
|---------------------------|-------------|
| * Methoxychlor, Technical | 50%         |
| INERT INGREDIENTS         | 50%         |
| <b>Total</b>              | <b>100%</b> |

\* Equivalent to 44% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 6% of related compounds.

**Directions for use:** For control of fruit flies and other insects, on citrus, grapes, peaches, pears, apples, and other fruit. For control of insects on ornamentals, apply at the rate of 100 to 150 gallons per acre. Apply on a thorough cover spray to upper and lower surfaces of leaves, on blossoms for insecticide. Start treatment at first sign of insects. Repeat at 7 to 10 day intervals as required.

Do not apply on tomatoes, peppers, cucumbers, and melons within 7 days of harvest, nor on beans and cabbage within 3 days of harvest.

### M.F.A. OIL COMPANY

Columbia, Missouri  
USDA Reg. No. 746.4

**CAUTION:** Remove spray deposits from plant products consumed as food.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Sep 11, 1970

704-4

MANUFACTURED FOR  
**M.F.A. OIL COMPANY**  
Columbia, Missouri  
Net Contents 1 Pound

## DIRECTIONS FOR USE

For control of fruit flies and other insects, on citrus, grapes, peaches, pears, apples, and other fruit. For control of insects on ornamentals, apply at the rate of 100 to 150 gallons per acre. Apply on a thorough cover spray to upper and lower surfaces of leaves, on blossoms for insecticide. Start treatment at first sign of insects. Repeat at 7 to 10 day intervals as required.

Do not apply on tomatoes, peppers, cucumbers, and melons within 7 days of harvest, nor on beans and cabbage within 3 days of harvest.

**CAUTION:** Remove spray deposits from plant products consumed as food.

**ACTIVE INGREDIENTS:**  
Methoxychlor, Technical

**INERT INGREDIENTS:**

**Total**

Equivalent to 44% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 6% of related compounds.





(50% Methoxychlor Wettable Powder)

**ACTIVE INGREDIENTS:**

|                                |      |
|--------------------------------|------|
| *Methoxychlor, Technical ..... | 50%  |
| INERT INGREDIENTS .....        | 50%  |
| Total .....                    | 100% |

\*Equivalent to 44% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 6% of related compounds.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**

SEE OTHER CAUTIONS ON SIDE PANEL.

**DIRECTIONS FOR USE:**

**DAIRY CATTLE:** For Horn Flies: Directly dust cow, using 1 tablespoonful per animal. Dust on poll, neck, back and upper sides. Rub in lightly to insure that the material is worked down beneath the hair. A single application controls horn flies for about 3 weeks.

**NOTE:** Use only the DIRECT DUSTING method of application on dairy animals.

**CATTLE LICE, HOG LICE AND SHEEP TICKS:** Direct spraying of livestock: Use 1 lb. per 10-12 gallons of water. Spray to insure thorough coverage of animals, using 2 quarts per hog and 4 quarts per beef animal. On sheep use 2 quarts, applying to give complete coverage of the entire animal. Repeat in 14-16 days if infestations warrant. Do not spray dairy cattle.

**CAUTION: KEEP OUT OF REACH OF CHILDREN.**  
SEE OTHER CAUTIONS ON SIDE PANEL.

(50% Methoxychlor Wettable Powder)

**ACTIVE INGREDIENTS:**

|                                |      |
|--------------------------------|------|
| *Methoxychlor, Technical ..... | 50%  |
| INERT INGREDIENTS .....        | 50%  |
| Total .....                    | 100% |

**DIRECTIONS FOR VEGETABLE USE:**

For control of flea beetles and blister beetles on potatoes, tomatoes, peppers, cucumbers, and melons, and for control of cabbage worms on cabbage, use 2 pounds 50% methoxychlor per 100 gallons of water or four teaspoons per gallon of water. Apply mixture at the rate of 100 gallons per acre. For control of Mexican bean beetles and tomato fruit worms on beans and tomatoes, apply above mixture at the rate of 100 to 150 gallons per acre. Apply as a thorough cover spray to upper and lower surfaces of leaves, on tomatoes for tomato fruit worm control apply directly to the fruit. Start treatment at first sign of insects. Repeat at 7 to 10 day intervals as required.

Do not apply on tomatoes, peppers, cucumbers and melons within 7 days of harvest, nor on beans and cabbage within 3 days of harvest.

MANUFACTURED FOR  
**MFA OIL COMPANY**  
Columbia, Missouri

**MFA OIL COMPANY**