

CAUTION:

KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON BACK PANEL



CHICKEN

DUSTER

Contains 4% Malathion

For control of flies on birds. Use on Dairy Cattle,
Beef Cattle, and on all types of Wild Weather as a Fly
Dip for Horn Fly Control on Dairy and Beef Cattle.

DIRECTIONS FOR USE

FLY MAGGOT CONTROL: Dust surfaces of animal refuse thoroughly and re-dust at 3 to 5 day intervals.

POULTRY: To control poultry lice, head lice, fowl mites, chicken red mites, and other parasites to birds, dust each nest and roost thoroughly. Repeat as necessary. Where poultry houses are infested, clean out and line floor. Then apply M.F.A. Stock and Poultry Dust according to directions.

BEEF CATTLE, NON-LACTATING DAIRY CATTLE, SWINE AND HORSES: To control house flies, thorough dusting of animals is necessary, including ears and hind quarters. Repeat on non-lactating dairy cattle every 10 days of freshening. Repeat on other animals every one month or less.

HORN FLY CONTROL: Use M.F.A. Stock and Poultry Dust on both sides of head, neck, and body, including and under the wings. Repeat every 14 days. Use of 4 to 5 lbs. M.F.A. Stock and Poultry Dust on back and neck of cow will control horn fly for 14 day intervals.

CAUTION: Avoid contact with eyes, nose, mouth, and contact with contaminated clothing. Do not use in confined areas. Do not use in areas where children play. Do not use in areas where food is prepared or stored. Do not use in areas where animals are kept.




CHICKEN DUSTER

Contains 4% Malathion

For control of Lice on Hogs, Lactating Dairy Cattle, Beef Cattle and Horses. Ideal in Cold Weather as a D.P. For Horn Fly Control on all Dairy and Beef Cattle.

ACTIVE INGREDIENTS:	
Malathion	4%
INERT INGREDIENTS:	96%
	<hr/>
Total	100%

*C. O. D. is the disodium salt of the active ingredient.



NET CONTENTS: 10 LBS.

are infested, clean out and burn litter and apply M.F.A. Stock and Poultry Dust according to directions.

BEEF CATTLE AND LACTATING DAIRY CATTLE, SWINE AND HORSES: For control of lice, thorough dusting of animals, including ears and between legs, 7 to 10 days or longer, as needed. Do not dust on non-lactating dairy cattle during period of freshening. Do not apply to cows within one month of calving.

HORN FLY CONTROL: For control of horn fly on both beef cattle and dairy cattle, dusting and spraying on every 4 to 7 days with 4% Malathion. Apply to the back and neck of animal and repeat every 14 day intervals.

CAUTION: Avoid contact with humans. Avoid contact with animals during milking. Avoid contamination of feed, bedding, water and water troughs. Do not allow animals to graze later than 2 hours after application during milking. Do not use this product on dairy animals as a spray or wash.

This product is toxic to fish. Do not contaminate water by washing equipment or disposal of wastes.

Do not reuse containers. Do not wash reusable

WARRANTY: M.F.A. Co. warrants that this product conforms to the description and is suitable for the purpose stated on the label and in accordance with directions and conditions of use. M.F.A. Co. warrants no other warranty, express or implied, fitness for a particular purpose or for any implied warranty, including but not limited to, merchantability, fitness for a particular purpose, or any other implied warranty.