

MFA

GARDEN
DUST

Insecticide - Fungicide

With 4% MALATHION

ACTIVE INGREDIENTS:

*Methoxychlor Technical	5.0%
**Captan	5.0%
***Malathion	4.0%
INERT INGREDIENTS:	<u>86.0%</u>
Total	100.0%

*Equivalent to 4.4% 2,2-bis (p-methoxyphenyl) -1,1,1-trichloroethane and 0.3% other isomers and reaction products
 **N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide.
 ***O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

KEEP OUT OF REACH OF CHILDREN (12 pt.)

DANGER (18 pt.)

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
 MANUFACTURED FOR MFA OIL COLUMBIA, MO 65201

NET CONTENTS:

EPA Reg. No. 746-29
 EPA Est. 407-IA-1

I-73-84-1

ACCEPTED
 SEP 05 1986

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Harmful if swallowed or inhaled. Avoid breathing of dust. May cause allergic skin reactions. Wash thoroughly after using. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse. Remove residues remaining at harvest by washing, wiping, or similar effective means.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not apply when weather conditions favor drift from areastreated. Do not apply where run off is likely to occur. Drift and runoff from treated ares may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan when using indoors or outdoors. When using outdoors wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

Do not rotate a foliar treated crop with crops other than those with registered captan uses.

Do not allow persons to enter treated areas within 4 days following application unless protective clothing is worn. Conspicuously post reentry information at site of application.

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container, in a cool, dry, secure area away from fertilizer, food, or feed. Keep container closed.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not re-use empty container. Wrap container and put in trash.

Start application as soon as plants are set or immediately after true leaves form. Repeat every 7 to 10 days and after every rain. Follow the caution statements as regards latest treatment possible to avoid residues at harvest. Apply Garden Dust to cover both sides of leaves and the stems as well as possible. Dust on wet or dry foliage but preferably when air is still.

GARDEN DUST will control insects, and plant diseases listed below:

BEANS: Aphids, Leafhoppers, Mexican Bean Beetle, Bean Leaf Beetle, Red Spider, Thrips, Downy Mildew.

BEETS, CARROTS: Garden Web Worm, Beet Web Worm, Greenhouse Leaf Tier, Early and Late Blight, Leaf Spot. If tops are to be used for food and feed, do not apply to Beets or Carrots within 14 days of harvest.

BEST AVAILABLE COPY

CANTALOUPE, WATERMELONS, CUCUMBERS: Melon Worm, Striped Cucumber Beetle, Spotted Cucumber Beetle, Flea Beetle, Melon Aphid, Bacterial Wilt, Leaf Spot, Downy Mildew or Blight.

PEPPERS: Spinach Aphids, Anthracnose and Cercospora.

POTATOES: Early Blight, Colorado Potato Beetle, Potato Aphid, Leafhopper, Potato Thrips, Potato Flea Beetle, Potato Psyllid.

TOMATOES: Leaf Spot, Anthracnose, Early and Late Blight, Fruit Spot, Tomato Horn Worm, Tomato Fruit Worm, Flea Beetles, Aphids, Leafhoppers, Psyllid.

ROSES: Leaf Spot, Aphids, Rose Chafer, Rose Leaf Roller, Rose Midge. Surface of beds should also be dusted.

FLOWER INSECTS: Chrysanthemum Thrips and Tier Climbing Cut Worm, Tussock Moth, Japanese Beetle, Thrips, Thrips on Snapdragons, White Fly, Sowbugs.

NOTICE: Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST AVAILABLE COPY

