




ACCEPTED

NET CONTENTS: 5 GAL.



2,4-D Amine Weed Killer

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL**



WHEAT, OATS, BARLEY, AND RYE

For the control of weeds such as mustard, pig weed, sunflower, lamb's quarter, ragweed, wild vetch, cocklebur, shepherd's purse, prickly lettuce, plantain, and certain other broad-leaved weeds, use 1 1/2 pints per acre, and apply at full tiller to joint stage. (The crop is usually 6 to 10 inches high.) Do not use unless damage or destruction of any legumes present can be tolerated.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, wash with soap and water. For eyes wash with plenty of water.

Avoid spray drift to susceptible plants such as cotton, sugarbeets, clover, soybeans, tomatoes, or certain ornamentals. Coarse sprays are less likely to drift. **DON'T OVERDOSE.** Vapors from this product may injure susceptible plants in the immediate vicinity. Avoid contamination of water used for irrigation or domestic purposes.

Do not store near fertilizers, seeds, insecticides, or fungicides. Do not use weed control equipment for applying insecticides or fungicides to susceptible crops.

NOTE: Store at temperatures above 40 degrees.

Do not reuse empty drum. Destroy by perforating or crush and burying in a safe place.

WARRANTY: M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company, and buyer assumes the risk of any such use. B71

GENERAL INFORMATION: Amine Weed Killer mixes readily with water and is best used for weed control under moderate conditions and where crop damage by ester formulations must be avoided.

DIRECTIONS FOR USE:—This product may be used for the control of many broad leaved weeds growing in wheat, oats, barley, rye, pastures, lawns, and corn.

In general, this product is most effective when weeds are small and growth active. It may be applied when temperatures range from 50 to 95° F. Results may be somewhat slower below 70° and above 90°. If possible, do not spray if rain is expected within 4 hours. Heavy dew does not usually affect results with low volume applications. This material may be applied with all types of spraying equipment. Mix only with water. Do not use on any grain crop within 5 days of harvest.

CORN AND SORGHUMS

POST EMERGENCE: For control of Ragweed, Morning Glory, Sunflower, Plantains, Lamb's Quarter, Mustard, Spanish Needle, Pig Weed, Cocklebur, and certain other broad leaved weeds. On field corn apply when crop is over 4 inches high. On sorghums apply when crop is 4 to 18 inches high. Use 1 pint per acre and spray to cover weed tops. Use nozzle drops when crop is taller than 10 inches to reduce contacting crop. Do not cultivate for ten days after spraying. Do not spray either crop after tasseling.

PASTURES AND LAWNS

For control of ragweed, bitterweed, cocklebur, mustard, sheep sorrel, wild carrot, pig weed, dandelion, and certain other broad leaved weeds. If legumes are present use 1 quart per acre. If ladino or Kozan, espedeza are present and some damage can be tolerated use 1/2 pint per acre. If other legumes are present do not use unless legume destruction can be tolerated.

Do not use on seed crops or creeping bent. New grass seedlings should not be sprayed until they are at least two months old. Do not use on St. Augustine grass and Dichondra.

Do not graze treated areas by dairy animals within 7 days of spraying.



ACCEPTED
 APR 2 1971
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND HERBICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 141120

NET CONTENTS: 5 GAL.



2,4-D Amine Weed Killer

**CAUTION: KEEP OUT OF REACH OF CHILDREN
 SEE OTHER CAUTIONS ON SIDE PANEL**



GENERAL INFORMATION: Amine Weed Killer mixes readily with water and is best used for weed control under moderate conditions, and where crop damage by ester formulations must be avoided.

DIRECTIONS FOR USE—This product may be used for the control of many broad-leaved weeds growing in wheat, oats, barley, rye, pastures, lawns, and corn.

In general, this product is most effective when weeds are small and growth is active. It may be applied when temperatures range from 50° to 95° F. Results may be somewhat slower below 70° and above 90°. If possible, do not spray if rain is expected within 4 hours. Heavy dew does not usually affect results with low volume applications. This material may be applied with all types of spraying equipment. Mix only with water. Do not use on any grain crop within 5 days of harvest.

CORN AND SORGHUMS

POST EMERGENCE: For control of Ragweed, Morning Glory, Sunflower, Plantains, Lamb's Quarter, Mustard, Spanish Needle, Pig Weed, Cocklebur, and certain other broad-leaved weeds. On field corn apply when crop is over 4 inches high. On sorghums apply when crop is 4 to 18 inches high. Use 1 pint per acre and spray to cover weed tops. Use nozzle drops when crop is taller than 10 inches to reduce contacting crop. Do not cultivate for ten days after spraying. Do not spray either crop after tasseling.

PASTURES AND LAWNS

For control of ragweed, bitterweed, cocklebur, mustard, sheep sorrel, wild carrot, pig weed, dandelion, and certain other broad-leaved weeds: If no legumes are present, use 1 quart per acre. If ladino or Korean lespedeza are present and some damage can be tolerated, use at 1 pint per acre. If other legumes are present, do not use unless legume destruction can be tolerated.

Do not use on seed crops or creeping bent. New grass seedlings should not be sprayed until they are at least two months old. Do not use on St. Augustine grass and Dichondra.

Do not graze treated areas by dairy animals within 7 days after spraying.

WHEAT, OATS, BARLEY, AND RYE

For the control of weeds such as mustard, pig weed, sunflower, lamb's quarter, ragweed, wild vetch, cocklebur, shepherd's purse, prickly lettuce, plantain, and certain other broad-leaved weeds, use 1 1/2 pints per acre, and apply at full tiller to joint stage. (The crop is usually 6 to 10 inches high.) Do not use unless damage or destruction of any legumes present can be tolerated.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin or clothing. In case of contact, wash with soap and water. For eyes wash with plenty of water.

Avoid spray drift to susceptible plants such as cotton, sugarbeets, clover, soybeans, tomatoes, or certain ornamentals. Coarse sprays are less likely to drift. **DON'T OVERDOSE.** Vapors from this product may injure susceptible plants in the immediate vicinity. Avoid contamination of water used for irrigation or domestic purposes.

Do not store near fertilizers, seeds, insecticides, or fungicides. Do not use weed control equipment for applying insecticides or fungicides to susceptible crops.

NOTE: Store at temperatures above 40 degrees.

Do not reuse empty drum. Destroy by perforating or crush and burying in a safe place.

WARRANTY: M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company, and buyer assumes the risk of any such use. B71

USDA Reg No 746 35