





9-12-22
746-40



WARNING KEEP OUT OF REACH OF CHILDREN
See other warnings on side panel

DIRECTIONS FOR USE M.F.A. LIVESTOCK TOXAPHENE

For controlling Horn Flies, Lice and Ticks on Beef Cattle, Hogs, Sheep and Goats dilute with water and use either as spray or dip.

Amount M.F.A. Livestock Toxaphene	Amount Water
1 gallon	150 gallons
1 quart	37 gallons
1 pint	19 gallons
1 cup	4 gallons
1 tablespoon	1 gallon

CAUTION: Do not apply to dairy animals within 4 weeks of lactation or animals under 6 months of age.

Apply to other animals during cold, stormy weather or during a period of stress.

HORN FLY CONTROL: Mix 1 part concentrate with 150 parts of water. Repeat when necessary. For Lice and Ticks: Apply 1 part concentrate to 37 parts of water. For Sheep and Goats: 1 part concentrate to 19 parts of water. For Hogs: 1 part concentrate to 4 parts of water. For Cattle: 1 part concentrate to 150 parts of water.

FOR BACKRUBBER USE: Dilute 1 part concentrate with 4 parts of water. Apply to the back of the animal. Repeat when necessary.

livestock[®] toxaphene

(emulsifiable concentrate)


(CONTAINS 6 POUNDS TECHNICAL TOXAPHENE PER GALLON)

ACTIVE INGREDIENTS	
Toxaphene - Technical Chlorinated Camphene containing 67% to 69% Chlorine	59.00%
Petroleum Distillate	28.50%
Xylene Range Solvent	6.63%
INERT INGREDIENTS:	5.87%
Total	100.00%

For Controlling Hornflies, Lice and Ticks on Beef Cattle, Hogs, Sheep and Goats

CONTENTS 1 GALLON



MANUFACTURED FOR



COLUMBIA MO. 65201

WARNING KEEP OUT OF REACH OF CHILDREN

See other warnings on side panel



Livestock[®] toxaphene

(emulsifiable concentrate)

CONTAINS 6 POUNDS TECHNICAL TOXAPHENE PER GALLON

ACTIVE INGREDIENTS:


Toxaphene (Technical Chlorinated Camphene containing 67% to 69% Chlorine)	59.00%
Petroleum Distillate	28.50%
Xylene Range Solvent	6.63%
INERT INGREDIENTS:	5.87%

Total 100.00%

For Controlling Hornflies, Lice and Ticks on Beef Cattle, Hogs, Sheep and Goats.

CONTENTS 1 GALLON

MANUFACTURED FOR



COLUMBIA, MO. 65201

DIRECTIONS FOR USE M.F.A. LIVESTOCK TOXAPHENE

Use only on Beef Cattle, Hogs, Sheep and Goats. Do not use on animals within 4 weeks of slaughter, or under six months old.

LICE: Dip or spray. Thorough coverage of the entire animal is necessary for control. Treat about November 1 and repeat in 15 days. Do not treat sows within 30 days of farrowing or while they are suckling young.

TICKS: Dip or spray for thorough animal coverage. Apply when ticks are first noticed and repeat as necessary for control.

WARNING: Do not take internally. May be fatal if swallowed.

ANTIDOTE: Give 1 tablespoon of table salt in glass of warm water and repeat until vomit is clear. Call physician. Avoid contamination of feed and food stuffs. Do not breathe spray mist. Do not get in eyes or on skin or clothing. In case of spillage on skin, wash with soap and water.

Do not use on household pets, dairy animals, animals being raised for slaughter, or humans. Do not use in dairy barns or milk rooms. Do not store near fire or flame. To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material or washing from spray tank. Do not reuse container. Destroy when empty.

WARRANTY: M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions or under conditions not reasonably foreseeable to M.F.A. Oil Company.