

AC

GENERAL INFORMATION: No. 6 Weed Kill mixes readily with water or oil. The ester forms of 2,4-D are generally effective for weed control under severe conditions, such as dry or very wet weather, and on hard-to-kill perennials.

DIRECTIONS FOR USE

This product may be used for the control of many broad-leaved weeds growing in wheat, oats, barley, rye, pastures, and corn. In general, this product is most effective when weeds are small and growth is active. It may be applied when temperatures range from 50° to 95°F. Results may be somewhat slower below 70° and above 90°F. If possible, do not spray if rain is expected within 2 hours. Heavy dew does not usually affect results with low volume applications. This material may be applied with all types of spraying equipment. Mix only with water for crop use. Do not use on any grain crop within 5 days of harvest.

CORN

Pre-Emergence: For control of Giant and Common Foxtail, Crabgrass, Ragweed, Pig Weed, Cockle Bur, and certain other annual grasses and broad-leaved weeds, after planting and before the crop comes up, apply 1/2 to 1 1/2 quarts per acre. Use 1/2 quarts on loam and 1 1/2 quarts on clay soils. If soil is disturbed, weed killing effect is lost. Do not use on sandy soil.

CORN AND SORGHUMS

Post-Emergence: For control of Field Bindweed, Ragweed, Morning Glory, Sunflower, Plantains, Curly Dock, Jimson Weed, Lamb's Quarter, Mustard, Spanish Needle, Pig Weed, Smartweed, Cockle Bur, and certain other broad-leaved weeds. On field corn apply when crop is over 4 inches high. On sorghums apply when crop is 4 to 38 inches high. Use 1/2 pint per acre and spray to cover weed tops. Use drop nozzles when crop is taller than weeds to reduce spray contacting crop. Use drop nozzles when crop is over 10 inches tall. Do not spray after either crop starts to tassel. Do not cultivate for 10 days after spraying.

CORN

Lay-by Spray: To reduce growth of both annual grasses and broad-leaved weeds after lay-by cultivation, apply 1/2 quart per acre. Use drop nozzles to direct spray on soil at base of corn, and between rows. Do not apply within 2 weeks of silking.

WHEAT, OATS, BARLEY AND RYE

For control of weeds such as mustard, pig weed, sunflower, lamb's quarter, ragweed, wild vetch, cockle bur, smart weed, hedge bindweed, shepherds purse, prickly lettuce, plantain, docks, and certain other broad-leaved weeds, use 1/2 pint per acre, and apply at full tiller to joint stage. (The crop is usually 6 to 10 inches high.) Do not use unless damage or destruction of any legume present can be tolerated. Do not forage or graze treated grain fields within 2 weeks after treatment. Do not feed treated straw to livestock.

ACCEPTED
MAY 29 1981
For the Federal Insecticide,
Fungicide and Fertilizer Act,
registered under the pesticide
EPA Reg. No. 146-43

NET CONTENTS

NO. 6

Weed Kill
(BUTYL ESTER 2,4-D)

ACTIVE INGREDIENTS:

Butyl Ester of 2,4-Dichlorophenoxyacetic acid* 74.4%

INERT INGREDIENTS:

Total

*Isomer Specific by AOAC Method No. 6. D01-5
*Equivalent to 59.5% 2,4-Dichlorophenoxyacetic acid or 1.5 lbs. per gallon.

GENERAL INFORMATION ON APPLICATIONS OF NO. 6 WEED KILL

CROP	AMOUNT NO. 6 WEED KILL PER ACRE
Corn - Sorghums - Susceptible Weeds	1/2 pint
For Pre-Emergence Spraying - Corn	1/2 to 1 1/2 quarts
Small Grains - Susceptible Weeds	1/2 pint
Pastures - Susceptible Weeds	1/2 pint
Pastures - Onions, Garlic, Canada Thistle	1/2 quart

CAUTION:

KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

Manufactured For



Columbia, Missouri 65201

86



NET CONTENTS



NO. 6

Weed Kill

(BUTYL ESTER 2,4-D)

ACTIVE INGREDIENTS:
Butyl Ester of 2,4-Dichlorophenoxyacetic acid* 74.4%

INERT INGREDIENTS: 25.6%

Total 100.0%

*Boomer Specific by AOAC Method No. 5. D01-5
*Equivalent to 59.5% 2,4-Dichlorophenoxyacetic acid or 5.0 lbs. per gallon.

GENERAL INFORMATION ON APPLICATIONS OF NO. 6 WEED KILL

CROP	AMOUNT NO. 6 WEED KILL PER ACRE
Corn - Sorghums - Susceptible Weeds	1/2 pint
For Pre-Emergence Spraying - Corn	1/2 to 1 1/2 quarts
Small Grains - Susceptible Weeds	1/2 pint
Pastures - Susceptible Weeds	1/2 pint
Pastures - Onion, Garlic, Canada Thistle	1/2 quart

CAUTION:
KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

Manufactured For  Columbia, Missouri 65201

86

PASTURES

For wild onion and garlic control. Apply 1 1/2 quarts per acre in early spring and again in late fall for 2 successive years. Do not use if legume destruction cannot be tolerated. Do not use on St. Augustine grass or Dichondra.

For control of sugarcane, bitterweed, cocklebur, nuttall's burdock and other broad-leaved weeds. If no legumes are present use 1/2 quart per acre. If no legumes are present and some damage can be tolerated use 3/4 pint per acre. If other legumes are present, do not use unless legume damage can be tolerated.

Do not use on seed crops or creeping bent. New grass seedlings should not be sprayed until they are at least two months old. Do not graze by dairy animals within 7 days after spraying.

WOODY PLANTS

Use from 1 1/2 to 2 1/2 quarts per 100 gallons of water for susceptible species such as willow, sumac, and buckbrush found in wasteland; for other species, use M.F.A. Super Brush-Kill.

CAUTION: Harmful if swallowed. Avoid contact with eyes, skin, or clothing. In case of contact, wash with soap and water. In case of contact with eyes, flush with plenty of water.

DON'T OVERDOSE: Avoid spray drift to susceptible plants such as cotton, sugarbeets, clover, soybeans, tomatoes, or certain ornamentals. Coarse sprays are less likely to drift. Vapors from this product may injure susceptible plants in the immediate vicinity.

Do not store near fertilizers, seeds, insecticides, or fungicides. Do not use weed control equipment for applying insecticides or fungicides to susceptible crops. Avoid contamination of water used for irrigation or domestic purposes.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not reuse container. Destroy by burying in a safe place. Do not burn.

NOTE: Store at temperatures above 60° F.

WARRANTY: M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company and buyer assumes the risk of any such use.

EPA Reg. No. 746-43
EPA Est. No. 407-1A-1

L-79-U-9902B-D

87