

JUN 3, 11

BEEF CATTLE

HORN FLY CONTROL SPRAY: Use 1 cup per 3 gallons or 2 gallons per 100 gallons of water. Thoroughly cover each animal using about 2 quarts of spray. Repeat after 2 to 3 weeks or when flies reappear. Do Not apply on dairy cattle.

HORN FLY CONTROL BACKRUBBER: Beef Cattle Use Only. Dilute 1 quart to 1 gallon of fuel oil. Locate backrubber in a place where cattle normally loaf during summer such as salt licks, shade, and water troughs. Initially charge backrubber to saturation. Recharge when necessary for control, or within two weeks. DO NOT let cattle use a backrubber that is dripping with the insecticide oil mixture.

BEEF CATTLE LICE, HOG LICE AND SHEEP TICKS:

Use 1 cup per 3 gallons or 2 gallons per 100 gallons of water. Cover animals thoroughly, including inside of ears and bush of tail using 2 quarts per hog or sheep and 4 quarts per bee animal. Repeat after 15 days except for sheep ticks.

ALFALFA AND CLOVER:

For control of Blister Beetle, Alfalfa Weevil, Clover Leaf Weevil, Leaf Hoppers, Webworms and Alfalfa Caterpillar. Apply 3 quarts per acre in water as determined by sprayer calibration. Apply to give complete coverage of foliage. Begin spraying at first signs of infestation, and repeat when needed. DO NOT spray during bloom nor apply within 7 days of cutting or grazing.



1 QT.

METHOXYCHLOR E.C.

**FOR USE ON:
FIELD CROPS GRAIN BINS
BEEF CATTLE**

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL
(CONTAINS 2 LBS. METHOXYCHLOR PER GAL.)**

ACTIVE INGREDIENTS:	
*Methoxychlor, Technical	24.0%
Aromatic Petroleum Hydrocarbons	72.5%
INERT INGREDIENTS:	3.5%
<hr/>	
Total	100.0%

*Consists of 21% 2, 2-Bis(p-Methoxyphenyl)-1, 1, 1-Trichloroethane and 3% other isomers and related compounds



Manufactured For **MFA OIL** Columbia, Missouri

USDA Reg. No. 746-60

GRAIN BINS

Apply as residual spray to aid in control of larvae and adult stages of Cadelle, Sawtoothed Grain Beetle, Bran bugs and certain other insects attacking stored grain. Mix Methoxychlor E.C. at 1 quart per 9 quarts, or 1 gallon per 9 gallons of water. Apply to the thoroughly cleaned bin until surfaces are wet but not until run-off. 2 gallons will spray about 1,000 square feet.

CAUTION: Harmful if swallowed. Avoid inhalation or contact with skin. Wash with soap and water after using. Do not use on food or feed crops within 7 days of harvest. Remove spray deposits from plant products consumed as food. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on the label. This product is toxic to bees and should not be applied when bees are actively visiting the area. Do Not Contaminate Water By Cleaning Of Equipment Or Disposal Of Wastes

WARRANTY: M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company, and buyer assumes the risk of any such use.

NOTICE: This product will tend to settle out under certain temperature conditions. Do not store in temperatures below 40° F. If it settles, warm to 80° F. and agitate thoroughly before using.

CONTAINER DISPOSAL Do not reuse container. Destroy when empty.