O Ш

DIRECTIONS FOR USE

[746-69]

Feed Storage-Far	m Buildings	Industrial Use	s
For control of Co	delle, Angoum	ols Grain Mo	th (Exposed
Stages) Confused	Flour Seetle,	Red Flour	Beetle, Flat
Grain Bootle, Saw	Indiana Orain	Beefle, Indian	Meal Moth
(Exposed Stages)	, Cockroaches	, Ants, Silv	erfish, and
Chanse Mites		۵۳۵، ۲۹۵، ۲۵۱، ۲۵۰، «یود پیش زاد کار ۲۵۱، ۲۵۱، ۲۵۰، ۲۵۰، ۲۵۰، ۲۵۰، ۲۵۰، ۲۵۰، ۲۵۰، ۲۵۰	

Spray all areas where insects hide, crawl, or rest, with part Pyrenone E.C. to 9 parts water, Apply at the rate of 1 gallon of mixture per 750 square feet of surface. for control of roaches, and silverfish, thorough and repeated spraying of the infested areas will be needed. Direct spraying of exposed insect should be done when possible. For control of ants follow the above directions for roaches and silverfish, plus treatment of ant trails and areas around door sills and window frames where the ants enter the premises will be needed. Repeated applications will be required in all cases.

DO NOT apply in the edible products area of food processing plants.

For fogging purposes, to kill exposed adults of Granary Weevil, Rice Weevil, Cadelle and Angoumois Moth, dilute with kerosene of water depending on instructions and calibration of the fogger. Apply at one pint of Pyrenone E. C. per 40,000 cubic feet, of space. Repeat as necessary.

DIRECTIONS FOR USE-LIVESTOCK

Stableflies Horseflies Deerflies Dilute 1 to 9 with water. Spray to wet hair over entire animal. Thoroughly cover legs and underline. On dairy animals make application 15 minutes before milking time. Spray applications on all livestock should be made 2 to 3 times a week or as needed.

As an alternative control method against attacks by horseflies, stableflies, and deerflies, on dairy cows and cattle, apply just enough undiluted Pyrenone spray to wet the. hair but not enough to soak the hide; using one or two ounces per animal the size of an adult cow. Apply once daily, repeating at daily intervals or as necessary to give continued protection.


TO CONTROL HORNFLIES AND MOSQUITOES: Dilute 1 to 19 with water and apply as above.

3



KEEP OUT OF REACH OF CHILDREN. SEE OTHER CAUTIONS ON SIDE PANEL.

CAUTION



LIVESTOCK, GRAIN ELEVATORS FARM BUILDINGS, FOOD PROCESS-ING PLANTS, INDUSTRIAL USES.

ACTIVE INGREDIENTS:

Petroleum Technical	Solvents Piperonyl															
Pyrethrins INERT INGRE	DIENTS .	• •	•	•	•	•	•	•	•	•	•	•	•	•	•	

*Equivalent to 8.0% (butylcarbityl) (6-propylpiperonyl) ether and 2.0% related compounds. *T. M. Reg. U. S. Pat. Off. by Food Machinery Corp.

Manufactured For







wet all parts of the animal. About $1\frac{1}{2}$ to 3 gallons is required for cattle and proportionally less for hogs and sheep. Repeat as needed.

FLEAS ON DOGS, HORSEL CATTLE, HOGS AND SHEEP

Dilute 1 to 100 with water. Theroughly wet coat of animal. Spray animal quarters, bedding and other areas where fleas occur. Repeat treatment as needed.

CHICKEN MITES AND BEDBUGS IN POULTRY HOUSES:

Ditute 1 10 9 with water. Thoroughly spray roosts, nests, litter, and cracks in floors and walls. Repeat treatment as needed.

R	ecommended	Use This Amt. of Concentrate to Make:							
	Dilution	1 Quart	1 Gallon	3 Gallon					
1	to 9	7 Tblsp.	3/4 Pint	21/4 Pints					
1	to 19	3 Thisp.	1/2 Pint	1½ Pints					
1	to 100	2 Teaspoons	21/2 Thisp.	1/4 Pint					

DILUTION TABLE

CAUTION: Harmful if swallowed. Keep out of reach of children. Wash hands after using, with soap and water. Do not take internally.

WARRANTY: M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company, and buyer assumes the risk of any such use. K-70

USDA Reg. No. 746-69

,100%

Columbia, Missouri