

MFA  
PYRENONE E.C.

PM 1

746-69  
P. 10/3

FOR:  
LIVESTOCK, GRAIN ELEVATORS, FARM BUILDINGS  
FOOD PROCESSING PLANTS, INDUSTRIAL USES.

ACTIVE INGREDIENTS:

|                                   |            |
|-----------------------------------|------------|
| Technical Piperonyl Butoxide..... | 10%        |
| Pyrethrins.....                   | 1%         |
| INERT INGREDIENTS.....            | 6%         |
| Petroleum Solvents.....           | <u>83%</u> |

TOTAL 100%

\*\*Equivalent to 8.0% (butyl carbityl)(6 propyl piperonyl) ether and 2.0% related compounds.

R Pyrenone -- trademark of Fairfield American Corporation

KEEP OUT OF THE REACH OF CHILDREN  
CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured for  
MFA OIL  
Columbia, MO 65201


NET CONTENTS \_\_\_\_\_

EPA Reg. No. 746-69  
EPA Est. No. 407-IA-1

B-71

1.17.95

Not Reviewed. Registrant claims to be in accordance with  
EPA guidance for Phase 2  
pesticide registration change in status.



4202

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if swallowed. Wash hands after using, with soap and water. Do not take internally.

**ENVIRONMENTAL HAZARDS**

Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE:** Store in original container, in a cool, dry, secure area, away from fertilizer, food or feed. Keep container closed.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water, food, or feed by storage or disposal.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then dispose of in a sanitary landfill or by other approved State and local procedures.

**Feed Storage -- Farm Buildings -- Industrial Uses.**


For control of Cadelle, Angoumois Grain Moth (Exposed Stages), Confused Flour Beetle, Red Flour Beetle, Flat Grain Beetle, Saw-toothed Grain Beetle, Indian Meal Moth (Exposed Stages), Cockroaches, Ants, Silverfish, and Cheese Mites.

Spray all areas where insects hide, crawl or rest, with 1 part Pyrenone E.C. to 9 parts water. Apply at the rate of 1 gallon of mixture per 750 square feet of surface. For control of roaches, and silverfish, thorough and repeated spraying of the infested areas will be needed. Direct spraying of exposed insects should be done when possible. For control of ants follow the above directions for roaches and silverfish, plus treatment of ant trails and areas around door sills and window frames where the ants enter the premises will be needed. Repeated application will be required in all cases.

For fogging purposes, to kill exposed adults of Granary Weevil, Rice Weevil, Cadelle and Angoumois Moth, dilute with kerosene or water depending on instructions and calibration of the fogger. Apply at one pint of Pyrenone E.C. per 40,000 cubic feet of space. Repeat as necessary.

**LIVESTOCK Horseflies -- Stableflies -- Deerflies**

Dilute 1 to 9 with water. Spray to wet hair over entire animal. Thoroughly cover legs and underline. On dairy animals make application 15 minutes before milking time. Spray applications on all livestock should be made 2 to 3 times a week or as needed.



As an alternative control method against attacks by horseflies, stableflies, and deerflies, on dairy cows and cattle, apply just enough undiluted Pyrenone spray to wet the hair but not enough to soak the hide; using one or two ounces per animal the size of an adult cow. Apply once daily, repeating at daily intervals or as necessary to give continued protection.

TO CONTROL HORNFLIES AND MOSQUITOES: Dilute 1 to 19 with water and apply as above.

LICE ON HORSES, CATTLE, HOGS AND SHEEP: Dilute 1 to 100 with water. Apply as a spray or dip to wet all parts of the animal. About 1 1/2 to 3 gallons is required for cattle and proportionally less for hogs and sheep. Repeat as needed.

FLEAS ON DOGS, HORSES, CATTLE, HOGS AND SHEEP: Dilute 1 to 100 with water. thoroughly wet coat of animal. Spray animal quarters, bedding and other areas where fleas occur. Repeat treatment as needed.

CHICKEN MITES AND BEDBUGS IN POULTRY HORSES: Dilute 1 to 9 with water. Thoroughly spray roosts, nests, litter, and cracks in floors and walls. Repeat treatment as needed.

**DILUTION TABLE**

| Recommended Dilution | Use this Amt. Concentrate to Make: |              |             |
|----------------------|------------------------------------|--------------|-------------|
|                      | 1 quart                            | 1 gallon     | 3 gallon    |
| 1 to 9               | 7 Tblsp.                           | 3/4 pint     | 2 1/4 pints |
| 1 to 19              | 3 Tblsp.                           | 1/2 pint     | 1 1/2 pints |
| 1 to 100             | 2 Teaspoons                        | 2 1/2 Tblsp. | 1/4 pint    |

