

**CAUTION: KEEP OUT OF REACH OF CHILDREN  
SEE OTHER CAUTIONS BELOW**

**CONTAINS 4% MALATHION  
FOR CONTROL OF:**

**Mites and lice on chickens, ducks, geese, turkeys, beef cattle, non-lactating dairy cattle, swine and horses. For control of Horn Fly on all dairy and beef cattle. Ideal for cold weather use.**

ACTIVE INGREDIENTS:  
Malathion\*

4%

INERT INGREDIENTS:

96%

Total

100%

\*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**GENERAL INSTRUCTIONS**

**POULTRY:** to control poultry lice, and northern fowl mites, apply to litter at 1 pound per 40 square feet to control chicken red mites apply at 1 pound per 60 square feet. Put a tablespoon of dust in each nest and dust roosts individually. Repeat as necessary. Dust individual birds with shaker can, puff type or rotary hand duster for direct application when needed.

**POULTRY DUSTING BOX:** For control of northern fowl mites and poultry lice, place one self-treatment dust box (18 x 12 x 3 inches deep), containing one pound Stock & Poultry Dust per each 30 hens. Allow hens to dust themselves freely. Repeat treatment as necessary. Avoid contaminating feed, feeding troughs, and water fountains.

**FLY MAGGOT CONTROL:** Dust surfaces of animal refuse thoroughly and repeat at 3 to 4 day intervals.

**CATTLE LICE:** For control of lice on both non-lactating dairy cattle and beef cattle, dust the animal thoroughly. Repeat in 10 days and as needed thereafter. Do not treat animals under one month of age. Do not apply on non-lactating dairy animals within 2 weeks of freshening.

**HORN FLY CONTROL:** For control of horn flies on both beef and dairy cattle,

apply 4 tablespoons of Stock and Poultry Dust per animal. Apply on the back and neck of animal and repeat in 10 to 14 day intervals.

**HOG LICE:** To control lice on hogs make thorough application to animals and dust pens thoroughly. Repeat in 10 days and thereafter as needed.

**CAUTION:** Avoid prolonged breathing of dust. Avoid contact with skin. Keep out of reach of children.

Avoid contamination of feed, feeding troughs and water fountains. Do not apply to dairy cattle within 5 hours before milking or during milking. Do not use Malathion on dairy animals as a spray or dip.

Do not reuse empty containers. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

**WARRANTY:** M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company, and buyer assumes the risk of any such use. C-71

Manufactured For  MFA OIL Columbia, Missouri

**TEN POUNDS NET CONTENTS**

USDA Reg. No. 746-85

ACCEPTED  
May 19, 7/46-85

# MFA STOCK AND POULTRY DUST