

WARNING
KEEP OUT OF REACH OF CHILDREN
 See other warnings on back panel.



25% LINDANE

25% WETTABLE POWDER

ACTIVE INGREDIENTS:

Gamma Isomer of Benzene Hexachloride from Lindane	25%
INERT INGREDIENTS	75%
	100%

FOR:

**CONTROL OF SARCOPTIC OR PSOROPTIC
 MANGE ON BEEF CATTLE AND SWINE —
 CONTROLLING LICE ON SWINE — LIMITED
 USE FOR CUTWORM CONTROL IN GARDENS
 — SEED TREATMENT AGAINST WIREWORM
 AND SQUASH BUG ON CUCURBITS.**

MANUFACTURED FOR  OIL Columbia, Missouri 65201

NET CONTENTS 1 POUND

M. F. A. 25% LINDANE

Directions for Use

SARCOPTIC OR PSOROPTIC MANGE ON BEEF CATTLE: (sponge use only) Sponge infected areas with 2 level tablespoons 25% Lindane in 1 gal. water. Do not use on new-born (very young) animals or calves under 3 months of age, or on dogs, cats or other household pets, or on poultry or in poultry houses. If calves between 3 to 6 months of age must be treated use 1 level tablespoon per gallon water and sponge infected area. Wear protective hand coverings. Repeat applications at 7 - 10 day intervals for 2 or more treatments as needed. Quarantine infected cattle immediately when scab is found and observe other cattle carefully for a spread of the condition. **DO NOT** treat within 30 days of slaughter. Treat calves only at lower dosage rate. **DO NOT** use on dairy animals. **DO NOT** treat overheated, weak or sick animals. Avoid treating animals in cold stormy weather.

HOG LICE AND SARCOPTIC MANGE: (spray or sponge use only) Thoroughly mix 1 to 2 pounds 25% Lindane in 100 gals. water (1 to 2 level tablespoons per gal.). Routinely treat pigs when taken from sows at weaning time, periodically treat sows 30 to 45 days before farrowing. Do not treat sows within 2 weeks of farrowing or 3 weeks thereafter. Treat boars before breeding season. Spray as soon as mange is found. Spray feeder pigs, in fall and other animals before they are added to herd. Thoroughly wet each animal. Repeat application at 7 - 10 day intervals for 2 or more treatments as needed. **DO NOT** spray within 30 days of slaughter. **DO NOT** treat pigs under three months of age. **DO NOT** treat overheated, sick or weak animals. Avoid treating animals in cold stormy weather.

CUTWORM CONTROL ON TOMATOES, CABBAGE AND PEPPERS: Mix 1 teaspoon 25% Lindane with each gallon of transplanting water.

SQUASH BUGS ON CUCURBITS, (CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS): Mix 1 teaspoon 25% Lindane and 4 teaspoons M.F.A. 50% Methoxychlor per gallon of water and spray affected plants thoroughly. Repeat every two weeks. Applications to cucumbers and squash after fruit forms may reduce the quality of harvest. There must be a one day pre-harvest interval on cucumbers and squash.

WARNING: May be fatal if swallowed. May be absorbed through the skin. Do not breathe dust or spray mist. Avoid contact with skin and eyes. Wash thoroughly after using. Keep out of reach of children. Avoid contamination of feed and foodstuffs.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish and wildlife. Keep out of lakes, streams and ponds. **DO NOT** contaminate water by cleaning of equipment or disposal of waste. Apply only as specified on label.

DO NOT reuse this container. Destroy by burying or burning in a safe place. Stay out of smoke or fumes.

NOTICE: M.F.A. Oil Company warrants that this product conforms to the name and description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extend to the use of this product contrary to label instructions, or under abnormal conditions.