


AM
 FUNG
 EC
 E



Caution: Keep Out of Reach
 of Children.

1 GAL.

DIRECTIONS FOR USE

ORANGE CROP INSECTS: Caution: Do not apply to crops in bloom. Do not apply within 7 days of harvest or pasturing.

ALFALFA: For control of young grasshoppers, aphids (including yellow clover aphid, spotted alfalfa aphid, pea aphid), potato leaf hopper, spider mites and alfalfa weevil larvae: Apply 1/5 gallon M.F.A. Malathion Emulsifiable Concentrate in sufficient water to cover an acre . . . when temperatures are over 65 degrees Fahrenheit and winds under 10 to 15 miles per hour.

CLOVER: For control of aphids, mites, leaf hoppers and young grasshoppers: Use 1/5 gallon M.F.A. Malathion Emulsifiable Concentrate in sufficient water to cover one acre. Repeat applications as necessary.

CORN LEAF APHIDS ON CORN: Use 1/5 gallon M.F.A. Malathion Emulsifiable Concentrate in sufficient water to cover one acre. If aphids are working in leaf sheath, plug nozzles between rows and use as much volume of water as possible . . . directing spray into whorl.

NON-WARRANTY - Seller (whether Manufacturer, Distributor or Other), makes no warranty, express or implied, regarding this product except the composition as set forth in the ingredient statement herein. Buyer or user assumes all risk of possession, handling or use, whether in accordance with any directions as contained herein or not.

MALATHION

EC


FOR POTENTIALLY HARMFUL CROP INSECTS:

Spotted Alfalfa Aphid, Pea Aphid, Yellow Clover Aphid, Cotton Aphid, Spider Mites, Potato Leaf Hopper and Certain Other Crop and Forage Insects and For Control of Certain Poultry Insects.

ACTIVE INGREDIENTS:

Malathion*	57%
Aromatic Petroleum Derivative Solvent	35%
INERT INGREDIENTS	8%
<hr/>	
Total	100%

*O, O-Dimethylthiophosphate of diethylmercaptosuccinate
 (Contains 5 pounds Malathion per gallon)

Manufactured For  M.F.A. Chemicals, Inc., Missouri

ACCEPTED WITH COMMENTS

Caution: Keep Out of Reach
of Children.

1 GAL.

MALATHION

EC

CONTROL OF INSECT PESTS:

Control Alfalfa Aphid, Pear Aphid, Yellow Clover Aphid, Cotton Aphid, Spider, Potato Leaf Hopper and Certain Crop and Forage Insects and For Control of Certain Poultry Insects.

ACTIVE INGREDIENTS:

Malathion*	57%
Aromatic Petroleum Derivative Solvent	35%
Other Inert INGREDIENTS	8%

Total.....100%

*O-Dimethyldithiophosphate of diethylmercaptosuccinate

(Contains 5 pounds Malathion per gallon)

Manufactured For



Chemical Company, Missouri

CUTWORMS ON PASTURES BEING GRAZED BY DAIRY CATTLE:

To control Black, Dingy, Variegated, and Bronze Cutworms in Orchard Grass, Fescue, Birdsfoot Trefoil, Alfalfa, Ladino Clover, Red Clover and similar crops, use 1 quart Malathion E. C. per acre. Apply in sufficient water to give complete coverage. Keep cattle off treated area for 7 days.

POULTRY APPLICATION FOR CONTROL OF NORTHERN FOWL MITE AND POULTRY LICE:

Mix 2 tablespoons of Malathion per gallon of water. Apply a gallon of mixture per 100 to 150 birds. Repeat in 4 to 8 weeks or as necessary. Also spray infested poultry houses with 4 tablespoons Malathion per gallon of water. Apply liberally to walls, ceilings, roosts, nests and adjacent areas forcing the spray into all cracks and crevices. For poultry tick control increase this dose rate to 1/2 pint per gallon of water. Repeat as necessary. Use a high pressure sprayer for application on surfaces.

COTTON:

For control of cotton aphid, use 1 pint M.F.A. Malathion Emulsifiable Concentrate per acre diluted with sufficient water for coverage. Apply when temperature is over 60 degrees Fahrenheit and wind is under 10 to 15 miles per hour.

CAUTION: HAZARDOUS IF SWALLOWED. DO NOT BREATHE

SPRAY MIST. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. If accidentally swallowed, give emetic of mustard and call physician. In case of spillage on skin, wash with soap and water. Do not use on pets, livestock or humans. Do not use in homes, milk houses or other buildings where food is processed. Do not store near fire or flame, to protect fish and wildlife, do not contaminate streams, lakes, or ponds with this material.

DIRECTIONS FOR USE

ALFALFA: For control of grasshoppers, aphids (including yellow clover aphid, spotted alfalfa aphid, pea aphid), armyworms, clover leaf weevil, leafhoppers, lygus bugs, spider mites and spittle bug; apply 1½ to 2 pints MFA Malathion Emulsifiable Concentrate in sufficient water to cover an acre . . . when temperatures are over 67 degrees Fahrenheit and winds under 10 to 15 miles per hour. May be used up until day of harvest.


CLOVER: For control of aphids, clover leaf weevil, grasshoppers, leafhoppers and mites, use 1½ to 2 pints MFA Malathion Emulsifiable Concentrate in sufficient water to cover one acre. Repeat applications as necessary. May be used up until day of harvest. **NOTE:** For protection of bees, apply to alfalfa and clover in bloom only in evening or early morning.

CORN: For control of corn leaf aphids, armyworms, cereal leaf beetle, corn rootworm adults, grasshoppers, sap beetle and thrips use 1½ pints MFA Malathion Emulsifiable Concentrate in sufficient water to cover one acre. If aphids are working in leaf sheath, plug nozzles between rows and use as much volume of water as possible . . . directing spray into whorl. Allow a 5 day pre-harvest interval.

PASTURES AND RANGE GRASSES: To control armyworm, use 1 quart in sufficient water to cover one acre. For aphids, grasshoppers and leafhoppers, use 1½ to 2 pints in sufficient water to cover one acre or 1½ pints in one gallon diesel oil per acre.

Note: A minimum of 3 to 10 gallons of water per acre consistent with crop growth and good coverage, is needed for airplane application.

NOTICE: MFA Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to MFA Oil Company, and buyer assumes the risk of any such use.



5 GALLONS

MALATHION E. C.

For Controlling Field Crop Insects and For
Control of Certain Poultry Insects

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

ACTIVE INGREDIENTS:

Malathion*	57%
Aromatic Petroleum Derivative Solvent	35%
INERT INGREDIENTS	8%
Total	100%

*O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

(Contains 5 Pounds Malathion Per Gallon)

WARNING:

FLAMMABLE, KEEP AWAY FROM HEAT AND OPEN FLAME

Manufactured
For



Columbia, Missouri
65201

ACCEPTED WITH COMMENTS