



5 GALLONS

MALATHION E. C.

For Controlling Field Crop Insects and For
Control of Certain Poultry Insects

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

ACTIVE INGREDIENTS:

Malathion*	57%
Aromatic Petroleum Derivative Solvent	35%
INERT INGREDIENTS	8%
Total	100%

* O, O-Dimethyl dithiophosphate of diethyl mercaptosuccinate

(Contains 5 Pounds Malathion Per Gallon)

WARNING:

FLAMMABLE, KEEP AWAY FROM HEAT AND OPEN FLAME

Manufactured
For



Columbia, Missouri
65201

POULTRY APPLICATION FOR CONTROL OF NORTHERN FOWL MITE, CHICKEN RED MITE, AND POULTRY LICE: Mix 2 tablespoons of Malathion per gallon of water. Apply a gallon of mixture per 100 to 150 birds. Repeat in 4 to 8 weeks or as necessary. Also spray infested poultry houses with 4 tablespoons Malathion per gallon of water. Apply liberally to walls, ceilings, roosts, nests and adjacent areas forcing the spray into all cracks and crevices. For poultry tick control increase the dose rate to 1/2 pint per gallon of water. Repeat as necessary. Use a high pressure sprayer for application on surfaces.

COTTON: For control of cotton aphid, boll weevil, cotton leaf perforator, cotton leaf worm and armyworm, use 1 to 2 quarts MFA Malathion Emulsifiable Concentrate per acre diluted with sufficient water for coverage. Apply when temperature is over 60 degrees Fahrenheit and wind is under 10 to 15 miles per hour.

CAUTION: HAZARDOUS IF SWALLOWED. DO NOT BREATHE SPRAY MIST. DO NOT GET IN EYES, ON SKIN OR ON CLOTHING. In case of spillage on skin, wash with soap and water. Do not use on pets, livestock or humans. Do not use in homes, milk houses or other buildings where food is processed.

NOTE TO PHYSICIAN: Malathion is a cholinesterase inhibitor. Atropine and 2-PAM are antidotal.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

Do not reuse empty container. Destroy or decontaminate by crushing. Bury or discard in a safe place.

EPA Reg. No. 100-87

© 1972 MFA OIL

SEPT 12, 72
746-89