



**5%**  
**SEVIN\* DUST**

**An Insecticide for Use on  
Poultry, Pets,  
Vegetables, Small Fruits,  
and Ornamentals**

**CAUTION:**  
**KEEP OUT OF REACH OF CHILDREN**  
**SEE OTHER CAUTIONS  
ON BACK PANEL**

**ACTIVE INGREDIENTS:**

Carbaryl (1-Naphthyl  
N-methylcarbamate)

**INERT INGREDIENTS:**

5%  
95%

**Total 100%**

\*Reg. T.M. Union Carbide Corp. for 1-Naphthyl  
N-methylcarbamate

**DIRECTIONS**

**DOGS AND CATS:** For control of brown dog ticks and fleas; Dust over entire animal, avoiding the eyes. Rub or fluff thoroughly into the skin. Repeat weekly if needed. Also dust sleeping quarters liberally by means of shaker can or hand

**CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS AND PIGEONS:** Direct Dust Application to Birds; Use one pound per 100 birds. Apply by shaker can or hand duster for control of northern fowl mite and lice and as a supplement to premises treatment for chicken mites and fleas. Directly dust each bird and fluff feathers. **DO NOT APPLY WITHIN 7 DAYS OF SLAUGHTER.** Repeat treatment in four weeks if necessary. Direct dusting is a supplement to litter treatment for control of the above insects.

**Litter Treatment;** Apply thoroughly to floors, roosts and interior surfaces, using one pound per 40 square feet to control chicken mites, fleas and bed bugs. Apply by shaker can or hand duster. Do not apply directly to eggs or nest litter. Do not house birds in treated houses within 7 days of slaughter. Do not contaminate feed or drinking water.

**Poultry Dusting Box;** For control of northern fowl mite, chicken mite, lice and fleas, place one self-treatment dust box (18 x 12 x 3 inches deep) containing 2½ pounds 5% Sevin Dust per each 50 birds. Allow birds to dust themselves. This is a supplement to litter treatment for chicken red mite and flea control. Remove birds from dust boxes 7 days before slaughter.

**VEGETABLE AND SMALL FRUITS:** Apply 5% Sevin Dust when insects or their damage appears. Treat both upper and lower surfaces of all foliage. Repeat as necessary, generally at 7 day intervals.

**Beans;** (Green Beans, Lima Beans, Snap Beans, Cowpeas, Black-Eyed Peas) Apply at 20 to 30 pounds per acre to control Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, stink bugs and tarnished plant bug.

**Sweet Corn;** Apply at 20 to 40 pounds per acre to control corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles and leafhoppers.

For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant.

As a silk dust: apply at 2 to 3 day intervals, starting with first silk and continuing until silks begin to dry.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

**Potatoes, Tomatoes, Eggplant, Peppers;** Apply at 20 to 40 pounds per acre to control Colorado potato beetle, leafhoppers, flea beetles, tomato hornworm, tomato fruitworm, stink bugs, lace bugs, and lygus bugs.

**Cucumbers, Melons, Squash, Pumpkin;** Apply at 20 pounds per acre to control pickleworm, melonworm, cucumber beetles, squash bugs, flea beetles and leafhoppers. Some foliage injury may result if treatments are made when tender foliage is wet or with the presence of prolonged high humidity. **DO NOT EXCEED DOSE RATE.**

**Leaf Lettuce, Carrots, Radishes, Mustard Greens and Beets;** Apply at 20 to 40 pounds per acre to control flea beetles, leafhoppers, harlequin bugs, corn earworm, cabbage caterpillar, armyworm, lygus bugs, and stink bugs. Do not use on leaf

**ACTIVE INGREDIENTS:**

**Carbaryl (1-Naphthyl  
N-methylcarbamate)**

5%  
95%

**INERT INGREDIENTS:**

**Total 100%**

\*Reg. T.M. Union Carbide Corp. for 1-Naphthyl  
N-methylcarbamate

**DIRECTIONS**

**DOGS AND CATS:** For control of brown dog ticks and fleas; Dust over entire animal, avoiding the eyes. Rub or fluff thoroughly into the skin. Repeat weekly if needed. Also dust sleeping quarters liberally by means of shaker can or hand duster. Repeat as necessary. Do not treat kittens less than 4 weeks of age.

**ORNAMENTALS:** Roses, Lilac, Junipers, Carnations and Gladiolus; For control of bagworms, blister beetles, boxelder bugs, flea beetles, leafhoppers, mealy bugs, rose aphids, rose slugs, scale insect, leaf rollers, and thrips. Dust thoroughly, covering both upper and lower surfaces of leaves as well as stems and buds; repeat at 7 day intervals, or as needed.

**WARRANTY:** M.F.A. Oil Company warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to M.F.A. Oil Company, and buyer assumes the risk of any such use.

**NET CONTENTS ONE POUND**

Manufactured For



Columbia, Missouri

U.S.D.A. Reg. No. 746-101

G-71

earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetles and leafhoppers.

For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant.

As a silk dust: apply at 2 to 3 day intervals, starting with first silk and continuing until silks begin to dry.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

**Potatoes, Tomatoes, Eggplant, Peppers;** Apply at 20 to 40 pounds per acre to control Colorado potato beetle, leafhoppers, flea beetles, tomato hornworm, tomato fruitworm, stink bugs, lace bugs, and lygus bugs.

**Cucumbers, Melons, Squash, Pumpkin;** Apply at 20 pounds per acre to control pickleworm, melonworm, cucumber beetles, squash bugs, flea beetles and leafhoppers. Some foliage injury may result if treatments are made when tender foliage is wet or with the presence of prolonged high humidity. **DO NOT EXCEED DOSE RATE.**

**Leaf Lettuce, Carrots, Radishes, Mustard Greens and Beets;** Apply at 20 to 40 pounds per acre to control flea beetles, leafhoppers, harlequin bugs, corn earworm, cabbage caterpillar, armyworm, lygus bugs, and stink bugs. Do not use on leaf lettuce or mustard greens within 14 days of harvest, nor on radishes or beets within 3 days of harvest.

**Cabbage, Broccoli and Cauliflower;** Apply at 20 to 40 pounds per acre to control cabbage caterpillars, corn earworm, flea beetles and harlequin bugs. When cabbage looper is the principal problem, use MFA Multi-Spray. For small loopers, after heads form, use 5% Sevin Dust every 5 to 7 days. Do not apply within 3 days of harvest.

**Grapes;** Apply at 40 pounds per acre, or to give plant foliage coverage, to control grape leafhoppers, grape leaf folder, grape berry moth, Japanese beetle, June beetle and red-banded leaf roller.

**Strawberries;** Apply at 20 to 40 pounds per acre to control strawberry weevil, meadow spittlebug and leaf rollers. Do not apply within one day of picking.

**CAUTION:**

**Harmful If Inhaled Or Swallowed:** Do not take internally. Avoid breathing of dust. Avoid skin contact. Wash hands before eating or smoking. Take shower after work. Wear long-sleeved clothing and change to clean clothes daily. Keep out of reach of children and animals. Avoid contamination of foodstuffs, feeds, feeding troughs and watering troughs.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**FOR PROTECTION OF HONEYBEES:** this product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

**DESTROY USED CONTAINER.** Do not reuse empty container. Destroy by burying in a safe place or burning. Keep out of smoke.