


POTATO SPROUT INHIBITOR

columbia/southern
chemicals



Net Contents 1 Gallon

SPROUT NIP

EMULSIFIABLE CONCENTRATE

CONTAINS 3 LBS. CHLORO IPC

ACTIVE INGREDIENT:

Isopropyl N (3-Chlorophenyl) Carbamate

INERT INGREDIENTS:

**CAUTION! HARMFUL IF SWALLOWED
DO NOT TAKE INTERNALLY**


Avoid prolonged breathing of vapor or spray mist.

Avoid contact with skin and eyes.

In case of skin contact, wash with plenty of soap and water and get medical attention.

Keep container closed when not in use.

Avoid prolonged exposure to air to prevent undue moisture pick up.



PITTSBURGH PLATE GLASS COMPANY

CHEMICAL DIVISION

PITTSBURGH 22, PENNSYLVANIA

Manufactured Under U. S. Patent No. 2,695,225

DIRECTIONS FOR USE

NOTICE

Sprout Nip Emulsifiable Concentrate is used for treating potatoes after storage and washing.

Sprout Nip Emulsifiable Concentrate will prevent wound periderm formation of potatoes, therefore, it should be used only after bruises and cuts have healed (normally minimum of two weeks).

Do not apply in field.

Do not use on seed potatoes. This inhibition of sprouting at recommended rates is usually effective up to a year.

Wash equipment thoroughly with soap and water before using for seed potatoes.

EQUIPMENT

Sprout Nip Emulsifiable Concentrate is applied as a 1% emulsion on the drying table after the potatoes have been washed.

Spray nozzle should be set to spray evenly across rolls moving the potatoes.

Dryer rolls will distribute Sprout Nip Emulsifiable Concentrate to assure complete coverage of each potato.

APPLICATION

Dilute Sprout Nip Emulsifiable Concentrate to a 1% emulsion by adding 1 gallon of Sprout Nip Emulsifiable Concentrate to 35 gallons of water.

Agitate enough to form a uniform emulsion.

Apply at rate of 1 quart of 1% emulsion per 20 bags of potatoes (100 pounds per bag).


748-182

1-20-1961

1/2

POTATO SPROUT INHIBITOR

columbia|southern
chemicals



Net Contents 1 Gallon

SPROUT NIP

EMULSIFIABLE CONCENTRATE

CONTAINS 3 LBS. CHLORO IPC

ACTIVE INGREDIENT:

Isopropyl N (3-Chlorophenyl) Carbamate 36.0%

INERT INGREDIENTS: 64.0%

100.0%

**CAUTION! HARMFUL IF SWALLOWED
DO NOT TAKE INTERNALLY**

Avoid prolonged breathing of vapor or spray mist.

Avoid contact with skin and eyes.

In case of skin contact, wash with plenty of soap and water and get medical attention.

Keep container closed when not in use.


Avoid prolonged exposure to air to prevent undue moisture pick up.

PITTSBURGH PLATE GLASS COMPANY

CHEMICAL DIVISION

PITTSBURGH 22, PENNSYLVANIA

Manufactured Under U. S. Patent No. 2,695,225



CONDITIONS OF SALE

Pittsburgh Plate Glass Company guarantees the material sold to be true to the label. Further than that, it makes no representation or warranty, and it shall not be held responsible for any injury to crops, person or property resulting from the use, storage or handling of said material whether in accordance with the directions or not. Nor does Pittsburgh Plate Glass Company guarantee that the residue of the material will not exceed any tolerance established by the appropriate governmental agencies. Application and storage conditions and other factors beyond Pittsburgh Plate Glass Company's control, having much to do with the results, the buyer and/or user accepts and uses this material subject to the above conditions of sale.

Form 23-8 (3-61)

2/6