

POTATO SPROUT INHIBITOR

EMULSIFIABLE CONCENTRATE

SPROUT NIP

ONE GALLON SPROUT
NIP EMULSIFIABLE
CONCENTRATE WILL
TREAT 2,880 CWT OF
POTATOES.

**HAZARD TO
HUMANS AND
DOMESTIC
ANIMALS**

Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Avoid breathing of vapor or spray mist.

**PRACTICAL
TREATMENT**

If swallowed: Induce vomiting by placing fingers in back of throat. Get immediate medical attention. Never give anything by mouth to an unconscious person.
If in eyes: Flush with plenty of water for at least 15 minutes and see a doctor.
If on skin: Wash with plenty of soap and water and see a doctor.

KEEP OUT OF REACH OF CHILDREN

**CAUTION! HARMFUL IF SWALLOWED
DO NOT TAKE INTERNALLY**

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
NET CONTENTS ONE GALLON

Manufactured by
PPG INDUSTRIES, Inc.
One Gateway Center
Pittsburgh, Pa. 15222

EPA REG. NO. 748-182
EPA EST. NO. 748-OH-01
EPA EST. NO. 32761 MO-3

LF 68LB-1A-782D

Biochemicals




INDUSTRIES



FLAMMABLE LIQUID


3

**FLAMMABLE LIQUID, N.O.S.
UN1993**



DIRECTIONS FOR USE

Violation of Federal law to use this product in a manner inconsistent with the label



CAUTION! ENVIRONMENTAL HAZARD

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTICE

Sprout Nip Emulsifiable Concentrate is used for treating potatoes after storage and washing. Sprout Nip Emulsifiable Concentrate will prevent wound periderm formation of potatoes, therefore, it should be used only after bruises and cuts have healed. (normally minimum of two weeks).

Do Not apply in field.

Do not use on seed potatoes. This inhibition of sprouting at recommended rates is usually effective up to a year. Wash equipment thoroughly with soap and water before using for seed potatoes.

EQUIPMENT

Sprout Nip Emulsifiable Concentrate is applied as a 1% emulsion on the drying table after the potatoes have been washed. Spray nozzle should be set to spray evenly across rolls moving the potatoes.

Dryer rolls will distribute Sprout Nip Emulsifiable Concentrate to assure complete coverage of each potato.

APPLICATION

Dilute Sprout Nip Emulsifiable Concentrate to a 1% emulsion by adding 1 gallon of Sprout Nip Emulsifiable Concentrate to 5 gallons of water. Agitate enough to form a uniform emulsion.

Apply at rate of 1 quart of 1% emulsion per 20 bags of potatoes (100 pounds per bag).

USE 1A 767D

STORAGE AND DISPOSAL

Keep container closed. Do not contaminate water, food, or feed by storage or disposal. This product inhibits germination of seed potatoes.

Avoid storage below freezing temperatures. Prolonged storage may cause crystal formation. Warming contents and agitation will restore material to usable condition, but do not heat above 250 F.

Any Sprout Nip or rinse water that can not be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

Do not reuse container. Triple-rinse then puncture and dispose of in a sanitary landfill or by other approved state and local procedures.

CONDITIONS OF SALE

PPG Industries, Inc. the seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.