

ACCE

6/11/55

## CHEM HOE 15-G GENERAL DIRECTIONS FOR USE

NOTICE: Use of this product in the manner or at a time or in quantities or on crops other than in accordance with the directions may produce serious plant injury.

### GENERAL

The following general directions are supplied for information purposes only. They are believed to be reliable but are in no way guaranteed. While these directions pertain to application under average conditions, ineffective weed control or crop damage may result from differing conditions of soil, temperature, moisture and crop variety. Therefore, before large scale use, this product should be applied on a trial basis under conditions actually encountered. Consult Agricultural Experiment Station Weed Specialists or Agricultural Extension Service Specialists for detailed recommendations of application under varying local conditions.

### WEEDS CONTROLLED

CHEM HOE 15-G controls many annual grasses and certain annual dicotyledonous weeds as they germinate. Used according to directions, it will normally give effective control of germinating seeds and young seedlings of the following weeds: Volunteer Small Grains, Annual Bluegrass, Annual Ryegrass, Foxtail Barkley, Fallbroomgrass, Chickweed, Small Nettle, Downy Broom (Chenopod), Wild Oats, Rattail Fescue and Corngrass.

In certain areas is effective post-emergence control of the above seedling weeds and may need to be used. No control of most weeds will be obtained normally. Do not expect results from treatments made after the weeds

This 70 pound bag of CHEM HOE 15-G contains 7.5 pounds of Isopropyl-Naphthyl Sulfonate.

### CONTAINER STORAGE AND HANDLING

Use in combination with fertilizers, lime, manure, and other soil amendments. Do not use in combination with herbicides. Do not use in combination with insecticides. Do not use in combination with fungicides.

### CONDITIONS OF SALE

THIS PRODUCT IS SOLD AS A FERTILIZER. IT CONFORMS TO THE REQUIREMENTS OF THE FERTILIZER ACT AND REGULATIONS THEREUNDER.

## DIRECTIONS FOR SPECIFIC APPLICATIONS

Rates given are for over all coverage except as otherwise indicated. For strip or band treatment reduce rates proportionately.

**ALFALFA, LINDO CLOVER, WHITE CLOVER, RED CLOVER or CRISON CLOVER.** Post-emergence. For control of certain annual weeds, apply 1.5 to 2.0 pounds of granules per acre. Make application in early morning or late afternoon. Do not apply if rain is expected within 24 hours. Do not apply if rain is expected within 24 hours after application. Do not apply if rain is expected within 24 hours after application. Do not apply if rain is expected within 24 hours after application.

**FLAX.** Post-emergence. For control of certain annual weeds, apply

1.5 to 2.0 pounds of granules per acre. Make application in early morning or late afternoon. Do not apply if rain is expected within 24 hours. Do not apply if rain is expected within 24 hours after application. Do not apply if rain is expected within 24 hours after application.

### APPLICATION

Application may be made by airplane, helicopter, fertilizer spreader or cyclone seeder. Apply the dry granules as they come from the package. Uniformity of coverage is essential. Before applying CHEM HOE 15-G to a crop area, check your distributor's equipment with this specific product. This can be done over the paper of the package, or a dark soil area. Do not assume a similar distribution pattern to another granular product which may vary in density, particle size, or bulk weight. An uneven distribution pattern with resulting high and low rates of application may cause crop injury or failure to control weeds.

### TEMPERATURE and MOISTURE CONSIDERATIONS

CHEM HOE 15-G breaks down rapidly in warm, moist soil, and therefore has proven most effective when applied during cool weather. One site of entry into plants is through the roots. Entry and activity depend on sufficient moisture to carry the chemical to the root zone. CHEM HOE 15-G also acts on germinating seeds and young seedlings. Therefore, moisture is also needed to cause weed seed germination at a time when maximum CHEM HOE 15-G effectiveness is present. If rainfall does not occur within five days after application, the field should be irrigated.

In its chemical description and is not suitable for the purpose of fertilizing. It is not suitable for use in combination with other fertilizers. It is not suitable for use in combination with other fertilizers. It is not suitable for use in combination with other fertilizers. It is not suitable for use in combination with other fertilizers.

For control of certain annual weeds, apply 1.5 to 2.0 pounds of granules per acre. Make application in early morning or late afternoon. Do not apply if rain is expected within 24 hours. Do not apply if rain is expected within 24 hours after application. Do not apply if rain is expected within 24 hours after application.

For control of certain annual weeds, apply 1.5 to 2.0 pounds of granules per acre. Make application in early morning or late afternoon. Do not apply if rain is expected within 24 hours. Do not apply if rain is expected within 24 hours after application. Do not apply if rain is expected within 24 hours after application.

USDA REG. No. 748-211



PPG INDUSTRIES, Inc.  
Chemical Division  
One Gateway Center  
Pittsburgh, Pa. 15222

Stamp on back  
of last  
page.

15 18 pt type

12 pt. type

14

10