

SECURITY

# PAN PEACH SPRAY

## WITHOUT LEAD ARSENATE

### CONTAINS BASIC ZINC SULPHATE AND SULPHUR

**ACCEPTED**

JAN 6 - 1969

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 769-29

**ACTIVE INGREDIENTS:**

Zinc, as metallic equivalent to 55% Basic Sulphate of Zinc.

as  $ZnSO_4 \cdot Zn(OH)_2$  .....

Sulphur, as elemental .....

**INERT INGREDIENTS:** .....

U. S. D. A. Reg. No. 769-29

### CAUTION!

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed. Avoid contamination of food

and clothing. Avoid spraying this material during very hot periods of the day.

### DIRECTIONS

**PEACHES:** 1. Apply 1.5 to 3.0 lbs. per acre and 2. Apply 1.5 to 3.0 lbs. per acre at the time of bloom. This material is effective against peach leaf curl, peach scab, and peach powdery mildew. It also acts as a fertilizer. Do not apply this product within 14 days of harvest. Do not apply this product within 14 days of harvest.

Have been killed and before the appearance of the first generation (the 2nd spray after check off) and again 10 days or 2 weeks before harvest. Do not apply within 14 days of harvest.

**DISPOSAL WARNING:** Dispose of waste properly and do not bury in non-crop land away from water supplies. Do not re-use container.

**NOTICE - DISCLAIMER OF WARRANTY, MERCHANTABILITY AND FITNESS:** If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then we disclaim all warranties, expressed or implied, including the warranty of merchantability.

## WOOLFOLK CHEMICAL WORKS, LTD.

MANUFACTURERS

760369

FORT VALLEY, GEORGIA