

SECURITY

ACCEPTED

Jul 29, 1971

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND FERTILIZER ACT
REGARDING POISON RESIDUES

USE NO. 769-32
with comment

PAN PEACH SPRAY

ACTIVE INGREDIENTS:

Zinc as metallic equivalent to 19.1% Basic Sulfate of Zinc as $ZnSO_4 \cdot Zn(OH)_2$	not less than	9.6%
Sulphur as elemental	not less than	34.0%
Lead Arsenate as $PbHAsO_4$	not less than	11.5%
INERT INGREDIENTS:	not more than	44.9%
	TOTAL	100.0%

U. S. D. A. Reg. No. 769-32

Total Arsenic as metallic not less than 2.1%

Arsenic in water soluble form as metallic not more than 0.2%

Lead, as metallic, 6.6%

Reg. U. S. Pat. Off.

U. S. Patent No. 1943181

WARNING: KEEP OUT OF REACH OF CHILDREN. May be Fatal if Swallowed! Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid breathing dust or spray mist. Keep away from children, pets, livestock and feedstuffs.

guaranteed. The number and time of applications will vary in different localities. Fruiting varieties will not require the full schedule. Consult your nearest State Agricultural Experiment Station or Extension Service for specific directions.

SPRAY SCHEDULE

PEACHES: (1) When 75% of petals (ink part of flower) have fallen; (2) When calyxes, or "shucks" are shedding, or when small fruit is exposed; (3) Two weeks after No. 1; (4) Two weeks after No. 3 (See Note Below); (5) 4 weeks before harvest; (6) 10 days before harvest (See Note Below).

NOTE: Use Pan Without Lead Arsenate or, if no zinc is desired, Nutonex wettable sulphur in spray No. 4 and 6 of the peach schedule, then three sprays at weekly intervals. Pan Peach Spray containing Lead Arsenate should not be used within 30 days of harvest.

CAUTION: Do not apply within 30 days of harvest. Any residue remaining at harvest must be removed by washing, brushing or other effective means. Do not spray when trees are wet or when temperature is above 90° F. Do not graze livestock on treated areas.

Dispose of waste pesticide and container by burying in non-crop land away from water supplies. Never re-use container.

ANTIDOTE: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give 2 tablespoonfuls of Epsom Salts or Milk of Magnesia in water, and plenty of milk and water. Have victim lie down and keep quiet. Call a Physician immediately.

This product is highly toxic to fish and wildlife. Keep out of creeks, ponds, and streams. Do not apply in any manner not specified on the label.

This product is highly toxic to trees. Avoid direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS

For control of Plum Curculio, Brown Rot, Scab, and Bacterial Spot use at the rate of 16 pounds per 100 gallons of water. The following schedule is suggested as a guide but is in no way

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or distributed in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. The liability for sale, use, storage, transport or distribution of this product in accordance with the instructions set forth on the label shall then and thereafter terminate and end Woolfolk Chemical Works, Ltd.'s liability for injury to crops, animals, personal property and real property of every kind and description and for injury to any person or persons and for any consequential damages arising out of or from the foregoing set forth. There are no warranties which extend beyond the words on this label.

WOOLFOLK CHEMICAL WORKS, LTD.
MANUFACTURERS 070216 **FORT VALLEY, GEORGIA**