

ACCEPTED

MAY 21 1981

SECURITY

769-34

WETTABLE SULPHUR

ACTIVE INGREDIENT:

Sulphur, as elemental not less than 90.0%

INERT INGREDIENTS: 10.0%

Sulphur - 95% Through 325 Mesh Screen TOTAL 100.0%
EPA Reg. No. 769-34 EPA. Est. 769-AL-1 EPA Est 769-GA-1

KEEP OUT OF REACH OF CHILDREN.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Avoid contamination of feed and foodstuffs.

FIRST AID: IF IN EYES - immediately flush with plenty of water for at least 15 minutes. Call a physician if irritation persists. FOR SKIN CONTACT - wash with soap and water. IF INHALED - and overcome by vapor or dust remove immediately from exposure and call a physician. Administer artificial respiration if breathing is irregular or has stopped. IF SWALLOWED - Drink 1 or 2 glasses of water and induce vomiting by sticking finger in back of throat. Never give anything by mouth to an unconscious person. Call a physician.

ENVIRONMENTAL HAZARDS

Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. **NOTE TO USER:** Do not use this product as a feed supplement. Do not apply this material to Cucumbers, Melons or Squash. Avoid applying this material during very hot periods of the day. When this product is used with Lead Arsenate, 8 pounds of Hydrated Lime or Basic Zinc Sulphate should be added for each 10 gallons of spray.

PEACHES and PLUMS: To control Brown Rot and Scab use 6 pounds per 100 gallons of water. Follow locally issued spray schedule for number and frequency of applications. Usually 6 applications are made. No. 1 — when 2/3 petals have fallen, No. 2 — when 3/4

shucks have shed, No. 3 — two weeks later, No. 4 — two weeks after No. 3, No. 5 — four weeks before harvest, No. 6 — two weeks before harvest.

FLOWERS and ORNAMENTALS: To control Red Spiders and the diseases listed below use Nutonex as a dust or spray using 6 pounds per 100 gallons of water. Applications should be made at 5 to 7 day intervals beginning when the disease first appears and continuing until all danger from the disease has passed. During periods of humid or rainy weather it may be necessary to make applications as often as every 2 to 3 days.

POWDERY MILDEW: Aster, Chrysanthemum, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden Fleece, Hibiscus, Honeysuckle, Hydrangea, Lady's Mantle, Ligustrum, Lilac, Myrtle, Phlox, Poplar, Silver Vine, Rose, Spiraea, Sunflower, Verbena, Willow and Zinnia.

LEAF SPOTS: Calendula, Cherry Laurel, Clematis, Columbine, English Ivy, Foxgloves, Hydrangea, Petunia, Phlox, Sage, Smilax and Snapdragon.

BLACK SPOT: Rose.

ROSE POWDERY MILDEW: Rose.

BOTRYTIS BLIGHT: Hydrangea.

RED SPIDERS: Arbovitae, Carnation, Cedars, Chrysanthemum, Clematis, Hydrangea, Rose, Snapdragon and Sweetpeas.

STORAGE AND DISPOSAL

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

NOTICE: DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY

If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Inc. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness. 182130

WOOLFOLK CHEMICAL WORKS, INC. 67
MANUFACTURERS FORT VALLEY, GEORGIA

769-34

05/21/1981

1/1