

ACCEPTED  
Oct. 20, 1967  
UNDER THE FEDERAL INSECTICIDE  
ACT

769-58

### DIRECTIONS

**APPLES:** To control scab use a delayed dormant application, when buds begin to break and show green, consisting of 5 gallons of Lime Sulphur per 100 gallons of water (2½ cupfuls to 3 gallons of water). A second application using 5 quarts to 100 gallons of water (¾ cupful to 3 gallons) should be made when buds are in the prepink stage before blossoms open. A third application using 5 quarts per 100 gallons of water should be made after blossoms fall.

**PEACHES:** Blossom Blight (Brown Rot) - Use 1 gallon per 100 gallons of water (½ cupful to 3 gallons of water) making the first application when blossoms first begin to open. Two or three additional applications at 3 day intervals should be made depending upon the length of the bloom period.

**DISPOSAL WARNING:** Never re-use container. Dispose of empty container in a safe place.

#### NOTICE

**Disclaimer of Warranty, Merchantability and Liability** - If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter Woolfolk Chemical Works, Ltd. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

SECURITY  
BRAND

## LIME SULPHUR SOLUTION 32° BÉ

#### ACTIVE INGREDIENT:

Calcium Polysulphide ..... 30.0%

INERT INGREDIENTS: . 70.0%

TOTAL 100.0%

**CAUTION: KEEP OUT OF REACH OF CHILDREN.** May cause irritation of skin. Avoid contact with skin. Wash thoroughly after handling.

Do not use this product on apples within 2 weeks of an oil spray. Lime Sulphur may burn foliage when temperature is high; do not apply at such times. In the South do not use on stone fruit during the growing period. Do not apply or allow to drift to painted surfaces or a permanent stain may result. U.S.D.A. Reg. No. 769-58

**WOOLFOLK CHEMICAL WORKS, LTD.**

MANUFACTURERS - FORT VALLEY, GA.

NET CONTENTS - ONE QUART

### DIRECTIONS

**DORMANT APPLICATIONS:** Against scale insects attacking deciduous fruit trees, ornamental shrubbery and berries, dilute with water at the rate of 1 part Lime Sulphur to 8 parts of water. Spray during the dormant season before growth starts in the spring. Do not apply dormant spray to peaches, if a sufficient number of dormant hours have occurred, immediately after or during periods of 5 days or more of unseasonably high temperatures.

#### **CITRUS, EVERGREENS AND BERRIES:**

To control red-spiders and rust mites use 1 gal. of Lime Sulphur plus 5 to 10 lbs. of Nutonex Sulphur per 100 gals. (½ cupful of Lime Sulphur plus ½ to 1 cupful of Nutonex Sulphur to 3 gals. water). Lime Sulphur should not be used on early and mid-season oranges or on tangerines in late winter or early spring.

**ROSES:** As a winter clean up and aid in control of diseases and for control of **Rose Scale, Oyster-Snail Scale** and **San Jose Scale**, use 1 part Lime Sulphur to 8 parts of water (1 pint to 1 gallon) and apply while plants are still dormant. To aid in control of Powdery Mildew use 1 part to 15 parts of water (1 pint to 2 gallons) and apply in early spring, immediately after pruning, while the plants are still dormant. 761002